

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/00, C07D 205/08, 223/16, 239/02, 211/26, 285/08, 285/14, 417/00, 513/00, 277/04, 277/18, 277/38, 263/04, 263/16, 295/00, 207/00, 207/08, 207/10, C07C 229/00, 241/00, 243/00, 249/00, 233/00, 235/00, 237/00, 239/00	A1	(11) International Publication Number: WO 95/23609 (43) International Publication Date: 8 September 1995 (08.09.95)
(21) International Application Number: PCT/US95/02558 (22) International Filing Date: 3 March 1995 (03.03.95) (30) Priority Data: 08/206,553 4 March 1994 (04.03.94) US (71) Applicant: ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). (72) Inventors: SMITH, Gerald, Floyd; 825 Queenswood Court, In- dianapolis, IN 46217 (US). WILEY, Michael, Robert; 7725 Langwood Drive, Indianapolis, IN 46268 (US). SCHACHT, Aaron, Leigh; 8061 Mallard Landing, Indianapolis, IN 46278 (US). SHUMAN, Robert, Theodore; 2596 Willow Street, Greenwood, IN 46143 (US). (74) Agents: LAMMERT, Steven, R. et al.; Barnes & Thornburg, 1313 Merchants Bank Building, 11 South Meridian Street, Indianapolis, IN 46204 (US).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LV, MD, MG, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, TJ, TT, UA, UG, UZ, VN, OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i>
(54) Title: ANTITHROMBOTIC AGENTS (57) Abstract This invention relates to thrombin inhibiting compounds having the formula (I): X-Y-NH-(CH ₂) _r -G where X, Y, r and G have the values defined in the description, as well as pharmaceutical formulations containing those compounds and methods of their use as thrombin inhibitors, coagulation inhibitors, and thromboembolic disorder agents.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

ANTITHROMBOTIC AGENTS

This invention relates to thrombin inhibitors which are useful anticoagulants in mammals. In particular it
10 relates to peptide derivatives having high antithrombotic activity, anticoagulant activity, and oral bioavailability.

The process of blood coagulation, thrombosis, is triggered by a complex proteolytic cascade leading to the formation of thrombin. Thrombin proteolytically removes
15 activation peptides from the A α -chains and B β -chains of fibrinogen, which is soluble in blood plasma, initiating insoluble fibrin formation.

Anticoagulation is currently achieved by the administration of heparins and coumarins.

20 Parenteral pharmacological control of coagulation and thrombosis is based on inhibition of thrombin through the use of heparins. Heparins act indirectly on thrombin by accelerating the inhibitory effects of endogenous antithrombin III (the main physiological inhibitor of
25 thrombin). Because antithrombin III levels vary in plasma and because surface-bound thrombin seems resistant to this indirect mechanism, heparins can be an ineffective treatment. Because coagulation assays are believed to be associated with efficacy and with safety, heparin levels must be monitored
30 with coagulation assays (particularly the activated partial thromboplastin time (APTT) assay). Coumarins impede the generation of thrombin by blocking the posttranslational

gamma-carboxylation in the synthesis of prothrombin and other proteins of this type. Because of their mechanism of action, the effect of coumarins can only develop slowly, 6-24 hours after administration. Further, they are not selective
5 anticoagulants. Coumarins also require monitoring with coagulation assays (particularly with the prothrombin time (PT) assay).

Recently, interest in small synthetic peptides that are recognized by proteolytic enzymes in a manner similar to
10 that of natural substrates has grown. Tripeptide aldehydes such as D-Phe-Pro-Arg-H, Boc-D-Phe-Pro-Arg-H, and D-MePhe-Pro-Arg-H, Bajusz et al., J. Med. Chem., 33, 1729-1735 (1990) demonstrate potent direct inhibition of thrombin. Early clinical studies which demonstrate that D-MePhe-Pro-Arg-H
15 sulfate is an anticoagulant in man have been reported, see Simoons et al., Circulation, 90, I-231, Abstr. 1241 (1994). Many investigators have synthesized analogs in an effort to develop pharmaceutical agents, for example Shuman et al., J. Med. Chem., 36, 314-319 (1993). U.S. Patent No. 4,346,078
20 teaches a series of anti-coagulant peptides containing an agmatine (1-amino-4-guanidinobutane) group. Agmatine derivatives and related compounds also are disclosed in the PCT application with International Publication Number WO 93/11152, as well as in European Patent Application,
25 Publication number 601459, published 15 June 1994. Such compounds differ from the former series in that the agmatine compounds lack a carbonyl moiety found in similar compounds containing an Arg group.


Although the heparins and coumarins are effective
30 anticoagulants, and no drug has yet emerged from the known tripeptide aldehydes, and despite the continuing promise for this class of compounds, there exists a need for anticoagulants that act selectively on thrombin, and independent of antithrombin III, exert inhibitory action
35 shortly after administration, preferably by an oral route, and do not interfere with lysis of blood clots, as required to maintain hemostasis.

The present invention is directed to the discovery that the compounds of the present invention, as defined below, are potent thrombin inhibitors that may have high bioavailability following oral administration. In addition, certain compounds of the present invention also may exhibit inhibition of factor Xa, which is involved in the coagulation cascade.

Accordingly, it is a primary object of the present invention to provide novel peptide derivatives that are potent thrombin inhibitors useful as anticoagulants.


Other objects, features, and advantages will be apparent to those skilled in the art from the following description and claims.

The present invention provides a thrombin inhibiting compound having the Formula I



wherein

X is prolinyl, homoprolinyl,
 $R^m-(CH_2)_g-NH-CH_2-C(O)-$,




in which

R^d is carboxy or methylsulfonyl;



R^e is $NHRC^c$, $NHCOR^c$ or $NHCOOR^c$; in which
 R^c is C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl or a (C_3 - C_8)
cycloalkyl (C_1 - C_6) alkyl radical of 4-10 carbons;
 T is C_3 - C_8 cycloalkyl, C_1 - C_8 alkyl,

5



- a is 0, 1 or 2; and
 Q is $-OH$, C_1 - C_4 alkoxy, or $-NH-A$;
 A is hydrogen, C_1 - C_4 alkyl, $R''SO_2-$, $R''OC(O)-$,
10 $R''C(O)-$, $R''C(O)-$ or $-(CH_2)_g-R^m$;
 g is 1, 2, or 3;
 B is hydrogen or C_1 - C_4 alkyl;
 R' is hydrogen or C_1 - C_4 alkyl;
 R'' is C_1 - C_4 alkyl, C_1 - C_4 perfluoroalkyl, $-(CH_2)_d-R^m$,
15 or unsubstituted or substituted aryl, where aryl is phenyl,
naphthyl, a 5- or 6-membered unsubstituted or substituted
aromatic heterocyclic ring, having one or two heteroatoms
which are the same or different and which are selected from
sulfur, oxygen and nitrogen, or a 9- or 10-membered
20 unsubstituted or substituted fused bicyclic aromatic
heterocyclic group having one or two heteroatoms which are
the same or different and which are selected from sulfur,
oxygen and nitrogen;
 R^m is $-COOR^b$, $-SO_2(C_1-C_4 \text{ alkyl})$, $-SO_3H$, $-P(O)(OR^b)_2$
25 or tetrazol-5-yl;
 R^n is $-COOR^b$ or tetrazol-5-yl;
each R^b is independently hydrogen or C_1 - C_4 alkyl;
 d is 1, 2, or 3;
 m is 0, 1, or 2;
30 n is 0, 1, or 2; and
 Z is hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, hydroxy,
halo or R_aSO_2NH- , where R_a is C_1 - C_4 alkyl;

5



in which

R^g is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, or $-(\text{CH}_2)_p\text{-L-}(\text{CH}_2)_q\text{-T}'$;

5 R^p is hydrogen, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, or $-(\text{CH}_2)_p\text{-L-}(\text{CH}_2)_q\text{-T}'$;

where p is 0, 1, 2, 3, or 4; L is a bond, $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$; q is 0, 1, 2 or 3; and T' is hydrogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, $-\text{COOH}$, $-\text{CONH}_2$, or Ar , where Ar is

10 unsubstituted or substituted aryl, where aryl is phenyl, naphthyl, a 5- or 6-membered unsubstituted or substituted aromatic heterocyclic ring, having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen, or a 9- or 10-membered
15 unsubstituted or substituted fused bicyclic aromatic heterocyclic group having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen;

R^y is $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$; and



20 R^z is a bond or, when taken with R^y and the three adjoining carbon atoms, forms a saturated carbocyclic ring of 5-8 atoms, one atom of which may be $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$;

r is 1, 2 or 3; and



G is $-(\text{CH}_2)_s\text{-R}$, where s is 0-5, $-\text{CH}=\text{CH}-(\text{CH}_2)_t\text{-R}$,

25 where t is 0-3, or G is

6



5



10

where D and E are each independently N or CH;



k is 0 or 1;

b is 0 or 1;

M is S, O, or NH;

each W is independently N or CH; and


15



20

and wherein one to all of the otherwise unsubstituted carbon atoms of the aromatic or heteroaromatic rings of

7




may bear a fluoro substituent;

or a pharmaceutically acceptable salt thereof;

5 or a pharmaceutically acceptable solvate of said compound or salt thereof;


provided that A is not hydrogen or t-butyloxycarbonyl when G is $-(CH_2)_s-NH-C(NH)NH_2$, Y is




unsubstituted prolinyl (R^D is hydrogen), and T is

10 and provided further that R is not amino or guanidino when $r=1$ and $s=0$;

and provided further that A is not hydrogen, C_1 - C_4 alkyl, methylsulfonyl or $-(CH_2)_g-R^m$ when G is



15

Y is  , unsubstituted prolinyl (R^D is hydrogen) or 4-hydroxyprolinyl (R^D is OH), R' is hydrogen, T is cyclohexyl and Q is $-NH-A$;


and provided further that $R''SO_2$ is not arylsulfonyl

20 when G is $-(CH_2)_s-R$ in which R is $-NH-C(=NH)-NH_2$, Y is

unsubstituted prolinyl (R^D is hydrogen) or 4-methylthioprolinyl (R^D is $-SCH_3$), and Q is $-NH-A$;


and provided further that A is not $R''SO_2-$ when G is

8




A particular group of the above compounds of
Formula I consists of those compounds of Formula I
5 wherein

X is prolinyl, homoprolinyl,



T is C₃-C₈ cycloalkyl, C₁-C₈ alkyl,

10



a is 0 or 1;

Q is -OH, C₁-C₄ alkoxy, or -NH-A;

A is hydrogen, C₁-C₄ alkyl, R"SO₂-, R"OC(O)-,

15 R"C(O)-, or -(CH₂)_g-COOH;

g is 1, 2, or 3;

B is hydrogen or C₁-C₄ alkyl;

R' is hydrogen or C₁-C₄ alkyl;

R" is C₁-C₄ alkyl, C₁-C₄ perfluoroalkyl,

20 -(CH₂)_d-COOH, or unsubstituted or substituted aryl, where aryl


is phenyl, naphthyl, a 5- or 6-membered unsubstituted or substituted aromatic heterocyclic ring, having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen, or a 9- or 10-membered unsubstituted or substituted fused bicyclic aromatic heterocyclic group having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen;

d is 1, 2, or 3;

10 m is 0, 1, or 2;

n is 0, 1, or 2; and

Z is hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, halo, or R_aSO₂NH-, where R_a is C₁-C₄ alkyl;



in which

R^g is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or $\text{--(CH}_2\text{)}_p\text{--L--(CH}_2\text{)}_q\text{--T'}$;

15 R^p is hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or $\text{--(CH}_2\text{)}_p\text{--L--(CH}_2\text{)}_q\text{--T'}$;

20 where p is 0, 1, 2, 3, or 4; L is a bond, -O-, -S-, or -NH-; q is 0, 1, 2 or 3, and T' is hydrogen, C₁-C₄ alkyl, C₃-C₈ cycloalkyl, -COOH, -CONH₂, or Ar, where Ar is

unsubstituted or substituted aryl, where aryl is phenyl, naphthyl, a 5- or 6-membered unsubstituted or substituted aromatic heterocyclic ring, having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen, or a 9- or 10-membered unsubstituted or substituted fused bicyclic aromatic


heterocyclic group having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen;

R^Y is $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$; and

5 R^Z is a bond or, when taken with R^Y and the three adjoining carbon atoms, forms a saturated carbocyclic ring of 5-8 atoms, one atom of which may be $-\text{O}-$, $-\text{S}-$, or $-\text{NH}-$;

r is 1 or 2; and

G is $-(\text{CH}_2)_s-\text{R}$, where s is 0-5, $-\text{CH}=\text{CH}-(\text{CH}_2)_t-\text{R}$,



where D and E are each independently N or CH ;

k is 0 or 1;

b is 0 or 1;

15 M is S , O , or NH ;

each W is independently N or CH ; and

R is $-\text{NH}_2$, $-\text{C}(\text{NH})\text{NH}_2$, $-\text{C}(\text{NOH})\text{NH}_2$, or $-\text{NH}-\text{C}(\text{NH})\text{NH}_2$;


or a pharmaceutically acceptable salt thereof;

or a pharmaceutically acceptable solvate of said compound

20 or salt thereof;

provided that A is not hydrogen or

t -butyloxycarbonyl when G is $-(\text{CH}_2)_s-\text{NH}-\text{C}(\text{NH})\text{NH}_2$, Y is



unsubstituted prolinyl (R^P is hydrogen), and T is

and provided further that R is not amino or

25 guanidino when $r=1$ and $s=0$;

and provided further that A is not hydrogen, C₁-C₄ alkyl, methylsulfonyl or -(CH₂)_g-COOH when G is -(CH₂)₅-R in

which R is $\begin{array}{c} \text{NH} \\ \parallel \\ -\text{C}-\text{NH}_2 \end{array}$ or $\begin{array}{c} \text{NH} \\ \parallel \\ -\text{NH}-\text{C}-\text{NH}_2 \end{array}$,



Y is $\begin{array}{c} \text{O} \\ \parallel \\ \text{---N---}^*\text{---C---} \end{array}$, unsubstituted prolinyl

5 (R^P is hydrogen) or 4-hydroxyprolinyl (R^P is OH), R' is hydrogen, T is cyclohexyl and Q is -NH-A;

and provided further that RⁿSO₂ is not arylsulfonyl when

G is -(CH₂)₅-R in which R is $\begin{array}{c} \text{NH} \\ \parallel \\ -\text{NH}-\text{C}-\text{NH}_2 \end{array}$, Y is unsubstituted prolinyl (R^P is hydrogen) or 4-methyl-thioprolinyl (R^P is -SCH₃), and Q is -NH-A;

and provided further that A is not RⁿSO₂- when G is



In addition to the compounds of Formula I, the present invention provides pharmaceutical formulations comprising a compound of Formula I in association with a pharmaceutically acceptable carrier, diluent or excipient.

The present invention also provides a method of inhibiting thrombosis in mammals comprising administering to a mammal in need of treatment, an antithrombotic dose of a compound of Formula I.

The present invention further provides a method of inhibiting thrombin comprising administering to a mammal in need of treatment, a thrombin inhibiting dose of a compound of Formula I.

This invention relates to new inhibitors of thrombin, pharmaceutical compositions containing the compounds as active ingredients, and the use of the compounds as anticoagulants for prophylaxis and treatment of thromboembolic diseases such as venous thrombosis, pulmonary embolism, arterial thrombosis, in particular myocardial ischemia, myocardial infarction and cerebral thrombosis, general hypercoagulable states and local hypercoagulable states, such as following angioplasty and coronary bypass operations, and generalized tissue injury as it relates to the inflammatory process.

The term "alkyl" by itself or as part of another substituent means a straight or branched chain alkyl radical having the stated number of carbon atoms such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl and sec-butyl. The term "perfluoroalkyl" by itself or as part of another substituent means a straight or branched chain alkyl radical having the stated number of carbon atoms in which each hydrogen atom is replaced with a fluorine atom such as trifluoromethyl, perfluoroethyl, perfluoro-n-propyl, perfluoroisopropyl, perfluoro-n-butyl, perfluoro-t-butyl, perfluoroisobutyl and perfluoro-sec-butyl.

The term "C₃-C₈ cycloalkyl" refers to the saturated alicyclic rings of three to eight carbon atoms such as cyclopropyl, methylcyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 4-methylcyclohexyl, cyclooctyl, and the like.

The term "alkoxy" means a straight or branched chain alkyl radical having the stated number of carbon atoms bonded to the parent moiety by an oxygen atom. The term "halo" means chloro, fluoro, bromo or iodo. The term "acetyl" means CH₃-C(O)-. The term "t-butyloxycarbonyl" means (CH₃)₃C-O-C(O)- and is abbreviated "Boc". The term "benzyloxycarbonyl" means C₆H₅CH₂-O-C(O)- and is abbreviated "Cbz".

The term "5- or 6-membered heterocyclic ring" means any 5- or 6-membered ring that will afford a stable structure containing one or two nitrogen atoms; one sulfur atom; one

oxygen atom; one nitrogen and one sulfur atom; or one nitrogen and one oxygen atom. The 5-membered ring has one or two double bonds and the 6-membered ring has two or three double bonds. Such heterocyclic systems include furyl, thienyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyranyl, pyridinyl, pyrimidinyl, pyrazinyl, oxazinyl and thiazinyl.


The term "9- or 10-membered heterocyclic ring" means any bicyclic group in which any of the above 5- or 6-membered rings is fused to a benzene ring or another 6-membered heterocyclic ring as defined above that will afford a stable structure. These heterocyclic systems include indolyl, benzothienyl, benzofuryl, benzoxazolyl, benzoisoxazolyl, benzopyrazolyl, quinolinyl, isoquinolinyl, benzimidazolyl and benzothiazolyl.

It will be appreciated that many of the above heterocycles may exist in tautomeric forms. All such forms are included within the scope of this invention.

All of the aromatic or heteroaromatic groups listed for the definition of Ar or R" are independently unsubstituted or substituted with one or two substituents that will afford a stable structure independently selected from halo, hydroxyl, C₁-C₄ alkyl, C₁-C₄ alkoxy, amino (-NH₂), mono(C₁-C₄ alkyl)amino, -(CH₂)_jCOOH, mercapto, -S(O)_h(C₁-C₄ alkyl), -NHS(O)_h(C₁-C₄ alkyl), -NHC(O)(C₁-C₄ alkyl), -S(O)_hNH₂, -S(O)_hNH(C₁-C₄ alkyl), or -S(O)_hN(C₁-C₄ alkyl)₂, h is 0, 1 or 2, and j is 0, 1, 2, 3, or 4. One particularly preferred such value for the substituent R"(C)O- is 1-methylindol-2-oyl.


In the representation of Formula I, the carbonyl functionality of group X is attached to the amine functionality of the Y group. The carbonyl functionality of Y is then attached to the amino group drawn in Formula I.

14





The group Z , where Z and A are both hydrogen, is referred to at times herein as phenylglycyl and abbreviated Phg. Compounds wherein A is, e.g., methyl, are referred to as the N^α -methyl-phenylglycyl group and

- 5 abbreviated MePhg. Substituted compounds wherein Z is other than hydrogen are referred to by the type and position of the substituent group, e.g., 3'-chlorophenylglycyl or Phg(3-Cl).




- The group Z , where Z and A are both hydrogen, is referred to at times herein as phenylalanyl and abbreviated Phe. Compounds wherein A is, e.g., methyl, are referred to as the N^α -methyl-phenylalanyl group and abbreviated MePhe. Substituted compounds wherein Z is other than hydrogen are referred to by the type and position of the substituent group, e.g., 3'-chlorophenylalanyl or Phe(3-Cl).
- 10
- 15




- The groups and , when R' is hydrogen, are referred to at times herein as 1- and 3-tetrahydro-isoquinolinecarbonyl, respectively, and are respectively abbreviated 1-Tiq and 3-Tiq.
- 20

15



The groups and , when


R' is hydrogen, are referred to at times herein as 1- and 3-perhydro-isoquinolinecarbonyl, respectively, and are respectively abbreviated 1-Piq and 3-Piq. As indicated by the crooked lines, various ring fusion isomers of these substituents exist -- this invention contemplates any individual isomer and combinations thereof.



The groups

and are

referred to as prolinyl and azetidine-2-carbonyl, respectively, and are respectively abbreviated Pro and Azt.




The group

represents a

saturated bicyclic system of the 4,5; 5,5; 6,5; 7,5; or 8,5 type. The stereochemistry at 3a is cis to the carbonyl; the other bridgehead bond may be either cis or trans except for the 4,5 and 5,5 systems must be cis at the bridgehead. The definitions of R^y and R^z provide that the variable ring, which includes the three carbon atoms shown, is a saturated carbocyclic system of 4-8 atoms. All of the ring atoms may be carbon, or one of the ring atoms may be a hetero atom selected from -O-, -S-, and -NH-. This definition includes

the preferred moiety derived from octahydroindole-2-carboxylic acid, abbreviated "Ohi", as represented by




The various cis and trans forms of this moiety are contemplated by this invention.

The asterisks in radical Y denote a chiral center that is (L). The asterisk in radical X denotes a chiral center that is (D) or (DL); the # in radical X denotes a chiral center that is (L).

In addition, diastereomers may exist depending upon branching of alkyl substituents. The compounds of the present invention include mixtures of two or more diastereomers as well as each individual isomer.


Preferred compounds of the present invention include those compounds of Formula I where X is



, homoprolinyl, 1- or 3-Tiq, or 1- or 3-Piq, and Y is prolinyl, and pharmaceutically acceptable salts and solvates thereof. In particular, compounds wherein Q is NHA and A is hydrogen or a sulfonamide (e.g., A = R"SO₂-), R' is hydrogen, Z is hydrogen, and B is hydrogen are all preferred. Also, those compounds wherein R is a guanidino, or especially, an amidino group are preferred.


One particularly preferred combination of substituents is where G is R-substituted phenyl (i.e., D=E=CH, k=0); especially preferred is where G is a 4-amidinophenyl group.

A preferred group of those compounds in which one to all of the otherwise unsubstituted carbon atoms of the aromatic or heteroaromatic rings of



bears a fluoro substituent is one in which no fluoro substituent is α - or γ - to D or E when D or E is N.


- 5 Another group of preferred compounds of the present invention includes those compounds of formula I as defined





- above where X is in which T is cyclohexyl, a is 1, R' is hydrogen and Q is -NH-A. One particular subgroup is one in which A is hydrogen. A second particular subgroup is one in which A is R"SO₂-, especially when R" is ethyl. A third particular subgroup is one in which A is -(CH₂)_g-COOH; preferably g is 1.

- Particular values of Y for a compound of Formula I in which X, r and G are defined as above include (L)-prolinyl (Pro), (S)-cis-octahydro-1H-indole-2-carbonyl (Ohi) and N-(2-phenylethyl)glycyl [N(PhCH₂CH₂)Gly].


- For a compound of Formula I in which R is -NH₂, it is preferred that the values of X and Y be selected from those defined above and the values of r and G be selected from


- a) r is 1 and G is  in which the anilino ring may bear one or two fluoro substituents;

- b) r is 1 and G is  ; and

- c) r is 1 or 2 and G is .

- One particularly preferred group of compounds of Formula I is one in which Y is (L)-prolinyl, r is 1 and G has


the value  $(CH_2)_k-R$ in which each of D and E is CH, k is 0 and R is amidino, and which may be represented by Formula Ia




wherein the benzamidine ring is unsubstituted or may bear one or two fluoro substituents, preferably meta- to the amidino radical and X has any of the values defined above.

A more particular value for a compound of Formula Ia is one in which the benzamidine ring is unsubstituted.

Another particularly preferred group of compounds of Formula I is one in which Y is (L)-prolinyl, r is 1 and G


has the value  in which M is S, each W is CH and R is amidino, and which may be represented by Formula Ib




wherein X has any of the values defined above.

An additional preferred group of compounds of formula I is one in which Y is (L)-prolinyl, r is 1 and G has

19


the value  in which D is N or CH, k is 0 and R is amidino, and which may be represented for Formula Ic



5

wherein X has any of the values defined above and D is N or CH.

A preferred value for X for a compound of formula



- 10 Ia, Ib or Ic is in which R' is hydrogen, a is 1, T is cyclohexyl or phenyl and Q is -NH-A. More particularly A is hydrogen, ethylsulfonyl or carboxymethyl. One particularly preferred value for X is N-carboxymethyl-D-cyclohexylalanyl. Another preferred value for X is
- 15 N-carboxymethyl-D-phenylalanyl.

Specific compounds of Formula I of the invention are described in the Examples. A preferred species, which may be employed as a pharmaceutically acceptable salt or solvate, may be selected from those disclosed as Examples 15, 20 18, 23, 44, 45, 46, 48, 49, 51, 52, 56, 65, 66, 68, 69, 70, 71, 72, 80, 86, 87, 88 and 92. A more preferred species may be selected from the compounds disclosed by Examples 45, 46, 48, 51, 65, 70, 71 and 72. One most preferred species based upon its unexpectedly superior properties is Example 48.

25 Another highly preferred species is Example 65.




As mentioned above, the invention includes pharmaceutically acceptable salts of the compounds defined by the above Formula I. A particular compound of this invention can possess one or more sufficiently basic functional groups,

and accordingly react with any of a number of inorganic and organic acids, to form a pharmaceutically acceptable salt. Acids commonly employed to form acid addition salts are inorganic acids such as hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid, phosphoric acid, and the like, and organic acids such as p-toluenesulfonic, methanesulfonic acid, oxalic acid, p-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid, acetic acid, and the like. Examples of such pharmaceutically acceptable salts thus are the sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, gamma-hydroxybutyrate, glycollate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate, and the like. Preferred pharmaceutically acceptable acid addition salts are those formed with mineral acids such as hydrochloric acid, hydrobromic acid and sulfuric acid.

The compounds of the present invention are known to form hydrates and solvates with appropriate solvents. Preferred solvents for the preparation of solvate forms include water, alcohols, tetrahydrofuran, DMF, and DMSO. Preferred alcohols are methanol and ethanol. Other appropriate solvents may be selected based on the size of the solvent molecule. Small solvent molecules are preferred to facilitate the corresponding solvate formation. The solvate or hydrate is typically formed in the course of recrystallization or in the course of salt formation. One useful reference concerning solvates is Sykes, Peter, A

Guidebook to Mechanism in Organic Chemistry, 6th Ed (1986, John Wiley & Sons, New York). As used herein, the term "solvate" includes hydrate forms, such as monohydrates and dihydrates.

- 5 The compounds of Formula I are prepared by known methods of peptide coupling. According to one such method the acid P-X'-COOH, where -X'-C(O)- is -X-, has the same meaning as defined for Formula I, and P is an amino protecting group, if necessary, is coupled with a carboxy
10 protected Y compound to form the dipeptide (a). The carboxy protecting ester group of the Y moiety is then removed (deblocked or de-esterified) and the free acid form of the dipeptide (b) is coupled with the protected reagent (d). The above reaction sequence is illustrated by the following
15 Scheme 1:



wherein G' is the same as G except R is -CN, -NHP,



- each P represents an amino protecting
20 group, if necessary, P' is H or P, alk is lower alkyl or some similar carboxylic acid protecting group, and -Y'- is the same as Y with the amino and carboxy functionalities visible, i.e., -Y- is the same as -N-Y'-C(O)-.

- If present, the cyano group in G' is elaborated to
25 a value of R; and the protecting groups in (c) are then

removed by procedures known to those skilled in the art such as hydrogenation over a metal catalyst to provide the compounds of Formula I.

The coupling of a P-X'-COOH compound with HN-Y'-COO-alk is carried out by first protecting the amino group of the amino acid, if any. Conventional amino protecting groups commonly used for temporary protection or blocking of the amino group are employed.

The amino-protecting group refers to substituents of the amino group commonly employed to block or protect the amino functionality while reacting other functional groups on the compound. Examples of such amino-protecting groups include the formyl group, the trityl group, the phthalimido group, the trichloroacetyl group, the chloroacetyl, bromoacetyl and iodoacetyl groups, urethane-type blocking groups such as benzyloxycarbonyl, t-butoxycarbonyl, 4-phenylbenzyloxycarbonyl, 2-methylbenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 4-fluorobenzyloxycarbonyl, 4-chlorobenzyloxycarbonyl, 3-chlorobenzyloxycarbonyl, 2-chlorobenzyloxycarbonyl, 2,4-dichlorobenzyloxycarbonyl, 4-bromobenzyloxycarbonyl, 3-bromobenzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, 4-cyanobenzyloxycarbonyl, 2-(4-xenyl)isopropoxycarbonyl, 1,1-diphenyleth-1-yloxycarbonyl, 1,1-diphenylprop-1-yloxycarbonyl, 2-phenylprop-2-yloxycarbonyl, 2-(p-toluy1)prop-2-yloxycarbonyl, cyclopentanyloxycarbonyl, 1-methylcyclopentanyloxycarbonyl, cyclohexanyloxycarbonyl, 1-methylcyclohexanyloxycarbonyl, 2-methylcyclohexanyloxycarbonyl, 2-(4-toluy1sulfonyl)ethoxycarbonyl, 2-(methylsulfonyl)ethoxycarbonyl, 2-(triphenylphosphino)ethoxycarbonyl, 9-fluoroenylmethoxycarbonyl ("Fmoc"), 2-(trimethylsilyl)ethoxycarbonyl, allyloxycarbonyl, 1-(trimethylsilylmethyl)prop-1-enyloxycarbonyl, 5-benzisoxalylmethoxycarbonyl, 4-acetoxybenzyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, 2-ethynyl-2-propoxycarbonyl,


cyclopropylmethoxycarbonyl, 4-(decyloxy)benzyloxycarbonyl, isobornyloxycarbonyl, 1-piperidyloxycarbonyl and the like; the benzoylmethylsulfonyl group, the 2-(nitro)phenylsulfonyl group, the diphenylphosphine oxide group, and the like amino-protecting groups. The species of amino-protecting group employed is not critical so long as the derivatized amino group is stable to the condition of subsequent reaction(s) on other positions of the molecule and can be removed at the appropriate point without disrupting the remainder of the molecule. Preferred amino-protecting groups are the benzyloxycarbonyl, allyloxycarbonyl, t-butoxycarbonyl, and trityl groups. Similar amino-protecting groups used in the cephalosporin, penicillin and peptide art are also embraced by the above terms. Further examples of groups referred to by the above terms are described by J. W. Barton, "Protective Groups in Organic Chemistry", J. G. W. McOmie, Ed., Plenum Press, New York, N.Y., 1973, Chapter 2, and T. W. Greene, "Protective Groups in Organic Synthesis", John Wiley and Sons, New York, N.Y., 1981, Chapter 7. The related term "protected amino" defines an amino group substituted with an amino-protecting group discussed above.

In carrying out the coupling reaction, an ester protecting group for HN-Y'-COOH is employed which is removable by conditions under which the amino protecting group remains intact. The amino protecting group of the acylating acid P-X'-COOH thus remains in place for protection of the amino group during the subsequent coupling with amine (d) to form (c).

The carboxy protecting ester group as used in the specification refers to one of the ester derivatives of the carboxylic acid group commonly employed to block or protect the carboxylic acid group while reactions are carried out on other functional groups on the compound. Examples of such carboxylic acid protecting groups include C₁-C₄ alkyl, benzyl, 4-nitrobenzyl, 4-methoxybenzyl, 3,4-dimethoxybenzyl, 2,4-dimethoxybenzyl, 2,4,6-trimethoxybenzyl, 2,4,6-trimethylbenzyl, pentamethylbenzyl,


- 3,4-methylenedioxybenzyl, benzhydryl,
 4,4'-dimethoxybenzhydryl, 2,2',4,4'-tetramethoxybenzhydryl,
 t-butyl, t-amyl, trityl, 4-methoxytrityl,
 4,4'-dimethoxytrityl, 4,4',4"-trimethoxytrityl, 2-phenylprop-
 5 2-yl, trimethylsilyl, t-butyldimethylsilyl, phenacyl,
 2,2,2-trichloroethyl, β -(trimethylsilyl)ethyl,
 β -(di(n-butyl)methylsilyl)ethyl, p-toluenesulfonylethyl,
 4-nitrobenzylsulfonylethyl, allyl, cinnamyl,
 1-(trimethylsilylmethyl)-prop-1-en-3-yl, and like moieties.
- 10 The species of carboxy-protecting group employed is not
 critical so long as the derivatized carboxylic acid is stable
 to the conditions of subsequent reaction(s) on other
 positions of the molecule and can be removed at the
 appropriate point without disrupting the remainder of the
 15 molecule. In particular, it is important not to subject the
 carboxy-protected molecule to strong nucleophilic bases or
 reductive conditions employing highly activated metal
 catalysts such as Raney nickel. (Such harsh removal
 conditions are also to be avoided when removing amino-
 20 protecting groups discussed below.) Further examples of
 these groups are found in E. Haslam, "Protective Groups in
 Organic Chemistry", J.G.W. McOmie, Ed., Plenum Press, New
 York, N.Y., 1973, Chapter 5, and T.W. Greene, "Protective
 Groups in Organic Synthesis", John Wiley and Sons, New York,
 25 N.Y., 1981, Chapter 5.

The compounds of Formula I can also be prepared by
 first synthesizing the $\text{HN-Y'-CONH(CH}_2)_r\text{-G'}$ amide precursor and
 then reacting with a protected X-moiety. According to one
 such method, (d) is prepared and coupled with PN-Y'-COOH (g)
 30 as shown below to afford the amide (h).



where P represents an amino protecting group such as the
 35 benzyloxycarbonyl (Cbz) group, t-butoxycarbonyl (Boc),
 p-toluenesulfonyl, and the like. Preferably the amino

protecting group used is removable by hydrogenation or treatment with mild acid (e.g. trifluoroacetic acid) or a strong acid (e.g. HCl). Examples of other suitable amino protecting groups are provided in "Protective Groups in Organic Synthesis", Second Edition, by T. W. Greene and Peter G. M. Wuts, Chapter 7, page 309-405 (1991), John Wiley & Sons, Inc., publishers. The Boc, or other suitable protecting group, is removed from the amino nitrogen of the Y-residue which is then acylated with the desired amino acid acyl group to afford the dipeptide shown below.







The cyano group, if present in G', is elaborated and protecting groups on (c) are removed as described earlier.

The coupling of a P-X'-COOH compound is carried out by first protecting the amino group of the amino acid, if any. Conventional amino protecting groups commonly used for temporary protection or blocking of the amino group are employed. Examples of such protecting groups are described above.

The coupling reactions described above are carried out in the cold preferably at a temperature between about -20 °C and about 15 °C. The coupling reactions are carried out in an inert organic solvent such as dimethylformamide, dimethylacetamide, tetrahydrofuran, methylene chloride, chloroform, and like common solvents or a mixture of such solvents. Generally, anhydrous conditions are used when, in the coupling reaction, an active ester of the acylating acid is used.

The intermediates (d) and (g) are prepared via standard techniques of organic chemistry as summarized in the following schemes:



where $-\text{K}-\text{R}$ is $-(\text{CH}_2)_r-\text{G}$.

According to the above sequences, protected guanidines can be prepared by doubly protecting S-methylisothiurea. A preferred blocking group is the t-butyloxycarbonyl (Boc) group which can be introduced by allowing S-methylisothiurea to react in the presence of di-t-butyl dicarbonate. Often an acid salt form of S-methylisothiurea is employed which can be sprung to the free base form in situ by dissolving the salt in water and treating with aqueous base. The di-t-butyl dicarbonate is then introduced to the reaction in a water miscible solvent, such as t-butanol, to give the doubly protected S-methylisothiurea. The desired doubly protected guanidine is then formed upon treatment with the appropriate diamine $\text{H}_2\text{N}-\text{K}-\text{NH}_2$ in a non reactive solvent or combination of solvents. Typically water miscible solvents such as dimethylformamide, or water, or mixtures thereof, are effectively employed. Such reaction is generally complete in about 3-72 hours. The resulting protected guanidine can then be coupled as previously described to provide the protected intermediates to the compounds of

Formula I wherein R is $-\overset{\text{NH}}{\underset{\parallel}{\text{C}}}-\text{NH}_2$.

For compounds of Formula I wherein $\text{R} = -\text{NH}_2$, the intermediate is a singly protected diamine. In most cases this

intermediate can be prepared by simply allowing an unprotected diamine to react with one molar equivalent of the protecting reagent. Other methods of introducing the final amine ($R = -NH_2$) will be apparent to organic chemists. For example, the amine may be obtained from some other precursor functionality, e.g., a nitro group or a cyano group. In the case of a nitro group, the transformation to an amino group is usually accomplished on those substrates where the nitro group is directly attached to an aromatic ring, particularly a phenyl group. In such cases, the nitrophenyl group is reduced to the corresponding aniline functionality by any of a number of methods known in the art. One particularly effective method is treatment of the nitro compound with sodium hydrosulfite in a non-reactive solvent, such as ethanol, water, or a mixture thereof. When the nitro compound is heated at reflux in a water/ethanol mixture in the presence of sodium hydrosulfite, reduction is usually complete within several hours. A cyano group can similarly be reduced, when appropriate, in the presence of a reducing agent such as lithium aluminum hydride, borane in a solvent such as tetrahydrofuran, or by metal promoted sodium borohydride reduction.




The amidines of this invention ($R = -C-NH_2$) can also be prepared from a nitrile precursor. A number of procedures for accomplishing this transformation are known in the art. In particular, the use of hydrogen sulfide in a mixture of pyridine and triethylamine, followed by treatment with acetone and methyl iodide, and finally ammonium acetate in methanol, is a preferred and efficient means for making this conversion. Alternatively, heating the nitrile with hydroxylamine hydrochloride and a base such as N,N-diisopropylethylamine in a hydroxylic solvent such as ethanol followed by catalytic hydrogenation (e.g., hydrogenolysis over palladium on carbon) can also be used to effect this transformation. This process

provides the hydroxy amidine ($R = \overset{\text{NOH}}{\parallel} \text{-C-NH}_2$) as an intermediate which can be isolated if desired.

The other compounds employed as initial starting materials in the synthesis of the compounds of this invention are well known and, to the extent not commercially available, are readily synthesized by standard procedures commonly employed by those of ordinary skill in the art.

The 4-substituted prolines (R^P is $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_8$ cycloalkyl, or $-(\text{CH}_2)_p\text{-L-(CH}_2)_q\text{-T'}$) used for making the compounds this invention are all of the *cis* configuration the 4-substituent relative to the carbonyl moiety. Intermediates for introducing this functionality into the compounds of Formula I are made by standard techniques.

For example, 4-substituted proline derivatives in which the R^P group contains a methylene group at the point of attachment to the proline ring can be prepared in the following manner:




where $R^4 = R^3\text{CH}_2 =$ an R^P group containing a methylene group at the point of attachment to the proline ring.

A 4-hydroxyproline (both the *cis* and *trans* forms are commercially available) is first protected with an amino-protecting group - the Cbz group is particularly useful in this sequence. The resulting intermediate is then esterified

(the methyl or especially ethyl esters are especially convenient) and then oxidized to give the corresponding ketone. This oxidation is accomplished under any of a number of oxidation conditions such as Jones oxidation or pyridinium chlorochromate; especially useful for this transformation is the use of pyridinium chlorochromate in a dry, non-reactive solvent such as dichloromethane. When allowed to react for 8-16 hours, this reaction is generally complete when performed at ambient temperature. This versatile ketone intermediate is then allowed to react with an appropriate Wittig reagent to give the desired olefin. Typically the appropriate R^P -substituted triphenylphosphonium halide is added to a dry inert solvent (e.g., tetrahydrofuran) which contains a strong base (e.g., potassium *t*-butoxide). The ketone is introduced and after approximately three hours at ambient temperature the desired olefin intermediate can be isolated. In order to obtain good yields of the olefin, it is preferred that a 0.4-0.6 molar excess of the Wittig reagent be employed relative to the ketone. The olefin is then reduced to the desired R^P -substituted proline by standard reduction techniques. Catalytic hydrogenation is the most facile method for accomplishing this transformation in the laboratory. Hydrogenation of the olefin in the presence of a catalyst (e.g., 5% palladium on carbon) in an inert solvent such as ethanol will be effective at atmospheric pressure. In the case of those intermediates in which the amino-protecting group is Cbz, hydrogenation also removes the protecting group which provides a compound which can be used for coupling to $P-X'-COOH$. As will be appreciated by those skilled in this art, this process will not be effective for preparing compounds where the R^P group is attached to the proline ring through a hetero atom or are an aromatic ring. Thus, in the above scheme, R^3 will be alkyl, aralkyl (e.g., benzyl), (cycloalkyl)alkyl, etc.


A related method for preparing these intermediates is summarized by the following scheme:

30



The above reaction scheme is an alternative to the Wittig reaction described earlier and is useful for preparing compounds for which Wittig reagents cannot be prepared. Thus, for preparing intermediates wherein R_a is alkyl, phenyl, and the like, the pyrrolidinone intermediate is allowed to react with an appropriate Grignard reagent. Typically a slight molar excess of the Grignard reagent is employed, usually at low temperatures (e.g., -80 to -60 °C) in a low freezing inert solvent such as tetrahydrofuran. After addition of the reagents, the reaction mixture can be permitted to warm to room temperature, after which time the reaction is usually complete within several hours. The resulting intermediate is dehydrated, for example, by treatment with trifluoroacetic acid. The 3,4-dehydro intermediate is then reduced to the desired *cis* intermediate using the same reductive conditions as described above for reduction of the olefin intermediate.

Intermediates wherein the hetero "L"-group is oxygen and is attached directly to the proline ring (i.e., p=0) can be prepared employing the Mitsunobu reaction (Mitsunobu, Synthesis, 1 (1981)):



25

In this reaction, the *trans* hydroxypyrrolidine-carboxylic ester is treated with triphenylphosphine in a


solvent such as tetrahydrofuran in the presence of Ar-O-H. The mixture is cooled to approximately 0 °C and diethyl azodicarboxylate is added. After warming to room temperature, the reaction is worked up to provide the desired
5 cis intermediate. While the scheme above depicts the reaction for compounds where L = -O-, p=q=0, and T=Ar, this sequence is useful for preparing other compounds where p=0 and L is -O-.

Intermediates wherein L is sulfur and is attached
10 directly to the ring can be prepared by first converting the hydroxy group to a tosylate or other similar leaving group and then displacing with a thiolate anion (see, e.g., Krapcho, et al., J. Med. Chem., 31, 1148-1160 (1988); Smith, et al., J. Med. Chem., 31, 875-855 (1988)).

15 Intermediates wherein L is nitrogen and is attached directly to the ring can be prepared by first converting the hydroxy group to a tosylate or other similar leaving group and then displacing with azide. The azide can be reduced using known methods and then alkylated to provide the desired
20 functionality (see, e.g., Smith, et al., J. Med. Chem., 31, 875-855 (1988)).


The compounds of this invention containing a cis-Ohi
Ohi functionality are prepared by preparing (S)-indoline
carboxylic acid ethyl ester from the corresponding acid (see,
25 Vincent, et al., Drug Design and Discovery, Vol. 9, pp 11-28 (1992)), and reducing this intermediate by hydrogenation over 5% Pd/C in ethanol to give the octahydroindole-2-carboxylic acid ester, generally referred to Ohi-ester as summarized below.

30



The compounds of this invention containing a trans-Ohi
functionality are prepared by the method of Vincent, et al., ...


Drug Design and Discovery, Vol. 9, pp 11-28 (1992)). This is summarized in the scheme shown below:



5

The compound of this invention containing a bicyclic system (with or without heteroatom) can be prepared by the method of Teetz, *et al.*, Tetrahedron Letters, **25**, 4479 (1984). Generally:


10



where P is a protecting group and R^x is alkyl.


The intermediates for introducing the N-substituted glycine functionality (Y) used for making the compounds this invention are made by standard techniques.

For example, a haloacetate ester, such as t-butyl bromoacetate, can be converted into the desired substituted upon treatment with the appropriate primary amine:



The t-butyl bromoacetate is allowed to react with the appropriate amine either neat or preferably in a non-reactive solvent, such as an alcohol. It is preferred that a molar excess of the amine is used to force the reaction to completion. Preferably the reaction mixture also contains a non-reactive acid scavenger, such as at least a molar equivalent of triethylamine. While the reactants are usually combined cooled (e.g., 0 °C), the reaction is usually allowed to warm to room temperature after which the reaction is usually complete within 24 hours. Although the bromoacetate is preferred, other haloacetates, such as iodoacetates and chloroacetates, can be employed for this transformation. Other ester groups can similarly be employed. The t-butyl ester is preferred because it can later be easily removed later upon treatment with anisole and trifluoroacetic acid.

A second method for preparing these intermediates is summarized by the following scheme:



where $\text{R}^o\text{-CH}_2\text{-}$ is an R^g -group having an unsubstituted methylene group adjacent to the point of attachment to the glycine moiety.

In the above reaction scheme, the appropriate aldehyde is mixed with glycine ester in a non-reactive solvent, such as methanol or ethanol. If a salt form of the glycine ester is used, a molar equivalent of a base, such as potassium hydroxide, can be added to allow generation of the

free base of the aminoester. The reaction of the aldehyde and glycine ester formed the intermediate Schiff base which can then be reduced *in situ* upon treatment with a reducing agent such as sodium cyanoborohydride. Formation of the Schiff base occurs in usually less than an hour; reduction is generally complete after 10-15 hours. The methyl or ethyl esters are particularly useful as these groups can be removed (deblocked) upon treatment with lithium hydroxide in aqueous dioxane. Employing an appropriate ketone instead of aldehyde R^O-CHO results in the preparation of intermediates wherein the methylene group attached to the glycine amine is substituted.

Alternatively, and especially for those compounds wherein R^G is Ar (i.e., without an intervening alkyl group), it is preferred to prepare the intermediate $P-X'-CONHAr$ by standard techniques (e.g., reacting an activated form of $P-X'-COOH$ with $ArNH_2$) and then reacting this intermediate with an alkyl haloacetate as described above to give $P-X'-CONHAr-CH_2-COO-alk$ which can then be further transformed in the usual way.


Many of the final compounds of this invention or intermediates thereto can be interconverted by standard techniques. For example, aryl compounds which are substituted with nitro can be reduced (e.g., in the presence of sodium hydrosulfite in a non-reactive solvent, such as ethanol, water, or a mixture thereof). When the nitro compound is heated at reflux in a water/ethanol mixture in the presence of sodium hydrosulfite, reduction is usually complete within several hours. The resulting amine may be present in the final product; if the amine is present in an intermediate, it may be desirable to convert it to its final desired form (e.g., acylation to provide the acylated amine) or protected to avoid side reactions during the subsequent chemistry. If the free amine is the desired compound, the Cbz protecting group is particularly useful in this regard. Other transformations and intraconversions of this type will be apparent to skilled organic chemists.

As will be appreciated by those skilled in this art, the above transformations can be performed on the starting materials noted above, or in most cases can also be accomplished on di- or tri-peptide intermediates containing the same respective functional group. In the latter cases, the need, or lack thereof, to protect the various groups may be negated; accordingly, the order and type of chemistry involved will dictate the need and type of protecting groups as well as the sequence for accomplishing the synthesis. As will also be appreciated by skilled artisans, one may choose other protecting groups so long as they serve the purpose of protecting the functional group during subsequent chemistry but can also be removed under appropriate conditions and in an appropriate order to allow for subsequent transformations. For example, in Scheme 1 above G' includes substituents wherein R is -CN; this nitrile group can be transformed into an amidine or reduced to the amine which can optionally be further elaborated to the guanidines of this invention.

The compounds of the invention are isolated best in the form of acid addition salts. Salts of the compounds of Formula I formed with acids such as those mentioned above are useful as pharmaceutically acceptable salts for administration of the antithrombotic agents and for preparation of formulations of these agents. Other acid addition salts may be prepared and used in the isolation and purification of the peptides. For example, the salts formed with the sulfonic acids such as methanesulfonic acid, n-butanesulfonic acid, p-toluenesulfonic acid and naphthalenesulfonic acid may be so used.

A compound of Formula I is prepared by:

a) removing simultaneously or sequentially the protecting group(s) P of a corresponding compound of Formula II



wherein (P)X represents a radical X which may bear one or more protecting groups P independently selected from an amino protecting group P for a compound of Formula I in which X

includes a basic NH moiety and a carboxy protecting group P for a compound of Formula I in which X includes a carboxy residue and G(P) represents a radical G which may bear one or more independently selected amino protecting groups P; or

5 b) for a compound of Formula I in which R is



, by hydrogenolysis of a corresponding compound of



Formula I in which R is $-\text{C}-\text{NH}_2$; and whereafter, when a salt of the compound of Formula I is required, forming the salt with a pharmaceutically acceptable acid.

10 It may be preferred to conduct process b) simultaneously with process a). For a compound of Formula I in which an acid protecting group is the t-butyl ester and/or an amino protecting groups is t-butyloxycarbonyl, the protecting group(s) may be removed by treatment with a
15 strong acid, such as trifluoroacetic acid or anhydrous hydrogen chloride in an inert solvent, such as dioxane or dichloromethane, in the presence of anisole. For a compound of Formula I in which an acid protecting group is the benzyl ester and/or an amino protecting group is
20 benzyloxycarbonyl, the protecting group(s) may be removed by hydrogenolysis, conveniently carried out in ethanolic hydrogen chloride over a palladium on carbon catalyst.

 The preferred method for purifying the compounds of Formula I, while at the same time preparing a desired stable
25 salt form, is that described in U.S. Patent 5,250,660. According to the method, stable sulfates or hydrochlorides are provided by preparative purification over C₁₈ reversed-phase chromatography in which the aqueous component comprises sulfuric acid or hydrochloric acid at pH 2.5 and acetonitrile
30 is the organic component. The pH of the acidic eluant is adjusted to between about pH 4 and about 6 with an anion exchange resin in the hydroxyl form, e.g. Bio-Rad AG-1X8. After adjustment of the pH, the solution of tripeptide sulfate or hydrochloride salt is lyophilized to provide the

pure salt in dry powder form. In an example of the process, crude D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂ sulfate can be dissolved in water and the solution is loaded on Vydac Cl₈ RP HPLC 5 cm X 50 cm column. A gradient of 2-10% B (A = 0.01% H₂SO₄; B =
5 acetonitrile) over 10 hours is used. Multiple fractions are collected and those containing product as determined by analytical RP HPLC are pooled. The pH of the pooled fractions is adjusted to pH 4.0 - 4.5 with AG-1X8 resin in hydroxide form (Bio-Rad, 3300 Ragatta Blvd., Richmond, CA
10 94804). The solution is filtered and the filtrate is lyophilized to provide the pure D-,L- diamide in the form of the sulfate salt.

The optically active isomers of the diastereomers at the radical X are also considered part of this invention.
15 Such optically active isomers may be prepared from their respective optically active precursors by the procedures described above, or by resolving the racemic mixtures. This resolution can be carried out by derivatization with a chiral reagent followed by chromatography or by repeated
20 crystallization. Removal of the chiral auxiliary by standard methods affords substantially optically pure isomers of the compounds of the present invention or their precursors. Further details regarding resolutions can be obtained in Jacques, et al., Enantiomers, Racemates, and Resolutions,
25 John Wiley & Sons, 1981.

The compounds employed as initial starting materials in the synthesis of the compounds of this invention are well known and, to the extent not commercially available, are readily synthesized by standard procedures commonly
30 employed by those of ordinary skill in the art.

The following Examples are provided to further describe the invention and to provide examples for comparison and are not to be construed as limitations thereof.

The abbreviations used in this specification have
35 the following meanings.


Amino acid residues: Arg = arginyl, Glu = glutamyl, Gly = glycyl, Pro = prolyl, hPro = homoprolyl, Azt

- = azetidine-2-carbonyl, Phg = phenylglycyl, Phe = phenylalanyl, hPhe = homophenylalanyl, 1-Tiq = 1,2,3,4-tetrahydroisoquinoline-1-carbonyl, 3-Tiq = 1,2,3,4-tetrahydroisoquinoline-3-carbonyl, Cha = β -cyclohexylalanyl, hCha = α -amino- γ -cyclohexylbutyryl, NMI = N-methylindol-2-oyl, Ohi = cis-octahydroindol-2-oyl, 1-Piq = perhydro-isoquinoline-1-carbonyl, 3-Piq = perhydro-isoquinoline-3-carbonyl, Met = methionyl, Met(O₂) = S,S-dioxomethionyl.
- 10 Agm = agmatine
Boc = t-butyloxycarbonyl
Bn = benzyl
Cbz = benzyloxycarbonyl
DCC = dicyclohexylcarbodiimide
- 15 DMF = dimethylformamide
Et = ethyl
DMSO = dimethylsulfoxide
EtOAc = ethyl acetate
Et₂O = diethyl ether
- 20 EtOH = ethanol
Fmoc = 9-fluorenylmethoxycarbonyl
FAB-MS = fast atom bombardment mass spectrum
FD-MS = field desorption mass spectrum
IS-MS = ion spray mass spectrum
- 25 HRMS = high resolution mass spectrum
HOBT = 1-hydroxybenzotriazole hydrate
IR = infrared spectrum
RPHPLC = Reversed Phase High Performace Liquid
Chromatography
- 30 Ph = phenyl
TFA = trifluoroacetic acid
THF = tetrahydrofuran
TLC = thin layer chromatography
- The following parameters for RPHPLC were employed:
- 35 Solvent A: 0.05% aqueous hydrochloric acid (1.5 mL concentrated hydrochloric acid in 3 L water); Solvent B: acetonitrile; Gradient: as defined in each Example; Method 1:

Column: Vydac C₁₈ - 2.5 cm x 25 cm; Flow rate: 5 mL/minute;
Method 2: Column: Vydac C₁₈ - 5 cm x 25 cm; Flow rate: 10
mL/minute; Method 3: Column: Vydac C₁₈ - 2.5 cm x 50 cm; Flow
rate: 10 mL/minute.

5 Unless otherwise stated, pH adjustments and work up
are with aqueous acid or base solutions.

 In the examples, where ¹H-NMR is shown, the product
afforded by the reaction was characterized by proton NMR to
confirm the indicated compound was obtained; IR without the
10 data similarly indicates a satisfactory infrared spectrum was
obtained. HRMS was used to confirm the exact mass of
compounds for which a satisfactory elemental analysis was not
obtained for the product of the described procedure; the
elemental composition of the observed ion (e.g., MH⁺) is
15 indicated.

Example 1

5

EtSO₂-D-Phe-Pro-Agm·HCl

A) Preparation of Boc-D-Phe-Pro-OBn.

To a solution of Boc-D-Phe-OH (89.1 g, 336 mmol), Pro-OBn hydrochloride (81.2 g, 336 mmol), HOBT (50 g, 370 mmol) and N,N-diisopropylethylamine (176 mL, 1,008 mmol) at 0 °C in dichloromethane (600 mL) was added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (71 g, 370 mmol). After stirring for 18 hours, the mixture was diluted with diethyl ether (1 L) and washed sequentially three times with 1 N citric acid (250 mL), once with water (250 mL), three times with saturated aqueous sodium bicarbonate (250 mL) and once with saturated aqueous sodium chloride (250 mL). The organic phase was dried (Na₂SO₄), filtered, and concentrated in vacuo to yield 140 g (92.5%) of a pale yellow foam.

FD-MS, m/e 452 (M⁺)¹H NMR

25

B) Preparation of D-Phe-Pro-OBn·TFA.

To a stirring solution of Boc-D-Phe-Pro-OBn (68 g, 150 mmol) in dichloromethane (50 mL) at 0 °C, was added anisole (20 mL) followed by trifluoroacetic acid (400 mL). After stirring for 3 hours, the solvents were evaporated in vacuo and the thick oily residue was dissolved in diethyl ether

(1.5 L) and refrigerated (72 hours). The white precipitate was filtered, washed with diethyl ether (300 mL) and dried to yield 59.4 g (85%) of white powder.

5 ¹H NMR

C) Preparation of EtSO₂-D-Phe-Pro-OBn.

To a stirring solution of D-Phe-Pro-OBn·TFA (12 g, 25.7 mmol) and triethylamine (7 mL, 50.2 mmol) in dichloromethane (200 mL) at -78 °C was added ethanesulfonyl chloride (2.65 mL, 28.3 mmol) dropwise via an addition funnel. The reaction vessel was warmed to 0 °C and after stirring 4 hours, water (10 mL) was added. The organic phase was washed three times with 1 N hydrochloric acid (100 mL), once with a saturated sodium chloride solution (100 mL) and then the solvent was removed *in vacuo*. The product was purified by flash chromatography over silica gel, eluting with ethyl acetate/hexanes (6:4). The product containing fractions (judged by TLC) were combined and concentrated to give 6.62 g (58%) of a yellow oil which solidified.

¹H NMR

25 FD-MS, m/e 445 (M⁺)

Analysis for C₂₃H₂₈N₂O₅S:

Calc: C, 62.14; H, 6.35; N, 6.30;

Found: C, 61.87; H, 6.37; N 6.18.

30 D) Preparation of EtSO₂-D-Phe-Pro-OH.

To a stirring solution of EtSO₂-D-Phe-Pro-OBn (4.5 g, 10.1 mmol) in p-dioxane (150 mL) was added a solution of lithium hydroxide monohydrate (2.1 g, 50.5 mmol) in water (75 mL). After stirring for 16 hours, the volume of the solution was reduced by half *in vacuo*, and the solution was diluted with water (300 mL) and 0.1 N NaOH (100 mL). The

aqueous phase was then washed twice with diethyl ether (250 mL), acidified with solid citric acid, and then extracted three times with ethyl acetate (150 mL). The combined ethyl acetate extracts were washed with saturated aqueous sodium chloride (200 mL), dried (MgSO_4), filtered and concentrated to give 3.6 g (90%) of white solid.

FD-MS, m/e 355 (M^+)

Analysis for $\text{C}_{16}\text{H}_{22}\text{N}_2\text{O}_5\text{S}$:

10 Calc: C, 54.22; H, 6.26; N, 7.90;
 Found: C, 54.40; H, 6.42; N, 7.85.

E) Preparation of N,N'-di-Boc-S-methylisothiurea.

15 To a stirring solution of di-t-butyl dicarbonate (100 g, 458 mmol) in t-butanol (300 mL) was added a solution of bis-S-methylisothiurea sulfate (32.7 g, 117 mmol) in water (150 mL), followed by a solution of sodium hydroxide (19.2 g, 480 mmol) in water (150 mL). After stirring for 48
20 hours, the mixture was concentrated to approximately one-third of the original volume in vacuo and diluted with diethyl ether (500 mL). The organic phase was washed once with water (250 mL), three times with 1 N citric acid (250 mL) and once again with water (250 mL). The organic phase
25 was then dried (MgSO_4), filtered and concentrated in vacuo to give 42 g (62%) of a white solid.

^1H NMR

30 F) Preparation of N⁹,N^{9'}-di-Boc-agmatine.

 To a stirring solution of 1,4-butanediamine (23 g, 258 mmol) in 2:1 dimethylformamide:water (300 mL) was added a solution of N,N'-di-Boc-S-methylisothiurea (15 g, 52 mmol)
35 in dimethylformamide (100 mL) via an addition funnel. After stirring for 2 hours, the solvents were removed in vacuo and the residue was dissolved in 1 N citric acid (250 mL),

diluted with water (250 mL) and washed with ethyl acetate (250 mL). The ethyl acetate phase was back extracted with 1 N citric acid (100 mL) and the combined aqueous phases were basified with sodium carbonate, saturated with solid sodium chloride, and extracted twice with ethyl acetate (250 mL). The combined ethyl acetate extracts were washed with saturated aqueous sodium chloride (200 mL), dried (MgSO₄), filtered and concentrated to give 12.5 g (73%) of a thick syrup.

10

¹H NMR

G) Preparation of EtSO₂-D-Phe-Pro-Agm(Boc)₂.

15

To a stirring solution of N⁹,N^{9'}-di-Boc-agmatine (2 g, 6 mmol) in dichloromethane (30 mL) was added EtSO₂-D-Phe-Pro-OH (2.1 g, 6 mmol), HOBT (810 mg, 6 mmol) and N,N-diisopropylethylamine (1.6 g, 12 mmol), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride

20

(1.4 g, 73 mmol). After stirring for 20 hours, the solution was diluted with ethyl acetate (300 mL) and washed three times with 1 N citric acid (150 mL), once with water (150 mL), and twice with saturated aqueous sodium bicarbonate. The organic phase was then dried (MgSO₄),

25

filtered and concentrated in vacuo. The residue was chromatographed over silica gel, eluting with a step gradient of ethyl acetate:hexanes (1:4) through ethyl acetate. The product containing fractions (based on TLC) were combined and concentrated to give 2.4 g (60%) of a

30

thick oil.

¹H NMRFD-MS, m/e 668 (MH⁺)

H) Preparation of EtSO₂-D-Phe-Pro-Agm·HCl.


A stirring suspension of EtSO₂-D-Phe-Pro-Agm(Boc)₂ (1.6 g, 2.4 mmol) in anisole (1 mL) was dissolved in trifluoroacetic acid (20 mL) and allowed to stir for 1 hour at room temperature. The solvent was then removed in vacuo and the residue was partitioned between water (100 mL) and diethyl ether (50 mL). The aqueous phase was washed again with diethyl ether (50 mL) and then partially concentrated and lyophilized to yield 1.4 g of crude trifluoroacetate salt. Half of this material was then dissolved in water and purified by RPHPLC (method 1; 98/2 (A/B); ramp to 50/50 (A/B), 60 minutes) to give 490 mg (81%) of white powder.

15 ¹H NMRFD-MS, m/e 467 (M⁺)Analysis for C₂₁H₃₄N₆O₄S·HCl·H₂O:

Calc: C, 48.41; H, 7.16; N, 16.13; Cl, 6.80;

Found: C, 48.01; H, 6.81; N, 16.15; Cl, 6.97.

20

Example 2

25

EtSO₂-D-Cha-Pro-Agm·HCl

A) Preparation of Boc-D-Cha-Pro-OBn.

By a method substantially equivalent to that described in Example 1-A, Boc-D-Cha-Pro-OBn was prepared from Boc-D-Cha-OH and Pro-OBn·HCl in 91% yield (109 g).

FD-MS, m/e 458 (M^+)

B) Preparation of D-Cha-Pro-OBn·TFA.

5 By a method substantially equivalent to that described in Example 1-B, D-Cha-Pro-OBn·TFA was prepared (116% of theoretical yield, 130 g).

1H NMR

10 FD-MS, m/e 359 (M^+)

C) Preparation of EtSO₂-D-Cha-Pro-OBn.

15 By a method substantially equivalent to that described in Example 1-C, EtSO₂-D-Cha-Pro-OBn was prepared (20% yield, 2.3 g).

1H NMR

FD-MS, m/e 450 (M^+)

20 Analysis for C₂₃H₃₄N₂O₅S:

Calc: C, 61.31; H, 7.61; N, 6.22;

Found: C, 61.55; H, 7.59; N, 6.28.

D) Preparation of EtSO₂-D-Cha-Pro-OH.

25

By a method substantially equivalent to that described in Example 1-D, EtSO₂-D-Cha-Pro-OH was prepared (48% yield, 0.78 g).

30 1H NMR

FD-MS, m/e 361 (M^+)

E) Preparation of EtSO₂-D-Cha-Pro-Agm(Boc)₂.

35 By a method substantially equivalent to that described in Example 1-G, 400 mg (40%) of EtSO₂-D-Cha-Pro-Agm(Boc)₂

were prepared from EtSO₂-D-Cha-Pro-OH and N⁹-N^{9'}-di-Boc-Agm.

¹H NMR

5 FD-MS, m/e 674 (MH⁺)

F) Preparation of EtSO₂-D-Cha-Pro-Agm·HCl.

10 By a method substantially equivalent to that described in Example 1-H, EtSO₂-D-Cha-Pro-Agm·HCl was prepared (45% yield, 100 mg). The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 50/50 (A/B), 60 minutes).

¹H NMR

15 FD-MS, m/e 473 (M⁺)


Analysis for C₂₁H₄₀N₆O₄S·1.2HCl·H₂O:

Calc: C, 47.20; H, 8.15; N, 15.73; Cl, 7.96;

Found: C, 47.47; H, 7.84; N, 16.10; Cl, 7.80.

20

Example 3



EtOCO-D-Phe-Pro-Agm·HCl

25

A) Preparation of EtOCO-D-Phe-Pro-OH.

30 By methods substantially equivalent to those described in Examples 1-C and 1-D using ethyl chloroformate in place of ethanesulfonyl chloride, 6.59 g (92%) of EtOCO-D-Phe-Pro-OH were prepared.

¹H NMRFD-MS, m/e 335 (M⁺)Analysis for C₁₇H₂₂N₂O₅:

Calc: C, 61.07; H, 6.63; N, 8.38;

5 Found: C, 60.88; H, 6.72; N, 8.14.


B) Preparation of EtOCO-D-Phe-Pro-Agm·HCl.

By a method substantially equivalent to that described
 10 in Example 1-G, 2.1 g (54%) of EtOCO-D-Phe-Pro-Agm(Boc)₂
 were prepared from EtOCO-D-Phe-Pro-OH and N⁹-N^{9'}-di-Boc-
 Agm. Then by a method substantially equivalent to that
 described in Example 1-H, 390 mg (77%) of EtOCO-D-Phe-Pro-
 Agm·HCl were prepared. The product was purified by RPHPLC
 15 (method 1, 98/2 (A/B), ramp to 50/50 (A/B), 60 minutes).

¹H NMRFD-MS, m/e 447 (M⁺)Analysis for C₂₂H₃₄N₆O₄·0.9HCl·0.2TFA·H₂O:

20 Calc: C, 51.70; H, 7.22; N, 16.15; Cl, 6.13;

Found: C, 51.73; H, 7.20; N, 16.54; Cl, 6.36.

Example 4

25

NMI-D-Phe-Pro-Agm·HCl

(N-[(1-methyl-1H-indol-2-yl)carbonyl]-D-phenylalanyl-N-
 [4-[(aminoiminomethyl)amino]butyl]-L-prolinamide
 30 monohydrochloride)

A) Preparation of NMI-D-Phe-Pro-OH.

To a solution of N-methylindole-2-carboxylic acid (2.6 g, 14.9 mmol) in dry tetrahydrofuran (45 mL) was added
5 pentafluorophenol (3 g, 16.5 mmol), followed by 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (3.2 g, 16.5 mmol). The mixture was allowed to stir at reflux for 3.5 hours and then cooled to room temperature. To this mixture was added a solution of D-Phe-Pro-OBn·TFA (7 g, 14.9 mmol)
10 and N,N-diisopropylethylamine (4 g, 30 mmol) in tetrahydrofuran (25 mL). After stirring for an additional 2 hours, the solvents were removed *in vacuo* and the residue was dissolved in ethyl acetate (500 mL), then washed three times with 0.1 N aqueous sodium bisulfate (250 mL) and
15 three times with 1 N aqueous potassium carbonate (250 mL). The organic phase was dried (Na₂SO₄), filtered, and concentrated *in vacuo* to give 6.5 g of amorphous solid (a mixture of the desired product, contaminated with pentafluorophenol). This crude product was then hydrolyzed
20 by a method substantially equivalent to that described in Example 1-D to yield 3.8 g (62%) of an off white solid.

¹H NMRFD-MS, m/e 419 (M⁺)

25

B) Preparation of NMI-D-Phe-Pro-Agm·HCl.

By a method substantially equivalent to that described in Example 1-G, 900 mg (20%) of NMI-D-Phe-Pro-Agm(Boc)₂
30 were prepared. Then, by a method substantially equivalent to that described in Example 1-H, 144 mg (31%) of NMI-D-Phe-Pro-Agm·HCl were prepared. The crude product was dissolved in glacial acetic acid and purified by RPHPLC (method 1, 90/10 (A/B), ramp to 40/60 (A/B), 80 minutes).
35

¹H NMRFD-MS, m/e 532 (M⁺)


Analysis for $C_{29}H_{37}N_7O_3 \cdot 0.9HCl \cdot 0.6TFA \cdot 0.5H_2O$:

Calc: C, 56.46; H, 6.29; N, 15.27; Cl, 4.97;

Found: C, 56.77; H, 6.58; N, 15.35; Cl, 5.28.

5

Example 5



D-Phe-Pro-NH(CH₂)₆NHC(NH)NH₂ · HCl

10

A) Preparation of Boc-D-Phe-Pro-OH.

To a solution of Boc-D-Phe-Pro-OBn (145 g, 320 mmol) in p-dioxane (660 mL) was added a solution of lithium hydroxide monohydrate (54 g, 1,280 mmol) in water (330 mL) with vigorous stirring. After 4 hours, the solution was concentrated in vacuo to about one-fourth the original volume and diluted with water (350 mL) and 0.1 N sodium hydroxide (100 mL). The aqueous phase was washed three times with diethyl ether (250 mL) and then acidified to pH 3 with solid citric acid which caused a precipitate to form. The solid was filtered, washed twice with water, and then dried under high vacuum, to yield 91 g (78%) of a white solid.

25

¹H NMR

FD MS, m/e 363 (M⁺)

B) Preparation of N⁹,N^{9'}-di-Boc-6-aminohexylguanidine.

By a method substantially equivalent to that described in Example 1-F, 4.7 g (66%) of N⁹,N^{9'}-di-Boc-6-amino-
5 hexylguanidine were prepared from 1,6-hexanediamine.

C) Preparation of Boc-D-Phe-Pro-NH(CH₂)₆NHC(NBoc)NH(Boc).

By a method substantially equivalent to that described
10 in Example 1-G, 1.3 g (62%) of Boc-D-Phe-Pro-NH(CH₂)₆NHC(NBoc)NH(Boc) were prepared from Boc-D-Phe-Pro-OH and N⁹,N^{9'}-di-Boc-6-aminohexylguanidine.

¹H NMR

15 FD-MS, m/e 703 (M⁺)

D) Preparation of D-Phe-Pro-NH(CH₂)₆NHC(NH)NH₂·HCl.


By a method substantially equivalent to that described
20 in Example 1-H, approximately 100 mg of D-Phe-Pro-NH(CH₂)₆NHC(NH)NH₂·HCl were prepared.

FD-MS, m/e 389 (M⁺)

Analysis for C₂₁H₃₄N₆O₂·0.9HCl·0.9TFA·0.5H₂O:

25 Calc: C, 49.97; H, 6.95; N, 15.34;

Found: C, 49.60; H, 7.13; N, 15.23.

Example 6

D-Phe-Pro-NH(CH₂)₅NHC(NH)NH₂·HCl

A) Preparation of N⁹,N^{9'}-di-Boc-5-aminopentylguanidine.

By a method substantially equivalent to that described
5 in Example 1-F, 1.73 g (72%) of N⁹,N^{9'}-di-Boc-5-amino-
pentylguanidine were prepared from 1,5-pentanediamine.

FD-MS, m/e 345 (M⁺)¹H NMR

10

B) Preparation of Boc-D-Phe-Pro-NH(CH₂)₅NHC(NBoc)NH(Boc).

By a method substantially equivalent to that described
in Example 1-G, 1.9 g (92%) of Boc-D-Phe-Pro-
15 NH(CH₂)₅NHC(NBoc)NH(Boc) were prepared from Boc-D-Phe-Pro-
OH and N⁹,N^{9'}-di-Boc-5-aminopentylguanidine.

¹H NMRFD-MS, m/e 689 (M⁺)

20


C) Preparation of D-Phe-Pro-NH(CH₂)₅NHC(NH)NH₂·HCl.

By a method substantially equivalent to that described
in Example 1-H, approximately 100 mg of D-Phe-Pro-
25 NH(CH₂)₅NHC(NH)NH₂·HCl were prepared. The product was
purified by RPHPLC (method 1, 98/2 (A/B), ramp to 40/60
(A/B), 40 minutes).

FD-MS, m/e 389 (M⁺)30 Analysis for C₂₀H₃₂N₆O₂·0.9HCl·0.9TFA·0.7H₂O:

Calc: C, 48.71; H, 6.79; N, 15.63;

Found: C, 48.34; H, 6.68; N, 16.01.

Example 7

5 D-Phe-Pro-NH(CH₂)₃NHC(NH)NH₂ · HCl

A) Preparation of Boc-D-Phe-Pro-NH(CH₂)₃NHC(NBoc)NH(Boc) .

To a solution of 1,3-diaminopropane (2.2 g, 30 mmol)
 10 in dimethylformamide (25 mL) was added a solution of N,N'-
 di-Boc-S-methylisothioureia (2.9 g, 10 mmol) in
 dimethylformamide (25 mL). After stirring for 1 hour, the
 mixture was diluted with dichloromethane (400 mL) and
 washed twice with a mixture of saturated aqueous sodium
 15 bicarbonate and saturated aqueous sodium chloride (200 mL),
 and once with saturated aqueous sodium chloride (250 mL).
 The organic phase was dried (MgSO₄), filtered, and
 partially concentrated in vacuo to a volume of about 200
 mL.

20 To this solution was then added Boc-D-Phe-Pro-OH (3.6
 g, 10 mmol), HOBT (1.3 g, 10 mmol) and
 N,N-diisopropylethylamine (1.3 g, 10 mmol), followed by
 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide · HCl (2.1 g,
 11 mmol). After stirring for 16 hours, the solvents were
 25 removed in vacuo and the residue was taken up in ethyl
 acetate (250 mL). The organic phase was washed three times
 with 1 N citric acid (200 mL), once with water (100 mL),
 twice with saturated aqueous sodium bicarbonate (200 mL)
 and once with saturated aqueous sodium chloride. The
 30 organic phase was then dried (MgSO₄), filtered and
 concentrated in vacuo. The residue was then
 chromatographed over silica gel, eluting with a step

gradient of ethyl acetate/hexanes (1:4) through ethyl acetate. The product containing fractions (judged by TLC) were concentrated to yield 2.6 g (40%) of thick colorless oil.

5

^1H NMR

FD-MS, m/e 661 (M^+)

B) Preparation of D-Phe-Pro-NH(CH₂)₃NHC(NH)NH₂·HCl.

10

By a method substantially equivalent to that described in Example 1-H, 460 mg (71%) of D-Phe-Pro-NH(CH₂)₃NHC(NH)NH₂·HCl were prepared. The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 40/60 (A/B), 40 minutes).

15

^1H NMR

FD-MS, m/e 361 (M^+)


Analysis for C₁₈H₂₈N₆O₂·HCl·1.1TFA·1.1H₂O:

20

Calc: C, 44.66; H, 6.20; N, 15.47;

Found: C, 44.69; H, 6.10; N, 15.19.

Example 8



25

D-Phe-Pro-NHCH₂-trans-CH=CHCH₂NHC(NH)NH₂·HCl

A) Preparation of N⁹,N^{9'}-di-Boc-4-amino-trans-2-butenylguanidine.

30

By a method substantially equivalent to that described in Example 1-F, 2.4 g (42%) of N⁹,N^{9'}-di-Boc-4-amino-trans-2-butenylguanidine were prepared from 1,4-diamino-trans-2-butene.

5

B) Preparation of Boc-D-Phe-Pro-NHCH₂-trans-CH=CHCH₂NHC(NBoc)NHBoc.

By a method substantially equivalent to that described in Example 1-G, 2.7 g (55%) of Boc-D-Phe-Pro-NHCH₂-trans-CH=CHCH₂NHC(NBoc)NHBoc were prepared from Boc-D-Phe-Pro-OH and N⁹,N^{9'}-di-Boc-4-amino-trans-2-butenylguanidine.

¹H NMR

15 FD-MS, m/e 673 (M⁺)

C) Preparation of D-Phe-Pro-NHCH₂-trans-CH=CHCH₂NHC(NH)NH₂·HCl.

20 By a method substantially equivalent to that described in Example 1-H, approximately 100 mg of D-Phe-Pro-NHCH₂-trans-CH=CHCH₂NHC(NH)NH₂·HCl were prepared. The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 40/60 (A/B), 40 minutes).

25


¹H NMR

FD-MS, m/e 373 (M⁺)

Analysis for C₁₉H₂₈N₆O₂·HCl·0.5TFA·2.5H₂O:

Calc: C, 47.01; H, 6.81; N, 16.45;

30 Found: C, 47.36; H, 6.53; N, 16.70.

Example 9

5 D-Phe-Pro-p-NHCH₂C₆H₄CH₂NHC(NH)NH₂ · 2TFA

A) Preparation of p-H₂NCH₂C₆H₄CH₂NHC(NBoc)NHBoc.

By a method substantially equivalent to that described
 10 in Example 1-F, 2.3 g (42%) of
 p-H₂NCH₂C₆H₄CH₂NHC(NBoc)NHBoc were prepared from
 p-xylenediamine.

¹H NMR

15

B) Preparation of Boc-D-Phe-Pro-p-NHCH₂-
 C₆H₄CH₂NHC(NBoc)NHBoc.

By a method substantially equivalent to that described
 20 in Example 1-G, 2.8 g (63%) of Boc-D-Phe-Pro-p-NHCH₂-
 C₆H₄CH₂NHC(NBoc)NHBoc were prepared from Boc-D-Phe-Pro-OH
 and p-H₂NCH₂C₆H₄CH₂NHC(NBoc)NHBoc.

¹H NMR

25 FD-MS, m/e 723 (M⁺)

C) Preparation of D-Phe-Pro-p-NHCH₂-
 C₆H₄CH₂NHC(NH)NH₂ · 2TFA.

30 By a method substantially equivalent to that described
 in Example 1-H, 725 mg (81%) of the title bis-TFA salt were
 prepared and was not purified further by RPHPLC.


¹H NMR

FD-MS, m/e 423 (M⁺)

Analysis for C₂₃H₃₀N₆O₂·2.1TFA·H₂O:

5 Calc: C, 48.05; H, 5.05; N, 12.36;
 Found: C, 48.06; H, 4.85; N, 12.28.

Example 10



D-Phe-Pro-p-NHCH₂C₆H₄CH₂NH₂·HCl

A) Preparation of N-Boc-p-(aminomethyl)benzylamine.

15

To a stirring solution of p-xylenediamine (10 g, 73 mmol) in dimethylformamide/water (1:1, 100 mL) was added di-t-butyl dicarbonate (8 g, 37 mmol). After stirring for 20 hours, the mixture was concentrated in vacuo and the residue was partitioned between diethyl ether (200 mL) and 1 N citric acid (200 mL). The aqueous phase was washed again with diethyl ether (200 mL), and then basified with solid sodium bicarbonate and saturated with solid sodium chloride. The aqueous phase was then extracted four times with ethyl acetate (200 mL). The combined ethyl acetate extracts were dried (MgSO₄), filtered and concentrated to give 2.1 g (24%) of a thick oil.

20

25

¹H NMR

30 FD-MS, m/e 237 (MH⁺)

Analysis for $C_{13}H_{20}N_2O_2$:

Calc: C, 66.07; H, 8.53; N, 11.85;

Found: C, 66.33; H, 8.44; N, 12.11.

5 B) Preparation of Boc-D-Phe-Pro-p-NHCH₂C₆H₄CH₂NHBoc.

By a method substantially equivalent to that described in Example 1-G, 1.1 g (63%) of Boc-D-Phe-Pro-p-NHCH₂-C₆H₄CH₂NHBoc were prepared from Boc-D-Phe-Pro-OH and N-Boc-p-(aminomethyl)benzyl amine.

¹H NMR

FD-MS, m/e 581 (M⁺)

Analysis for $C_{32}H_{44}N_4O_6$:

15 Calc: C, 66.19; H, 7.64; N, 9.65;

Found: C, 65.99; H, 7.63; N, 9.42.

C) Preparation of D-Phe-Pro-p-NHCH₂C₆H₄CH₂NH₂·HCl.

20 By a method substantially equivalent to that described in Example 1-H, about 100 mg of D-Phe-Pro-p-NHCH₂-C₆H₄CH₂NH₂·HCl were prepared. The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 40/60 (A/B), 40 minutes).

25


¹H NMR

FD-MS, m/e 381 (M⁺)

Analysis for $C_{22}H_{28}N_4O_2 \cdot HCl \cdot 1.1TFA \cdot H_2O$:

Calc: C, 51.87; H, 5.77; N 10.00;

30 Found: C, 51.78; H, 5.88; N 10.28.

Example 11

5 D-Phe-Pro-m-NHCH₂C₆H₄CH₂NH₂ · HCl

A) Preparation of N-Boc-m-(aminomethyl)benzylamine.

By a procedure substantially equivalent to that
 10 described in Example 10-A, 2.6 g (30%) of
 N-Boc-m-(aminomethyl)benzylamine were prepared from
 m-xylenediamine.

¹H NMR

15 FD-MS, m/e 237 (MH⁺)

Analysis for C₁₃H₂₀N₂O₂:

Calc: C, 66.07; H, 8.53; N, 11.85;

Found: C, 65.81; H, 8.48; N, 11.98.

20 B) Preparation of Boc-D-Phe-Pro-m-NHCH₂C₆H₄CH₂NHBoc.

By a method substantially equivalent to that described
 in Example 1-G, 1.6 g (95%) of Boc-D-Phe-Pro-m-NHCH₂-
 C₆H₄CH₂NHBoc were prepared from Boc-D-Phe-Pro-OH and N-Boc-
 25 m-(aminomethyl)benzylamine.

¹H NMR


FD-MS, m/e 581 (M⁺)

C) Preparation of D-Phe-Pro-m-NHCH₂C₆H₄CH₂NH₂·HCl.

By a method substantially equivalent to that described in Example 1-H, about 100 mg of D-Phe-Pro-m-NHCH₂-
5 C₆H₄CH₂NH₂·HCl were prepared.

¹H NMRFD-MS, m/e 381 (M⁺)Analysis for C₂₂H₂₈N₄O₂·HCl·TFA·H₂O:

10 Calc: C, 52.51; H, 5.87; N 10.21;
 Found: C, 52.13; H, 6.21; N 10.48.

Example 12D-Phe-Pro-NHCH₂-trans-4-(aminomethyl)cyclohexane·HCl

A) Preparation of N-Boc-trans-4-
20 (aminomethyl)cyclohexanecarboxylic acid.

To a solution of trans-4-(aminomethyl)cyclohexane-carboxylic acid (50 g, 318 mmol) in 1 N sodium hydroxide (334 mL, 334 mmol) and t-butanol (400 mL) was added a
25 solution of di-t-butyl dicarbonate (73 g, 334 mmol) in tetrahydrofuran (50 mL). After stirring for 20 hours, the solvents were removed in vacuo and the residue was partitioned between water (500 mL) and diethyl ether (250 mL). The aqueous phase was washed again with diethyl ether
30 (250 mL) and then acidified with solid citric acid, which resulted in the formation of a white precipitate. The

solid was filtered, washed twice with water (100 mL) and dried in vacuo to yield 48 g (59%) of white powder.

¹H NMR

5

B) Preparation of HOCH₂-trans-4-(N-Boc-aminomethyl)cyclohexane.

To a stirring solution of N-Boc-trans-4-(aminomethyl)cyclohexanecarboxylic acid (15 g, 58 mmol) in tetrahydrofuran (150 mL) at 0 °C, was added N-methylmorpholine (5.9 g, 58 mmol), followed by ethyl chloroformate (6.3 g, 58 mmol). After stirring for 30 minutes, sodium borohydride (6.5 g, 175 mmol) was added, and then methanol (300 mL) was added via an addition funnel over 5 minutes. The mixture was allowed to stir for 1 hour and then the solvents were removed in vacuo. The residue was dissolved in ethyl acetate (500 mL) and washed twice with 1 N citric acid (250 mL), once with water (100 mL), twice with saturated aqueous sodium bicarbonate (250 mL) and once with saturated aqueous sodium chloride (250 mL). The organic phase was dried (MgSO₄), filtered and concentrated to give 13 g (91%) of the title compound.

25 ¹H NMR

C) Preparation of NH₂CH₂-trans-4-(N-Boc-aminomethyl)cyclohexane.

To a stirring solution of HOCH₂-trans-4-(N-Boc-aminomethyl)cyclohexane (13 g, 53 mmol) and triphenylphosphine (21 g, 80 mmol) in tetrahydrofuran (300 mL) was added diethyl azodicarboxylate (13.9 g, 80 mmol), followed by a solution of diphenylphosphoryl azide (22 g, 80 mmol) in tetrahydrofuran (100 mL). After stirring for 16 hours, the solvents were removed in vacuo and the residue was chromatographed over silica gel, eluting with a

step gradient of ethyl acetate/hexanes (1:3) through ethyl acetate/hexanes (3:1). The product containing fractions (judged by TLC), were combined and concentrated to give 17.4 g of crude product (contaminated with higher R_f compound). The crude azide was dissolved in methanol (200 mL) and this solution was added to a stirring suspension of finely ground $\text{Na}_2\text{S} \cdot 9\text{H}_2\text{O}$ (51 g, 212 mmol) and triethylamine (1 g, 11 mmol) in methanol (100 mL). The resulting mixture was heated to reflux (16 hours), then cooled to room temperature and the solvents were removed *in vacuo*. The residue was diluted with water (250 mL) and acidified with solid citric acid. The aqueous phase was washed twice with ethyl acetate (250 mL), basified with solid sodium bicarbonate, and saturated with solid sodium chloride. The aqueous phase was then extracted three times with ethyl acetate (200 mL) and the combined extracts were dried (MgSO_4), filtered and concentrated to give 6.4 g (45%) of a thick oil.

^1H NMR

D) Preparation of N-Boc-D-Phe-Pro-NHCH₂-trans-4-(N-Boc-aminomethyl)cyclohexane.

By a method substantially equivalent to that described in Example 1-G, 4.5 g (74%) of N-Boc-D-Phe-Pro-NHCH₂-trans-4-(N-Boc-aminomethyl)cyclohexane were prepared from Boc-D-Phe-Pro-OH and NH₂CH₂-trans-4-(N-Boc-aminomethyl)cyclohexane.

^1H NMR

FD-MS, m/e 587 (M^+)

E) Preparation of D-Phe-Pro-NHCH₂-trans-4-(aminomethyl)cyclohexane·HCl.

By a method substantially equivalent to that described in Example 1-H, 588 mg (75%) of D-Phe-Pro-NHCH₂-trans-4-(aminomethyl)cyclohexane·HCl were prepared. In this case, the analytical RPHPLC of the intermediate TFA salt was very
5 clean, but the salt was hygroscopic. So the salt was dissolved in 0.1 N HCl (20 mL), the pH was adjusted to 5, and the sample was lyophilized again to yield the stable white solid, hydrochloride salt.

10 ¹H NMR

FD-MS, m/e 387 (M⁺)


Analysis for C₂₂H₃₄N₄O₂·HCl·TFA·2H₂O:

Calc: C, 50.30; H, 7.04; N, 9.78;

Found: C, 50.44; H, 7.20; N, 9.62.

15

Example 13



20

D-Phe-Pro-p-NHCH₂C₆H₄NH₂·HCl

A) Preparation of Boc-D-Phe-Pro-p-NHCH₂C₆H₄NH₂.

By a method substantially equivalent to that described
25 in Example 1-G, 8 g of Boc-D-Phe-Pro-p-NHCH₂C₆H₄NO₂ were prepared from Boc-D-Phe-Pro-OH and p-NO₂-benzylamine·HCl. The intermediate was dissolved in ethanol (250 mL) and heated to reflux. To this stirring solution was added a solution of Na₂S₂O₄ (12.3 g, 70 mmol) in water (125 mL).
30 After stirring at reflux for 2 hours, the solvents were removed in vacuo and the residue was partitioned between ethyl acetate (250 mL) and water (250 mL). The aqueous

phase was extracted again with ethyl acetate (250 mL) and the combined organic phase was dried (MgSO_4), filtered, and concentrated in vacuo to give 2.4 g (21%) of a light yellow solid.

5

^1H NMR

FD-MS, m/e 466 (M^+)

Analysis for $\text{C}_{26}\text{H}_{34}\text{N}_4\text{O}_4$:

Calc: C, 66.93; H, 7.34; N, 12.01;

10 Found: C, 66.69; H, 7.32; N, 12.28.

B) Preparation of D-Phe-Pro-p-NHCH₂C₆H₄NH₂·HCl.

By a method substantially equivalent to that described
15 in Example 1-H, 180 mg (60%) of D-Phe-Pro-p-NHCH₂-C₆H₄NH₂·HCl were prepared. The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 60/40 (A/B), 60 minutes).

20 ^1H NMR


FD-MS, m/e 366 (M^+)

Analysis for $\text{C}_{21}\text{H}_{26}\text{N}_4\text{O}_2 \cdot \text{HCl} \cdot 0.6\text{TFA} \cdot 0.5\text{H}_2\text{O}$:

Calc: C, 55.51; H, 6.00; N, 11.66;

25 Found: C, 55.16; H, 6.14; N, 11.57.

Example 14



30

D-Phe-Pro-p-NHCH₂CH₂C₆H₄NH₂·HCl

A) Preparation of Boc-D-Phe-Pro-p-NHCH₂CH₂C₆H₄NH₂.

By a method substantially equivalent to that described in Example 13-A, 3 g (23%) of Boc-D-Phe-Pro-p-NHCH₂CH₂-C₆H₄NH₂ were prepared from Boc-D-Phe-Pro-OH and p-NO₂-phenethylamine·HCl.

¹H NMR

FD-MS, m/e 480 (M⁺)

Analysis for C₂₇H₃₆N₄O₄:

10 Calc: C, 67.48; H, 7.55; N, 11.66;
 Found: C, 67.30; H, 7.54; N, 12.34.

B) Preparation of D-Phe-Pro-p-NHCH₂CH₂C₆H₄NH₂·HCl.

15 By a method substantially equivalent to that described in Example 1-H, 175 mg (58%) of D-Phe-Pro-p-NHCH₂CH₂-C₆H₄NH₂·HCl were prepared. The product was purified by RPHPLC (method 1, 98/2 (A/B), ramp to 60/40 (A/B), 60 minutes).

20


¹H NMR

FD-MS, m/e 380 (M⁺)

Analysis for C₂₂H₂₈N₄O₂·HCl·0.7TFA·0.7H₂O:

 Calc: C, 55.18; H, 6.15; N, 11.00;
25 Found: C, 55.12; H, 6.18; N, 10.99.

Example 15



30

D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂ · HCl

(D-phenylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide hydrochloride)

A) Preparation of p-(aminomethyl)benzonitrile·TFA.

5

To a stirring suspension of sodium hydride (2.2 g, 56 mmol, 60% dispersion in oil) in tetrahydrofuran (100 mL) was added 4-(bromomethyl)benzonitrile (10 g, 51 mmol). To this mixture was added slowly via an addition funnel a solution of di-t-butyl iminodicarboxylate (12.2 g, 56 mmol). After stirring for 16 hours, the mixture was diluted with diethyl ether (300 mL) and washed twice with water (150 mL). The organic phase was then dried (MgSO₄), filtered and concentrated. The resulting solid was then dissolved in a minimum amount of dichloromethane. Anisole (10 mL) was added, and the solution was cooled to 0 °C. The solution was then diluted with trifluoroacetic acid (200 mL) and allowed to stir for 1 hour. The solvent was then removed in vacuo and the oily residue was stirred vigorously with diethyl ether (100 mL) and after a few minutes, the product solidified. The precipitate was filtered, washed with diethyl ether, and dried in vacuo to give 11.3 g (90%) of white powder.

25 IR

¹H NMR

FD-MS, m/e 132 (M⁺)

B) Preparation of Boc-D-Phe-Pro-p-NHCH₂C₆H₄CN.

30

By a method substantially equivalent to that described in Example 1-G, 7.4 g (78%) of Boc-D-Phe-Pro-p-NHCH₂C₆H₄CN were prepared from Boc-D-Phe-Pro-OH and p-(aminomethyl)benzonitrile·TFA. In this case, the product was purified by recrystallization from diethyl ether.

35

IR

 ^1H NMRFD-MS, m/e 476 (M^+)5 C) Preparation of Boc-D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂.

Hydrogen sulfide gas was bubbled through a solution of Boc-D-Phe-Pro-p-NHCH₂C₆H₄CN (2 g, 4.2 mmol) in pyridine (25 mL) and triethylamine (2.5 mL) for 30 minutes. The
10 reaction vessel was then sealed and allowed to stand at room temperature for 2 days. The solution was then diluted with water (100 mL) and extracted twice with ethyl acetate (200 mL). The combined organic phase was washed twice with saturated aqueous sodium chloride, dried (MgSO₄), filtered
15 and concentrated *in vacuo*.

The residue was dissolved in acetone (50 mL), methyl iodide (10 mL) was added, and the solution was heated to reflux (2 hours). The solvents were removed *in vacuo*, the residue was dissolved in methanol (20 mL), NH₄OAc (712 mg,
20 9.2 mmol) was added, and the solution was heated to reflux (12 hours). The solvent was again removed *in vacuo*, the residue was dissolved in 1 N citric acid (100 mL) and the aqueous phase was washed twice with ethyl acetate (200 mL), then basified with solid sodium bicarbonate, saturated with
25 solid sodium chloride, and extracted twice with ethyl acetate (200 mL). The combined ethyl acetate extracts were dried (MgSO₄), filtered and concentrated to give 1.4 g (67%) of thick oil.

30 ^1H NMRFD-MS, m/e 494 (M^+)D) Preparation of D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl.

35 By a method substantially equivalent to that described in Example 1-H, 7.7 g (57%) of D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl were prepared. The product was

purified by RPHPLC (method 3, 98/2 (A/B), ramp to 70/30 (A/B), 300 minutes).

^1H NMR

5 FD-MS, m/e 394 (M^+)


Analysis for $\text{C}_{22}\text{H}_{27}\text{N}_5\text{O}_2 \cdot \text{HCl} \cdot 1.4\text{TFA} \cdot 0.5\text{H}_2\text{O}$:

Calc: C, 49.76; H, 5.12; N, 11.70;

Found: C, 49.75; H, 5.19; N, 11.58.

10

Example 16



D-Phe-Pro-m-NHCH₂C₆H₄C(NH)NH₂ · 0.75HCl

15

A) Preparation of m-(aminomethyl)benzonitrile · TFA.

By a method substantially equivalent to that described in Example 15-A, 10.8 g (86%) of m-(aminomethyl)-benzonitrile · TFA were prepared from m-(bromomethyl)-benzonitrile.

IR

^1H NMR

25 FD-MS, m/e 132 (M^+)

B) Preparation of Boc-D-Phe-Pro-m-NHCH₂C₆H₄CN.

By a method substantially equivalent to that described in Example 1-G, 7.5 g (79%) of Boc-D-Phe-Pro-m-NHCH₂C₆H₄CN were prepared from Boc-D-Phe-Pro-OH and

m-(aminomethyl)benzonitrile·TFA. In this case, the product was purified by recrystallization from diethyl ether.

IR

5 ¹H NMR

FD-MS, m/e 476 (M⁺)

Analysis for C₂₇H₃₂N₄O₄:

Calc: C, 68.05; H, 6.77; N, 11.76;

Found: C, 68.27; H, 6.82; N, 11.96.

10

C) Preparation of Boc-D-Phe-Pro-m-NHCH₂C₆H₄C(NH)NH₂.

By a method substantially equivalent to that described in Example 15-C, 1.1 g (53%) of Boc-D-Phe-Pro-m-NHCH₂C₆H₄C(NH)NH₂ were prepared.

FD-MS, m/e 494 (M⁺)

20

D) Preparation of D-Phe-Pro-m-NHCH₂C₆H₄C(NH)NH₂·0.75HCl.

By a method substantially equivalent to that described in Example 1-G, 0.65 g (63%) of D-Phe-Pro-m-NHCH₂C₆H₄C(NH)NH₂·0.75 HCl were prepared. The product was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 75/25 (A/B), 120 minutes).


FD-MS, m/e 394 (M⁺)

Analysis for C₂₂H₂₇N₅O₂·0.75HCl·1.2TFA·0.5H₂O:

30

Calc. C, 51.72; H, 5.33; N, 12.36; Cl, 4.69;

Found: C, 51.79; H, 4.93; N, 11.96; Cl, 4.82.

Example 17

5 D-hPro-Pro-p-NHCH₂C₆H₄C(NH)NH₂ · HCl
 (D-homopropyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-
 prolinamide hydrochloride)

A) Preparation of Cbz-D-hPro-OH.

10

D-hPro-OH (5.0 g, 38.7 mmol) was dissolved in tetrahydrofuran (100 mL) and water (30 mL). The pH of the solution was adjusted to 9.5 with 2 N sodium hydroxide and benzyl chloroformate (5.5 mL, 38.7 mmol) was added dropwise
 15 and the pH maintained at 9.5 with 2 N sodium hydroxide. The reaction was stirred for an additional 1 hour at room temperature. The organic solvent was evaporated in vacuo, diethyl ether (100 mL) and water (50 mL) was added to the residue. The aqueous layer separated, the pH of the
 20 solution was adjusted to 2.8 with 3 N hydrochloric acid and ethyl acetate (150 mL) was added. The organic layer was separated and dried (MgSO₄); the filtrate was concentrated in vacuo to give 9.6 g (95%) of a clear oil.

25 ¹H NMR

FD-MS, m/e 264 (MH⁺)

B) Preparation of Cbz-D-hPro-Pro-OH.

30

Cbz-D-hPro-OH (9.5 g, 36 mmol) was dissolved in ethyl acetate (100 mL) and the solution cooled to 0 °C. Added to the solution was 2,4,5-trichlorophenol (7.1 g, 36 mmol) and

1,3-dicyclohexylcarbodiimide (7.4 g, 36 mmol). The reaction was stirred for 1 hour at 0 °C and 1 hour at room temperature. The precipitate was filtered and the filtrate concentrated in vacuo to an oil. The oil was dissolved in
5 pyridine (100 mL), Pro-OH (4.2 g, 36 mmol), and triethylamine (5.0 mL, 36 mmol) were added. The reaction was stirred at room temperature (24 hours). The reaction solvent was removed in vacuo to an oil. The residue was dissolved in water (100 mL), diethyl ether (50 mL) was
10 added and the pH adjusted to 9.5 with 2 N sodium hydroxide. The aqueous layer extracted twice with diethyl ether. The aqueous layer separated, the pH adjusted to 2.8 with 3 N hydrochloric acid and ethyl acetate (150 mL) was added. The organic layer was separated, dried (MgSO₄), and the
15 filtrate evaporated in vacuo to an amorphous solid (11.4 g, 88%).

FD-MS 361 (M⁺);

20 Analysis for C₁₉H₂₄N₂O₅:

Calc: C, 63.32; H, 6.71; N, 7.77;

Found: C, 63.42; H, 6.84; N, 7.96.

C) Preparation of Cbz-D-hPro-Pro-p-NHCH₂C₆H₄CN.

25

By a method substantially equivalent to that described in Example 1-G, 2.2 g (84%) of Cbz-D-hPro-Pro-p-NHCH₂C₆H₄CN were prepared from Cbz-D-hPro-Pro-OH and p-NH₂CH₂C₆H₄CN·TFA.

30

¹H NMR

FD-MS, m/e 474 (M⁺)

Analysis for C₂₇H₃₀N₄O₄:

Calc: C, 68.34; H, 6.37; N, 11.81;

35 Found: C, 68.36; H, 6.47; N, 11.57.

D) Preparation of D-hPro-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl.

By a method substantially equivalent to that described in Example 15-C, Cbz-hPro-Pro-p-NHCH₂C₆H₄C(NH)NH₂ (28 mmol, 5 theoretical) was prepared. This crude material was then dissolved in acetic acid (350 mL) and HBr gas was bubbled through the solution for 30 minutes. After stirring for an additional 1 hour, the solvent was removed in vacuo and the residue was dissolved in water (200 mL) and washed twice 10 with ethyl acetate (100 mL). The aqueous phase was then adjusted to pH 4 with ion exchange resin (Bio Rad AG1-X8, in basic form) and lyophilized to give a fluffy white solid. The product was then redissolved in water (25 mL) and purified by preparative RPHPLC (method 3, 98/2 (A/B), 15 ramp to 70/30 (A/B), 300 minutes) to give 5 g (41%) of hPro-Pro-p-NHCH₂C₆H₄C(NH)NH₂·0.9HCl·0.9HBr·0.5H₂O.


¹H NMRFD-MS, m/e 357 (M⁺)20 Analysis for C₁₉H₂₇N₅O₂·0.9HCl·0.9HBr·0.5H₂O:

Calc: C, 48.34; H, 6.36; N, 14.83; Cl, 6.76; Br, 15.23;

Found: C, 48.66; H, 6.36; N, 14.62; Cl, 7.14; Br, 14.90.

Example 18

25

1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·2HCl

(N-[[4-(aminoiminomethyl)phenyl]methyl]-1-[[(4aS,8aS)-
30 decahydro-1(R)-isoquinolinyl]carbonyl]-L-prolinamide
dihydrochloride)

A) Preparation of Cbz-D-1-Piq-Pro-OH.

A solution of 1-isoquinolinecarboxylic acid (50 g, 0.288 mol) in EtOH (150 mL) and 60 mL of 5N HCl was reduced over 5% Rh/Al₂O₃ (14 g) at 52 bar (750 psi) of hydrogen in a high pressure apparatus at 50 °C for 17 hours. The reaction mixture was filtered through a pad of diatomaceous earth, and the filtrate was concentrated in vacuo. The solid was triturated with water, filtered and dried to give DL-perhydro-1-isoquinolinecarboxylic acid (DL-1-Piq-OH) (30 g, 48%) FD-MS 184 (MH⁺).

DL-1-Piq-OH (30.2 g, 137 mmol) was dissolved in tetrahydrofuran (150 mL) and water (150 mL). The pH of the solution was adjusted to 9.8 with 5N NaOH, and benzyl chloroformate (21.6 mL, 151 mmol) was added dropwise and the pH maintained at 9.5 with 2N NaOH. The reaction was stirred for an additional 2 hours at room temperature. The organic solvent was evaporated in vacuo, and diethyl ether (150 mL) and water (50 mL) were added to the residue. The aqueous layer was separated, the pH of the solution was adjusted to 2.5 with 5N HCl, and ethyl acetate (200 mL) was added. The organic layer was separated and dried (MgSO₄) and the filtrate was concentrated in vacuo to give a clear oil. The oil was dissolved in diethyl ether (150 mL) and the solution allowed to stand at room temperature (24 hours). The precipitate was filtered and dried to give 2-Cbz-DL-perhydro-1-isoquinolinecarboxylic acid (Cbz-DL-1-Piq-OH) (32 g, 75%) FD-MS 318 (MH⁺).

Cbz-DL-1-Piq-OH (31.8 g, 100 mmol) was dissolved in DMF (100 mL) and cooled to 0 °C. To the reaction was added proline t-butyl ester (17.1 g, 100 mmol), 1-hydroxybenzotriazole (13.5 g, 100 mmol), and DCC (20.6 g, 100 mmol). The reaction was stirred for 3 hours at 0 °C and 24 hours at room temperature. The reaction precipitate was filtered and the filtrate concentrated in vacuo to an oil. The oil was dissolved in EtOAc (200 mL) and water (100 mL). The organic layer was separated, and washed sequentially with

1N NaHCO₃, water, 1.5N citric acid, and water. The organic layer was dried (MgSO₄), and the filtrate evaporated to an oil which was dried to give 2-Cbz-DL-perhydro-1-isoquinoline-carbonyl-L-prolyl t-butyl ester (Cbz-DL-1-Piq-Pro-O-t-Bu) (47.0 g, 100%) FAB-MS 470 (MH⁺).

Cbz-DL-1-Piq-Pro-O-t-Bu (47.0 g, 100 mmol) was placed in a round bottom flask containing trifluoroacetic acid (100 mL), CH₂Cl₂ (35 mL), anisole (5 mL) and stirred at room temperature (1 hour). The reaction was concentrated in vacuo without heating, and diethyl ether (100 mL) and water (100 mL) were added. The pH of the solution was adjusted to 9.8 with 5N NaOH. The aqueous layer was separated, the pH of the solution was adjusted to 2.5 with 5N HCl, and ethyl acetate (200 mL) was added. The organic layer was separated and dried (MgSO₄) and the filtrate was concentrated in vacuo to give a clear oil. The oil was dissolved in diethyl ether (700 mL) and (L)-(-)- α -methylbenzylamine added to the solution. The solution was allowed to stand at room temperature for 5 days. The resulting solid was filtered and washed with diethyl ether. The filtrate was washed with 1.5N citric acid, and water. The organic layer was dried (MgSO₄), and the filtrate evaporated to an oil. The oil was dissolved in diethyl ether (400 mL) and allowed to stand at room temperature (48 hours). The resulting solid was filtered, washed with diethyl ether, and dried to give 2-Cbz-D-perhydro-1-isoquinolinecarbonyl-L-proline (Cbz-D-1-Piq-Pro-OH) (5.86 g, 36%) FAB-MS 415 (MH⁺); [α]_D = -34.2° (C= 0.5, MeOH).

30

B) Preparation of N-Boc-p-(aminomethyl)benzonitrile.

To a stirring suspension of sodium hydride (4.6 g, 115 mmol, 60% dispersion in oil) in tetrahydrofuran (150 mL) was added 4-(bromomethyl)benzonitrile (20.5 g, 105 mmol). To this mixture was added (slowly via an addition funnel) a solution of di-t-butyl iminodicarboxylate (25 g, 115 mmol).

After stirring for 16 hours, the mixture was diluted with diethyl ether (500 mL) and washed twice with water (250 mL). The organic phase was then dried (MgSO_4), filtered and concentrated to give 40.2 g of crude solid.

5 The resulting solid (28.3 g, 85 mmol) was then dissolved in tetrahydrofuran (150 mL) and a solution of sodium hydroxide (3.4 g, 85 mmol) in methanol (300 mL) was added. After stirring overnight, the solution was concentrated to about one-half volume and water was added
10 to promote precipitation of the product. The precipitate was filtered and dried *in vacuo* to give 18.5 g (94%) of a white solid.

IR

15 ^1H NMR

FD-MS, m/e 232 (M^+)

Analysis for $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_2$:

Calc: C, 67.22; H, 6.94; N, 12.06;

Found: C, 67.19; H, 7.16; N, 11.82.

20

C) Preparation of $p\text{-(BocNHCH}_2\text{)C}_6\text{H}_4\text{C(NH)NHCBz}$.

By a method substantially equivalent to that described in Example 15-C, N-Boc-p-(aminomethyl)benzonitrile (32.7 g,
25 140 mmol) was elaborated to $p\text{-(BocNHCH}_2\text{)C}_6\text{H}_4\text{C(NH)NH}_2$. The residue from this procedure was dissolved in dimethylformamide (700 mL) and N,N-diisopropylethylamine (72 g, 560 mmol) was added. To this stirring solution was added dropwise benzyl chloroformate (48 g, 280 mmol).
30 After stirring for 16 hours, water (100 mL) was added and then the solvents were removed *in vacuo*. The residue was partitioned between water (250 mL) and ethyl acetate (500 mL). The phases were separated and the organic phase was washed three times with saturated aqueous ammonium chloride
35 (250 mL), once with water (200 mL) and twice with saturated aqueous sodium bicarbonate (250 mL). The organic phase was then dried (MgSO_4), filtered and concentrated and the

product was recrystallized from diethyl ether to give 14 g (26%) of white solid.

¹H NMR

5 FD-MS, m/e 384 (M⁺)

D) Preparation of p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

To a solution of p-(BocNHCH₂)C₆H₄C(NH)NHCbz (11 g, 28.7 mmol) in dichloromethane (125 mL) at 0 °C, was added anisole (10 mL) followed by trifluoroacetic acid (125 mL). After stirring for 2 hours, the solvents were removed in vacuo and the residue was dissolved in 1 N hydrochloric acid (50 mL) and washed twice with diethyl ether (50 mL). The pH was adjusted to 3 with ion exchange resin (Bio Rad AG1-X8, in basic form) and the solution was lyophilized to give 9.2 g (90%) of white powder.

¹H NMR

20 FD-MS, m/e 284 (M⁺)

E) Preparation of Cbz-1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NHCbz.

By a method substantially equivalent to that described in Example 1-G, 4.4 g (79%) of Cbz-1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NHCbz were prepared from Cbz-1-Piq-Pro-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl. In this case, the reaction was performed in dimethylformamide due to solubility problems with p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

¹H NMR

FD-MS, m/e 681 (MH⁺)

Analysis for C₃₉H₄₅N₅O₆:

Calc: C, 68.91; H, 6.67; N, 10.30;

Found: C, 68.71; H, 6.93; N, 10.38.

F) Preparation of 1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·2HCl.


To a solution of Cbz-1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NHCbz (4.2 g, 6.1 mmol) in ethanol (200 mL), was added 1 N hydrochloric acid (18.3 mL, 18.3 mmol) and water (100 mL). To this stirring solution was added 5% Pd/C (1 g), and hydrogen gas was bubbled through the solution for a period of 2 hours. The mixture was then flushed with nitrogen, and then filtered through a pad of diatomaceous earth. The filtrate was then concentrated in vacuo, redissolved in water (25 mL) and purified by RPHPLC (method 2, 98/2 (A/B), ramp to 60/40 (A/B), 300 minutes), to give 1.3 g (53%) of 1-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·2HCl.

15 ¹H NMRFD-MS, m/e 412 (M⁺)Analysis for C₂₃H₃₃N₅O₂·1.9HCl·2.5H₂O:

Calc: C, 52.53; H, 7.65; N, 13.32; Cl, 12.81;

Found: C, 52.63; H, 7.36; N, 13.47; Cl, 12.95.

20

Example 19

25

D-3-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·3HCl

A) Preparation of Cbz-D-3-Piq-Pro-OH.

D-Phenylalanine (50 g, 302 mmol) was reacted with a 37% solution of formaldehyde (120 mL) and concentrated HCl (380 mL) at reflux. After 30 minutes an additional 50 mL of formaldehyde was added and reaction continued for 3 hours.

The reaction cooled to -10 °C and the precipitate was filtered. The solid was dried in vacuo to give D-1,2,3,4-tetrahydro-3-isoquinolinecarboxylic acid (24.2 g, 45%) FD-MS 178 (MH⁺).

5 A solution of D-1,2,3,4-tetrahydro-3-isoquinoline-carboxylic acid (17 g, 96 mmol) in water (200 mL) and 20 mL of 5N HCl was hydrogenated over 5% Rh/Al₂O₃ (8.5 g) at 138 bar (2000 psi) in a high pressure apparatus at 120 °C for 16 hours. The reaction mixture was filtered through a pad of
10 diatomaceous earth and the filtrate was freeze-dried to give D-perhydro-3-isoquinolinecarboxylic acid (D-3-Piq-OH) (21 g, 100%) FD-MS 184 (MH⁺).

 D-3-Piq-OH (21.0 g, 95.8 mmol) was dissolved in tetrahydrofuran (75 mL) and water (50 mL). The pH of the
15 solution was adjusted to 10.0 with 5N NaOH, and benzyl chloroformate (16.4 mL, 115 mmol) was added dropwise and the pH maintained at 9.5 with 2N NaOH. The reaction was stirred for an additional 1 hour at room temperature. The organic solvent was evaporated in vacuo, and diethyl ether (100 mL)
20 and water (50 mL) were added to the residue. The aqueous layer was separated, the pH of the solution was adjusted to 3.0 with 3N HCl, and ethyl acetate (250 mL) was added. The organic layer was separated and dried (MgSO₄). The filtrate was concentrated in vacuo to give a clear oil of 2-Cbz-
25 D-perhydro-3-isoquinolinecarboxylic acid (Cbz-D-3-Piq-OH) (25.8 g, 85%) FD-MS 318 (MH⁺).

 Cbz-D-3-Piq-OH (17.2 g, 54 mmol) was dissolved in DMF (50 mL) and cooled to 0 °C. To the reaction was added proline t-butyl ester (9.2 g, 54 mmol), 1-hydroxybenzo-
30 triazole (7.3 g, 54 mmol), and DCC (11.1 g, 54 mmol). The reaction was stirred for 3 hours at 0 °C and 24 hours at room temperature. The reaction precipitate was filtered and the filtrate concentrated in vacuo to an oil. The oil was dissolved in EtOAc (200 mL) and water (100 mL). The organic
35 layer was separated, and washed sequentially with 1N NaHCO₃, water, 1.5N citric acid, and water. The organic layer was dried (MgSO₄), and the filtrate evaporated to an oil which

was dried to give 2-Cbz-D-perhydro-3-isoquinolinecarbonyl-L-proline t-butyl ester (Cbz-D-3-Piq-Pro-O-Bu) (23.8 g, 94%) FAB-MS 471 (MH⁺).

5 Cbz-D-3-Piq-Pro-O-t-Bu (31.2 g, 66.3 mmol) was placed in a round bottom flask containing trifluoroacetic acid (100 mL), anisole (5 mL), and stirred at room temperature (1 hour). The reaction was concentrated in vacuo without heating, and diethyl ether (150 mL), and water (100 mL) were added. The pH of the solution was adjusted to 9.8 with 5N
10 NaOH. The aqueous layer was separated, the pH of the solution was adjusted to 2.8 with 3N HCl, and ethyl acetate (200 mL) was added. The organic layer was separated, dried (MgSO₄), and filtered. The filtrate was concentrated in vacuo to give a clear oil. The oil was dissolved in diethyl
15 ether (300 mL) and the solution was allowed to stand at room temperature (24 hours). The resulting solid was filtered, washed with diethyl ether, and dried to give 2-Cbz-perhydro-3-isoquinolinecarbonyl-L-proline (Cbz-D-3-Piq-Pro-OH) (13.5 g, 49%) FAB-MS 415 (MH⁺).

20 Analysis for C₂₃H₃₀N₂O₅:

Calc: C, 66.65; H, 7.29; N, 6.76;

Found: C, 66.90, H, 7.33, N, 6.81.

B) Preparation of Cbz-D-3-Piq-Pro-p-NHCH₂C₆H₄C(NH)NHCbz.

25

By a method substantially equivalent to that described in Example 18-E, 1.6 g (49%) of Cbz-D-3-Piq-Pro-p-NHCH₂-C₆H₄C(NH)NHCbz were prepared from Cbz-D-3-Piq-Pro-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

30

FD-MS, m/e 680 (M⁺)

C) Preparation of D-3-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·3HCl.

35

By a method substantially equivalent to that described in Example 17-C, 150 mg of D-3-Piq-Pro-p-NHCH₂C₆H₄C(NH)NH₂·3HCl were prepared. The product was

purified by RPHPLC (method 2, 98/2 (A/B), ramp to 60/40 (A/B), 240 minutes).

^1H NMR

5 FD-MS, m/e 412 (M^+)


Analysis for $\text{C}_{23}\text{H}_{33}\text{N}_5\text{O}_2 \cdot 3\text{HCl} \cdot 0.5\text{H}_2\text{O}$:

Calc: C, 52.13; H, 7.04; N, 13.22;

Found: C, 52.35; H, 7.23; N, 12.95.

10

Example 20



D-hPro-Ohi-p-NHCH₂C₆H₄C(NH)NH₂ · 3HCl

15 ((S-cis)-N-[[4-(aminoiminomethyl)phenyl]methyl]-octahydro-1-D-homopropyl-1H-indole-2-carboxamide trihydrochloride)

A) Preparation of Cbz-D-hPro-Ohi-OH.

20 HCl gas was bubbled through a stirring suspension of (S)-indoline-2-carboxylic acid (20 g, 110 mmol) in ethanol (500 mL). When the acid was completely dissolved, the solution was brought to reflux. After 16 hours, the solution was cooled and the solvent removed in vacuo. The residue was
25 triturated with diethyl ether and the resulting off-white solid was collected by filtration, washed with hexanes and dried overnight in a vacuum oven at 30 °C to give (S)-indoline-2-carboxylic acid ethyl ester · HCl (25.7 g, 78%).

The solid was dissolved in ethanol (800 mL), 5% Pd/C (25
30 g) was added, and the resulting suspension was hydrogenated on a Parr shaker for 8 hours (4.1 bar, 60 psi). The solution was filtered and the solvent was removed in vacuo. The

residue was dissolved triturated with diethyl ether and 18.8 g (73%) of an off white solid (cis-Ohi-OEt·HCl) were collected by filtration.

By a method substantially equivalent to that described in Example 1-A, 13.5 g (93%) of Cbz-D-hPro-cis-Ohi-OEt were prepared from Cbz-D-hPro-OH and cis-Ohi-OEt·HCl.

¹H NMR

FD-MS, m/e 442 (M⁺)

10 Analysis for C₂₅H₃₄N₂O₅:

Calc: C, 67.85; H, 7.74; N, 6.33;

Found: C, 67.59; H, 7.72; N, 6.48.

By a method substantially equivalent to that described in Example 1-D, 12.5 g (102%) of Cbz-D-hPro-cis-Ohi-OH were prepared.

¹H NMR

FD-MS, m/e 414 (M⁺)

20 Analysis for C₂₃H₃₀N₂O₅:

Calc: C, 66.65; H, 7.29; N, 6.76;

Found: C, 66.46; H, 7.30; N, 6.86.

B) Preparation of Cbz-D-hPro-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz.

25

By a method substantially equivalent to that described in Example 18-E, 3.3 g (67%) of Cbz-D-hPro-Ohi-p-NHCH₂-C₆H₄C(NH)NHCbz were prepared from Cbz-D-hPro-Ohi-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

30

¹H NMR

FD-MS, m/e 681 (MH⁺)

C) Preparation of D-hPro-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·3HCl.

35

By a method substantially equivalent to that described in Example 18-F, 2.2 g (66%) of D-hPro-Ohi-p-NHCH₂-

$C_6H_4C(NH)NH_2 \cdot 3HCl$ were prepared. The product was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 60/40 (A/B), 3000 minutes).

5 1H NMR

FD-MS, m/e 412 (M^+)


Analysis for $C_{23}H_{33}N_5O_2 \cdot 3HCl \cdot 0.5H_2O$:

Calc: C, 52.13; H, 7.04; N, 13.22;

Found: C, 51.98; H, 7.04; N 13.35.

10

Example 21



15

D-hPro-N(PhCH₂CH₂)Gly-p-NHCH₂C₆H₄C(NH)NH₂ · 2HCl
(D-homoprolyl-N(α)-(2-phenylethyl)-N-[[4-(aminoiminomethyl)phenyl]methyl]glycinamide dihydrochloride)

20

A) Preparation of Cbz-D-hPro-N(PhCH₂CH₂)Gly-OH.

To a solution of phenethylamine (58 mL, 461 mmol) and triethylamine (21 mL, 154 mmol) in ethanol (200 mL) at 0 °C was added a solution of t-butyl bromoacetate (30 g, 154 mmol) in ethanol (50 mL) over 1 hour. The cold bath was left
25 unattended and the solution was allowed to warm to room temperature. After stirring overnight, the solvents were removed in vacuo and the residue was dissolved in 1N citric acid. The aqueous solution was washed twice with diethyl
30 ether, basified with solid sodium bicarbonate, and then extracted three times with ethyl acetate (200 mL). The

combined ethyl acetate extracts were dried (MgSO_4), filtered and allowed to stand for 24 hours. The resulting precipitate was filtered, washed with diethyl ether, and dried to give 10.5 g of white solid. The mother liquor was concentrated to a volume of about 100 mL and then diluted with diethyl ether (400 mL). After standing for 30 minutes, the solution was filtered to yield an additional 23.5 g of white solid for a combined total of 34 g (94%) of $\text{N}(\text{PhCH}_2\text{CH}_2)\text{Gly-O-t-Bu}$.

10 ^1H -NMR

FD-MS, m/e 235 (M^+)

By a method substantially equivalent to that described in Example 1-A, 10.8 g (56%) of $\text{Cbz-D-hPro-N}(\text{PhCH}_2\text{CH}_2)\text{Gly-O-t-Bu}$ were prepared from Cbz-D-hPro-OH and $\text{N}(\text{PhCH}_2\text{CH}_2)\text{Gly-O-t-Bu}$.

^1H NMR

FD-MS, m/e 480 (M^+)

20 Analysis for $\text{C}_{28}\text{H}_{36}\text{N}_2\text{O}_5$:

Calc: C, 69.98; H, 7.55; N, 5.83;

Found: C, 69.68; H, 7.56; N, 5.77.

By a method substantially equivalent to that described in Example 18-A, for the deprotection of $\text{Cbz-DL-1-Piq-Pro-O-t-Bu}$, 9.2 g (100%) of $\text{Cbz-D-hPro-N}(\text{PhCH}_2\text{CH}_2)\text{Gly-OH}$ were prepared.

^1H NMR

30 FD-MS, m/e 425 (M^+)

Analysis for $\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_5$:

Calc: C, 67.91; H, 6.65; N, 6.60;

Found: C, 68.19; H, 6.68; N, 6.71.

35 B). Preparation of $\text{Cbz-D-hPro-N}(\text{PhCH}_2\text{CH}_2)\text{Gly-p-NHCH}_2\text{-C}_6\text{H}_4\text{C}(\text{NH})\text{NHCbz}$.

By a method substantially equivalent to that described in Example 18-E, 3.2 g (55%) of Cbz-D-hPro-N(PhCH₂CH₂)Gly-p-NHCH₂C₆H₄C(NH)NHCbz were prepared from Cbz-D-hPro-N(PhCH₂CH₂)Gly-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

5

¹H NMRFD-MS, m/e 690 (M⁺)Analysis for C₄₀H₄₃N₅O₆:

Calc: C, 69.65; H, 6.28; N, 10.15;

10 Found: C, 69.80; H, 6.46; N, 10.14.

C) Preparation of D-hPro-N(PhCH₂CH₂)Gly-p-NHCH₂-C₆H₄C(NH)NH₂·2HCl.

15


By a method substantially equivalent to that described in Example 19-F, 770 mg (54%) of D-hPro-N(PhCH₂CH₂)Gly-p-NHCH₂C₆H₄C(NH)NH₂·2HCl were prepared. The product was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 85/15 (A/B), 120 minutes).

20

¹H NMRFD-MS, m/e 423 (MH⁺)Analysis for C₂₄H₃₁N₅O₂·2HCl:

Calc: C, 58.30; H, 6.73; N, 14.16;

25 Found: C, 58.05; H, 6.60; N, 14.28.

Example 22

D-hPro-Pro(4-cis-PhO)-p-NHCH₂C₆H₄C(NH)NH₂·2HCl
(cis-D-homopropyl-N-[[4-(aminoiminomethyl)-phenyl]methyl]-
4-phenoxy-L-prolinamide dihydrochloride)

5

A) Preparation of Cbz-D-hPro-Pro(4-cis-PhO)-OH.

To a solution of Cbz-Pro(4-trans-OH)-Et (58.8 g, 200 mmol), triphenylphosphine (65.6 g, 250 mmol), and phenol
10 (23.5 g, 250 mmol) in tetrahydrofuran (500 mL) at 0 °C, was added (dropwise over 1 hour) a solution of diethyl azodicarboxylate (40 mL, 250 mmol) in tetrahydrofuran (50 mL). The cold bath was then removed and the solution was allowed to warm to room temperature (16 hours). The solvent
15 was then removed in vacuo and the remaining amber syrup was triturated with diethyl ether. The white solid was removed by filtration and the filtrate was concentrated. The residue was then chromatographed over silica gel (1 Kg), eluting with a step gradient from hexanes through 1:1 ethyl
20 acetate/hexanes. The fractions containing pure product (as judged by TLC) were combined and concentrated in vacuo to give 36.3 g (50%) of Cbz-Pro(4-cis-phenoxy)-OEt as a colorless syrup.

25 ¹H NMRFD-MS, m/e 369 (M⁺)Analysis for C₂₁H₂₃NO₅:

Calc: C, 68.28; H, 6.28; N, 3.79;

Found: C, 68.38; H, 6.30; N, 3.89.

30

To a solution of Cbz-Pro(4-cis-phenoxy)-OEt (25 g, 67.7 mmol) in ethanol (400 mL) was added 5% Pd/C (5 g). After bubbling hydrogen through the solution for 3 hours, the solution was filtered through a pad of diatomaceous earth, 3
35 mL of concentrated hydrochloric acid were added, and the solution was concentrated in vacuo. The residue was suspended in diethyl ether with vigorous stirring and then

filtered and dried to give 14.2 g (77%) of Pro(4-cis-phenoxy)-OEt·HCl as a white solid.

¹H NMR

5 FD-MS, m/e 235 (M⁺)

Analysis for C₁₃H₁₈NO₃Cl:

Calc: C, 57.46; H, 6.68; N, 5.15;

Found: C, 57.68; H, 6.78; N, 5.18.

10 By a method substantially equivalent to that described in Example 1-A, 19.4 g (100%) of Cbz-D-hPro-Pro(4-cis-phenoxy)-OEt were prepared from Cbz-D-hPro-OH and Pro(4-cis-phenoxy)-OEt·HCl.

15 ¹H NMR

FD-MS, m/e 480 (M⁺)

Analysis for C₂₇H₃₂N₂O₆:

Calc: C, 67.48; H, 6.71; N, 5.83;

Found: C, 67.71; H, 6.79; N, 5.89.

20

By a method substantially equivalent to that described in Example 1-D, 16 g (100%) of Cbz-D-hPro-Pro(4-cis-phenoxy)-OH were prepared.

25 ¹H NMR

FD-MS, m/e 452 (M⁺)

Analysis for C₂₅H₂₈N₂O₆:

Calc: C, 66.36; H, 6.24; N, 6.19;

Found: C, 66.22; H, 6.18; N, 6.17.

30

B) Preparation of Cbz-D-hPro-Pro(4-cis-PhO)-p-NHCH₂-C₆H₄C(NH)NHCbz.

35 By a method substantially equivalent to that described in Example 18-E, 4.55 g (75%) of Cbz-D-hPro-Pro(4-cis-PhO)-p-NHCH₂C₆H₄C(NH)NHCbz were prepared from Cbz-D-hPro-Pro(4-cis-PhO)-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

¹H NMR

FD-MS, m/e 718 (M⁺)

- 5 C) Preparation of D-hPro-Pro(4-cis-PhO)-p-NHCH₂-C₆H₄C(NH)NH₂·2HCl.

By a method substantially equivalent to that described in Example 18-F, 873 mg (40%) of D-hPro-Pro(4-cis-PhO)-p-NHCH₂C₆H₄C(NH)NH₂·2HCl were prepared. The product was
 10 purified by RPHPLC (method 2, 98/2 (A/B), ramp to 85/15 (A/B), 120 minutes).

¹H NMR

15 FD-MS, m/e 451 (MH⁺)


Analysis for C₂₅H₃₁N₅O₃·2HCl:

Calc: C, 57.47; H, 6.37; N, 13.40;

Found: C, 57.22; H, 6.29; N, 13.47.

20

Example 23



EtSO₂-D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl

25

(N-(ethylsulfonyl)-D-phenylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide hydrochloride)

- A) Preparation of p-NH₂CH₂-C₆H₄CN·HCl.

30

HCl gas was bubbled through a stirring solution of N-Boc-p-aminomethylbenzonitrile (15 g, 64.6 mmol) in ethyl

acetate (400 mL) at 0 °C for 10 minutes. The cold bath was removed and after stirring for 1.5 hours, the solvent was removed in vacuo and the residue was suspended in diethyl ether, filtered, washed with diethyl ether and dried to give
5 10.1 g (93%) of white solid.

IR

¹H NMR

FD-MS, m/e 132 (M⁺)

10 Analysis for C₈H₉N₂Cl:

Calc: C, 56.98; H, 5.38; N, 16.61; Cl, 21.02;

Found: C, 56.36; H, 5.46; N, 16.22; Cl, 21.31.

B) Preparation of EtSO₂-D-Phe-Pro-p-NHCH₂-C₆H₄CN.

15

By a method substantially equivalent to that described in Example 1-G, 1.5 g (80%) of EtSO₂-D-Phe-Pro-p-NHCH₂-C₆H₄CN were prepared from EtSO₂-D-Phe-Pro-OH and p-NH₂CH₂-C₆H₄CN·HCl.

20 IR

¹H NMR

FD-MS, m/e 468 (M⁺)

Analysis for C₂₄H₂₈N₄O₄S:

Calc: C, 61.52; H, 6.02; N, 11.90;

25 Found: C, 61.23; H, 6.13; N, 11.80.

C) Preparation of EtSO₂-D-Phe-Pro-p-NHCH₂-C₆H₄C(=NOH)NH₂·HCl.

To a solution of EtSO₂-D-Phe-Pro-p-NHCH₂-C₆H₄CN (1 g, 2.1
30 mmol) in absolute ethanol (35 mL) was added
N,N-diisopropylethylamine (0.47 mL, 2.7 mmol) followed by
hydroxylamine hydrochloride (185 mg, 2.7 mmol) and the
solution was brought to reflux. After 16 hours, the solution
was cooled and the solvents were removed in vacuo. 250 mg of
35 this material were taken on to the next step and the
remaining material was purified by RPHPLC (method 1, 90/10
(A/B); ramp to 60/40 (A/B) over 200 minutes).

IR

 ^1H NMRFD-MS, m/e 501 (M^+)Analysis for $\text{C}_{24}\text{H}_{31}\text{N}_5\text{O}_5\text{S} \cdot 1.2\text{HCl} \cdot \text{H}_2\text{O}$:

- 5 Calc: C, 51.17; H, 6.12; N, 12.42; Cl, 7.55;
 Found: C, 51.04; H, 5.81; N, 12.39; Cl, 7.18.

D) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2 \cdot \text{HCl}$.

- 10 To a solution of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2 \cdot \text{HCl}$ (250 mg, 0.52 mmol) in ethanol (40 mL) and water (19 mL) was added 1 N HCl (1 mL) followed by 5% Pd on carbon (250 mg). The stirring suspension was placed under an atmosphere of hydrogen for 18 hours and was then filtered,
 15 concentrated and purified by RPHPLC (method 1, 90/10 (A/B), ramp to 60/40 (A/B) over 200 minutes) to give 140 mg (52%) of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2 \cdot \text{HCl}$.

 ^1H NMR

- 20 FD-MS, m/e 486 (M^+)


Analysis for $\text{C}_{24}\text{H}_{31}\text{N}_5\text{O}_4\text{S} \cdot \text{HCl} \cdot 1.5\text{H}_2\text{O}$:

- Calc: C, 52.50; H, 6.42; N, 12.75;
 Found: C, 52.56; H, 6.19; N, 12.59.

- 25 Another 5 g of product were prepared by the methods described in Example 15 and purified by RPHPLC (method 3; 98/2 (A/B), 60 minutes, ramp to 60/40 (A/B) 300 minutes).

Example 24

30

 $\text{EtSO}_2\text{-D-Phe-Pro-m-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2 \cdot \text{HCl}$

A) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-m-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2$.

By methods substantially equivalent to those described in Example 23, $\text{EtSO}_2\text{-D-Phe-Pro-m-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2$ was prepared using $\text{m-BrCH}_2\text{-C}_6\text{H}_4\text{CN}$ in place of $\text{p-BrCH}_2\text{-C}_6\text{H}_4\text{CN}$. 140 mg (13%) of this crystalline intermediate was held back and the rest of the material was taken on to step B.

 ^1H NMRFD-MS, m/e 502 (M^+)Analysis for $\text{C}_{24}\text{H}_{31}\text{N}_5\text{O}_5\text{S}$:

Calc: C, 57.47; H, 6.23; N, 13.96;

Found: C, 57.28; H, 6.21; N, 13.66.


B) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-m-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2\cdot\text{HCl}$.

By a method substantially equivalent to that described in Examples 23-C and 23-D, 0.27 g (28%, 2 steps) of $\text{EtSO}_2\text{-D-Phe-Pro-m-NHCH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2\cdot\text{HCl}$ were prepared.

 ^1H NMRFD-MS, m/e 486 (M^+)Analysis for $\text{C}_{24}\text{H}_{31}\text{N}_5\text{O}_4\text{S}\cdot 1.1\text{HCl}\cdot 2\text{H}_2\text{O}$:

Calc: C, 51.32; H, 6.48; N, 12.47; Cl, 6.94;

Found: C, 51.33; H, 6.09; N, 12.20; Cl, 6.66.

Example 25D-1-Piq-Pro-m-NHCH₂-C₆H₄C(=NH)NH₂·HCl

By a method substantially equivalent to that described in Example 23, 0.86 g of D-1-Piq-Pro-m-NHCH₂-C₆H₄C(=NH)NH₂·HCl were prepared from Cbz-D-1-Piq-Pro-OH and m-NH₂CH₂-C₆H₄CN·HCl.

5


¹H NMR

FD-MS, m/e 412 (M⁺)

Analysis for C₂₃H₃₃N₅O₂·2.5HCl·0.5H₂O:

10 Calc: C, 53.99; H, 7.19; N, 13.69;
 Found: C, 54.19; H, 7.02; N, 13.81.

Example 26



15

EtSO₂-D-Phe-Pro-p-NHCH₂CH₂-C₆H₄C(=NH)NH₂·HCl

A) Preparation of methyl p-cyano-trans-cinnamate.

20

To a stirring suspension of NaH (6.1 g of 60% oil suspension, 153 mmol) and p-cyanobenzaldehyde (20 g, 153 mmol) in tetrahydrofuran (250 mL) at 0 °C was added via addition funnel a solution of trimethyl phosphonoacetate (28 g, 153 mmol) in tetrahydrofuran (50 mL). After stirring for 48 hours, the solvent was removed in vacuo and the crude residue was dissolved in ethyl acetate (500 mL). The ethyl acetate solution was washed once with water, three times with saturated aqueous NaHSO₃ and once with brine. The organic

25

phase was then dried with MgSO_4 , filtered and concentrated in vacuo to give 28 g (98%) of a white solid.

IR

5 ^1H NMR

FD-MS, m/e 187 (M^+)

B) Preparation of methyl p-cyano-dihydrocinnamate.

10 To a solution of methyl p-cyano-trans-cinnamate (13.6 g, 73 mmol) in toluene (485 mL) was added 5% Pd/ BaSO_4 (2.7 g). After 9 hours under hydrogen gas at 4 bar (60 psi), the solution was filtered, concentrated in vacuo and chromatographed over silica gel, eluting with a step gradient
15 of hexanes through 30% ethyl acetate/hexanes. The product containing fractions were combined and concentrated to give 10.6 g (77%) of colorless oil.

IR

20 ^1H NMR

FD-MS, m/e 189 (M^+)

C) Preparation of p-cyano-dihydrocinnamic acid.

25 By methods substantially equivalent to those described in Example 1-D, using 1.1 equivalent of $\text{LiOH}\cdot\text{H}_2\text{O}$, 5.1 g (58%) of p-cyano-dihydrocinnamic acid were prepared from methyl p-cyano-dihydrocinnamate.

30 IR

^1H NMR

FD-MS, m/e 175 (M^+)

D) Preparation of Boc-p-NHCH₂CH₂-C₆H₄CN.

35

To a solution of p-cyano-dihydrocinnamic acid (6.7 g, 38.2 mmol) and triethylamine (5.9 mL, 42 mmol) in t-butanol

(150 mL) was added diphenylphosphoryl azide (11.6 g, 42 mmol) and the solution was brought to reflux. After stirring overnight, the solution was cooled and the solvent was removed in vacuo. The residue was dissolved in ethyl acetate and washed three times with 1 N citric acid, once with brine, twice with saturated aqueous NaHCO_3 , and then dried (MgSO_4), filtered and concentrated in vacuo. The residue was then chromatographed over silica gel, eluting with 10% ethyl acetate/hexanes through 50% ethyl acetate/hexanes. The product containing fractions as judged by TLC were combined and concentrated to give 5.4 g (57%) of white solid.

IR

^1H NMR

15 FD-MS, m/e 246 (M^+)

Analysis for $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}_2$:

Calc: C, 68.27; H, 7.37; N, 11.37;

Found: C, 68.39; H, 7.50; N, 11.40.

20 E) Preparation of $\text{p-NH}_2\text{CH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}\cdot\text{HCl}$.

By methods substantially equivalent to those described in Example 23-A, 3.6 g (98%) of $\text{p-NH}_2\text{CH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}\cdot\text{HCl}$ were prepared.

25

^1H NMR

FD-MS, m/e 147 (MH^+)

Analysis for $\text{C}_9\text{H}_{11}\text{N}_2\text{Cl}$:

Calc: C, 59.18; H, 6.07; N, 15.34; Cl, 19.41;

30 Found: C, 58.90; H, 6.16; N, 15.20; Cl, 19.30.

F) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}$.

By methods substantially equivalent to those described in Example 1-G, 1.5 g of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}$ were prepared from $\text{EtSO}_2\text{-D-Phe-Pro-OH}$ and $\text{p-NH}_2\text{CH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}\cdot\text{HCl}$.

IR

 ^1H NMRFD-MS, m/e 482 (M^+)

5

G) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2\cdot\text{HCl}$.

To a stirring solution of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{CN}$ (1 g, 2.07 mmol) and N,N -diisopropylethylamine (0.45 mL, 2.59 mmol) was added hydroxylamine hydrochloride (180 mg, 2.59 mmol) and the solution was brought to reflux. After 18 hours, the solution was cooled, the solvent was removed in vacuo and the residue was dissolved in acetic acid (15 mL) and purified by RPHPLC (Method 2, 90/10 (A/B); ramp to 60/40 (A/B) over 200 minutes). The fractions containing pure $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2\cdot\text{HCl}$ as determined by analytical RPHPLC were combined and pH adjusted as described above and lyophilized to give 0.35 g (31%) of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2\cdot\text{HCl}$.

 ^1H NMRFD-MS, m/e 516 (M^+)Analysis for $\text{C}_{25}\text{H}_{33}\text{N}_5\text{O}_5\text{S}\cdot\text{HCl}\cdot\text{H}_2\text{O}$:

25 Calc: C, 52.67; H, 6.36; N, 12.28; Cl, 6.22;
 Found: C, 52.40; H, 6.10; N, 12.25; Cl, 6.51.

H) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2\cdot\text{HCl}$.

30

By a method substantially equivalent to that described in Example 23-D, 0.098 g (50%) of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NH)NH}_2\cdot\text{HCl}$ were prepared from $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{CH}_2\text{-C}_6\text{H}_4\text{C(=NOH)NH}_2\cdot\text{HCl}$.

35

 ^1H NMRFD-MS, m/e 500 (M^+)


Analysis for $C_{25}H_{33}N_5O_4S \cdot 2.6HCl \cdot H_2O$:

Calc: C, 49.03; H, 6.19; N, 11.44;

Found: C, 48.87; H, 5.79; N, 11.15.

5

Example 27



$EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NH)NH_2 \cdot HCl$

10

A) Preparation of $EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NOH)NH_2$.

By a method substantially equivalent to that described in Examples 26-A through 26-E and 24-A, 0.15 g of $EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NOH)NH_2$ were prepared from m-cyanobenzaldehyde.

15

1H NMR

FD-MS, m/e 516 (M^+)

20 Analysis for $C_{25}H_{33}N_5O_5S$:

Calc: C, 58.23; H, 6.45; N, 13.50;

Found: C, 57.99; H, 6.57; N, 13.28.

B) Preparation of $EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NH)NH_2 \cdot HCl$.

25


By a method substantially equivalent to that described in Example 24-B, 0.21 g (20%) of $EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NH)NH_2 \cdot HCl$ were prepared from $EtSO_2-D-Phe-Pro-m-NHCH_2CH_2-C_6H_4C(=NOH)NH_2$.

30

¹H NMRFD-MS, m/e 500 (M⁺)Analysis for C₂₅H₃₃N₅O₄S·2.1HCl·0.7H₂O

Calc: C, 51.00; H, 6.25; N, 11.89;

5 Found: C, 50.79; H, 5.86; N, 11.54.

Example 2810 D-1-Piq-Pro-p-NHCH₂CH₂-C₆H₄C(=NH)NH₂·2HCl


By a method substantially equivalent to that described in Example 23, 0.85 g of 1-Piq-Pro-p-NHCH₂CH₂-C₆H₄C(=NH)NH₂·2HCl were prepared from Cbz-D-1-Piq-Pro-OH and

15 p-NH₂CH₂CH₂-C₆H₄CN·HCl.

¹H NMRFD-MS, m/e 426 (M⁺)Analysis for C₂₄H₃₅N₅O₂·2HCl·2H₂O:

20 Calc: C, 53.93; H, 7.73; N, 13.10;

Found: C, 53.94; H, 7.60; N, 13.06.

Example 2925 D-1-Piq-Pro-m-NHCH₂CH₂-C₆H₄C(=NH)NH₂·2HCl

By a method substantially equivalent to that described in Example 23, 0.8 g of 1-Piq-Pro-m-NHCH₂CH₂-C₆H₄C(=NH)NH₂·2HCl were prepared from Cbz-D-1-Piq-Pro-OH and
 5 m-NH₂CH₂CH₂-C₆H₄CN·HCl.


¹H NMR

FD-MS, m/e 426 (M⁺)

Analysis for C₂₄H₃₅N₅O₂·2HCl·2H₂O:

10 Calc: C, 53.93; H, 7.73; N, 13.10;
 Found: C, 53.62; H, 7.57; N, 13.18.

Example 30



EtSO₂-D-Phe-Pro-p-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂ · HCl

A) Preparation of p-HOCH₂CH₂CH₂-C₆H₄CN.

20

To a stirring solution of methyl p-cyano-dihydrocinnamate (10 g, 53 mmol) in tetrahydrofuran (150 mL) was added LiBH₄ (1.15 g, 53 mmol) and the solution was heated to reflux. After 2 hours, the solution was cooled, and
 25 sodium phosphate buffer (pH 7) was added dropwise. After gas evolution was complete, ethyl acetate and water were added and the layers were separated. The aqueous phase was extracted once with ethyl acetate and the combined ethyl acetate phases were washed with brine, then dried with MgSO₄,
 30 filtered and concentrated to give 8.1 g (95%) of a thick colorless oil.

IR

 ^1H NMRFD-MS, m/e 161 (M^+)

5

B) Preparation of p-BrCH₂CH₂CH₂-C₆H₄CN.

To a stirring solution of p-HOCH₂CH₂CH₂-C₆H₄CN (8.1 g, 50 mmol) in tetrahydrofuran (100 mL) was added
10 triphenylphosphine (14.4 g, 55 mmol) followed by carbon tetrabromide (18.2 g, 55 mmol). After stirring for 18 hours, the solvent was removed in vacuo and the residue was chromatographed over silica gel eluting with a step gradient of hexanes through 20% ethyl acetate/hexanes. The product
15 containing fractions as judged by TLC were combined and concentrated to give 7.3 g (65%) of a thick colorless oil.

IR

 ^1H NMR20 FD-MS, m/e 223 (M^+)Analysis for C₁₀H₁₀BrN:

Calc: C, 53.60; H, 4.50; N, 6.25;

Found: C, 53.90; H, 4.67; N, 6.24.

25 C) Preparation of p-Boc₂NCH₂CH₂CH₂-C₆H₄CN.

To a stirring suspension of NaH (1.4 g of 60% oil dispersion, 34 mmol) in DMF (100 mL) was added dropwise a solution of di-t-butyl iminodicarboxylate (7.4 g, 34 mmol) in
30 DMF (20 mL) via an addition funnel. After gas evolution was complete, a solution of p-BrCH₂CH₂CH₂-C₆H₄CN (7 g, 31 mmol) in DMF was added via an addition funnel and the solution was heated to 70 °C. After stirring for 12 hours, the solution was cooled and the solvent was removed in vacuo. The residue
35 was dissolved in diethyl ether and washed three times with water. The organic phase was dried with MgSO₄, filtered and concentrated and the residue was chromatographed over silica

gel, eluting with hexanes through 20% ethyl acetate/hexanes. The product containing fractions were combined and concentrated to give 9.38 g (84%) of white solid.

5 IR

¹H NMR

FD-MS, m/e 361 (M⁺)

Analysis for C₂₀H₂₈N₂O₄:

Calc: C, 66.64; H, 7.83; N, 7.77;

10 Found: C, 66.40; H, 7.81; N, 7.57.

D) Preparation of p-NH₂CH₂CH₂CH₂-C₆H₄CN·HCl.

By methods substantially equivalent to those described
15 in Example 23-A, 4.3 g (84%) of p-NH₂CH₂CH₂CH₂-C₆H₄CN·HCl were prepared.

IR

¹H NMR

20 FD-MS, m/e 160 (M⁺)

E) Preparation of EtSO₂-D-Phe-Pro-p-NHCH₂CH₂CH₂-
C₆H₄C(=NOH)NH₂·HCl.

25 By methods substantially equivalent to those described
in Examples 1-G and 26-G, 0.32 g of EtSO₂-D-Phe-Pro-p-
NHCH₂CH₂CH₂-C₆H₄C(=NOH)NH₂·HCl were prepared from EtSO₂-D-Phe-
Pro-OH and p-NHCH₂CH₂CH₂-C₆H₄CN·HCl.

30 ¹H NMR

FD-MS, m/e 530 (M⁺)

Analysis for C₂₆H₃₅N₅O₅S·1.2HCl·H₂O:

Calc: C, 52.88; H, 6.51; N, 11.84;

Found: C, 52.71; H, 6.26; N, 11.76.

35

F) Preparation of EtSO₂-D-Phe-Pro-p-NHCH₂CH₂CH₂-
C₆H₄C(=NH)NH₂·HCl.

By a method substantially equivalent to that described in Example 23-D, 0.13 g (67%) of EtSO₂-D-Phe-Pro-p-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl were prepared from EtSO₂-D-Phe-Pro-p-NHCH₂CH₂CH₂-C₆H₄C(=NOH)NH₂·HCl.


¹H NMR

FD-MS, m/e 514 (M⁺)

Analysis for C₂₆H₃₅N₅O₄S·1.5HCl·2H₂O:

10 Calc: C, 51.67; H, 6.75; N, 11.59;
 Found: C, 51.36; H, 6.46; N, 11.28.

Example 31



15

EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl

A) Preparation of EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NOH)NH₂·HCl.

20

By methods substantially equivalent to those described in Examples 1-G and 26-G, 0.32 g of EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NOH)NH₂·HCl were prepared from m-cyanodihydrocinnamic acid.

25

¹H NMR

FD-MS, m/e 530 (M⁺)

Analysis for C₂₆H₃₅N₅O₅S·HCl·1.1H₂O:

30 Calc: C, 53.30; H, 6.57; N, 11.95; Cl, 6.05;
 Found: C, 52.97; H, 6.19; N, 11.96; Cl, 6.13.

B) Preparation of EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl.

5 By a method substantially equivalent to that described in Example 23-D, 0.12 g (62%) of EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl were prepared from EtSO₂-D-Phe-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NOH)NH₂·HCl.

10 ¹H NMR

FD-MS, m/e 514 (M⁺)


Analysis for C₂₆H₃₅N₅O₄S·1.5HCl·H₂O:

Calc: C, 53.26; H, 6.62; N, 11.94;

Found: C, 53.19; H, 6.25; N, 12.00.

15

Example 32



20

1-Piq-Pro-p-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·2HCl

By a method substantially equivalent to that described in Example 23, 0.66 g (48%) of 1-Piq-Pro-p-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl were prepared from 1-Piq-Pro-OH and

25 p-NH₂CH₂CH₂CH₂-C₆H₄CN·HCl.

¹H NMR


FD-MS, m/e 440 (M⁺)

Analysis for C₂₅H₃₇N₅O₂·2.1HCl·H₂O:


30

Calc: C, 56.21; H, 7.75; N, 13.11;

Found: C, 56.36; H, 7.44; N, 12.79.

Example 33

5



By a method substantially equivalent to that described in Example 23, 0.64 (46%) of 1-Piq-Pro-m-NHCH₂CH₂CH₂-C₆H₄C(=NH)NH₂·HCl were prepared from 1-Piq-Pro-OH and

10 m-NH₂CH₂CH₂CH₂-C₆H₄CN·HCl.


¹H NMRFD-MS, m/e 440 (M⁺)15 Analysis for C₂₅H₃₇N₅O₂·2HCl·H₂O:

Calc: C, 56.60; H, 7.79; N, 13.20;

Found: C, 56.92; H, 7.55; N, 13.26.

Example 34

20

25 A) Preparation of Boc-p-NHCH₂C₆H₄NO₂.

To a stirring solution of 4-nitrobenzylamine hydrochloride (15 g, 79 mmol) and N,N-diisopropylethylamine (14 mL, 79 mmol) in dichloromethane (200 mL) was added di-*t*-butyl dicarbonate (17 g, 79 mmol). After 48 h, the solvent
5 was removed in *vacuo* and the residue was dissolved in ethyl acetate (500 mL) and washed twice with 1 M citric acid, once with water, and once with saturated aqueous NaHCO₃. The organic phase was dried with MgSO₄, filtered and concentrated in *vacuo* to give an off white solid which was recrystallized
10 from chloroform/hexanes. Three crops were combined, washed with hexanes and dried in *vacuo* to give 11.5 g (58%) of a white solid.

IR

15 ¹H NMR

FD-MS, m/e 252 (M⁺)

Analysis for C₁₂H₁₆N₂O₄:

Calc: C, 57.13; H, 6.39; N, 11.10;

Found: C, 57.27; H, 6.60; N, 11.13.

20

B) Preparation of p-BocNHCH₂C₆H₄NH₂.

To a stirring solution of p-BocNHCH₂C₆H₄NO₂ (7.5 g, 29.7 mmol) and NiCl₂·6H₂O (17.7 g, 74.3 mmol) in methanol (150 mL)
25 at 0 °C was added NaBH₄ (5.6 g, 149 mmol) in small portions over 30 min. After complete addition of NaBH₄ and 15 min, the solvent was evaporated in *vacuo* and the residue was dissolved in conc. ammonium hydroxide and extracted twice with dichloromethane. The combined organic extracts were
30 washed with brine, dried with MgSO₄, filtered and concentrated in *vacuo* to give 6.4 g (97%) of a white solid.

IR

¹H NMR

35 FD-MS, m/e 222 (M⁺)

Analysis for $C_{12}H_{18}N_2O_2$:

Calc: C, 64.84; H, 8.16; N, 12.60;

Found: C, 65.10; H, 8.42; N, 12.76.

5 C) Preparation of N,N'-di-Cbz-S-methylisothiurea.

To a stirring suspension of bis-S-methylisothiurea sulfate (20 g, 144 mmol) in dichloromethane (200 mL), was added 5 N sodium hydroxide (16 mL). The solution was
10 cooled to 0 °C and benzyl chloroformate (41 mL, 288 mmol) was added dropwise. At the same time, 2 N sodium hydroxide was added at a rate which kept the pH of the solution at approximately 11. The cold bath was then removed and after warming to room temperature, the phases were separated and
15 the aqueous phase was extracted with dichloromethane (250 mL). The combined organic phases were then washed twice with 0.1 N HCl (250 mL) and once with brine (250 mL). The organic phase was then dried with $MgSO_4$, filtered and concentrated in vacuo to give 41 g (79%) of a thick,
20 colorless syrup.

1H NMR

25 D) Preparation of p-BocNHCH₂C₆H₄NHC(NCbz)NHCbz.

To a stirring solution of p-BocNHCH₂C₆H₄NH₂ (5 g, 22.5 mmol) in tetrahydrofuran (50 mL) was added N,N'-di-Cbz-S-methylisothiurea (8.9 g, 24.7 mmol). After 48 h, the solvent was removed in vacuo and the residue was dissolved in
30 chloroform. Silica gel was added and the solvent was removed in vacuo to give an off white powder which was then dry loaded onto a silica gel column. The column was then eluted with a step gradient of 5% ethyl acetate/hexanes through 30% ethyl acetate/hexanes. The product containing fractions
35 (determined by TLC) were combined and concentrated in vacuo to give 7.6 g (63%) of a white solid.

104

IR

 ^1H NMRFD-MS, m/e 532 (M^+)Analysis for $\text{C}_{29}\text{H}_{32}\text{N}_4\text{O}_6$:

5 Calc: C, 65.40; H, 6.06; N, 10.52;
 Found: C, 65.66; H, 6.35; N, 10.59.

E) Preparation of $\text{HCl} \cdot \text{p-NH}_2\text{CH}_2\text{C}_6\text{H}_4\text{NHC}(\text{NCbz})\text{NHCbz}$

10 By a method substantially equivalent to that described
 in example 23-A, 4.7 g (89%) of $\text{HCl} \cdot \text{p-NH}_2\text{CH}_2\text{C}_6\text{H}_4\text{NHC}-$
 $(\text{NCbz})\text{NHCbz}$, a white solid, was prepared from
 $\text{p-BocNHCH}_2\text{C}_6\text{H}_4\text{NHC}(\text{NCbz})\text{NHCbz}$.

15 IR

 ^1H NMRFD-MS, m/e 433 (MH^+)F) Preparation of $\text{EtSO}_2\text{-D-Phe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{NHC}(\text{NH})\text{NH}_2 \cdot \text{HCl}$.

20

 By methods substantially equivalent to those described
 in example 1-G and example 18-F, 1.1 g of $\text{EtSO}_2\text{-D-Phe-Pro-}$
 $\text{p-NHCH}_2\text{C}_6\text{H}_4\text{NHC}(\text{NH})\text{NH}_2 \cdot \text{HCl}$ was prepared from $\text{EtSO}_2\text{-D-Phe-ProOH}$
 and $\text{HCl} \cdot \text{p-NH}_2\text{CH}_2\text{C}_6\text{H}_4\text{NHC}(\text{NCbz})\text{NHCbz}$.


25

IR

 ^1H NMRFD-MS, m/e 501 (M^+)Analysis for $\text{C}_{24}\text{H}_{32}\text{N}_6\text{O}_4\text{S} \cdot \text{HCl} \cdot \text{H}_2\text{O}$:

30 Calc: C, 51.73; H, 6.36; N, 15.14;
 Found: C, 52.32; H, 5.99; N, 14.79.

105

Example 35

5 EtSO₂-D-Phe-Pro-p-NHCH₂CH₂C₆H₄NHC(NH)NH₂ · HCl

By methods substantially equivalent to those described
in example 34, 1.8 g of EtSO₂-D-Phe-Pro-
p-NHCH₂CH₂C₆H₄CNH(NH)NH₂ · HCl was prepared from
10 4-nitrophenethylamine · HCl.


IR

¹H NMR

FD-MS, m/e 515 (MH⁺)

15 HRMS (FAB), m/e calcd. for C₂₅H₃₅N₆O₄S: 515.2441

Found: 515.2483

Example 36

20

EtSO₂-D-Phe-Pro-m-NHCH₂C₆H₄NHC(NH)NH₂ · HCl

By methods substantially equivalent to those described
25 in examples 30-C and 34-B through 34-F, 1.8 g of EtSO₂-D-Phe-

Pro-m-NHCH₂C₆H₄NHC(NH)NH₂·HCl was prepared from 3-nitrobenzyl bromide.

IR

5 ¹H NMR


FD-MS, m/e 501 (MH⁺)

HRMS (FAB), m/e calcd. for C₂₄H₃₃N₆O₄S: 501.2284

Found: 501.2280

10

Example 37



EtSO₂-D-Phe-Pro-m-NHCH₂CH₂C₆H₄NHC(NH)NH₂·HCl

15

By methods substantially equivalent to those described in examples 26-D, 26-B (using 5% Pd/C in place of Pd/BaSO₄ and ethyl acetate in place of toluene), and 34-D through 34-F, 0.85 g of EtSO₂-D-Phe-Pro-m-NHCH₂CH₂C₆H₄NHC(NH)NH₂·HCl was prepared from 3-nitrocinnamic acid.

20

¹H NMR

FD-MS, m/e 515 (MH⁺)


Analysis for C₂₅H₃₄N₆O₄S·2.2HCl·0.5H₂O:

25

Calc: C, 49.73; H, 6.21; N, 13.92;

Found: C, 49.45; H, 5.82; N, 13.55.

107

Example 38

5 D-1-Piq-Pro-p-NHCH₂C₆H₄NHC(NH)NH₂ · 2HCl

By methods substantially equivalent to those described in example 34, 0.94 g of D-1-Piq-Pro-p-NHCH₂C₆H₄NHC(NH)NH₂ · 2HCl was prepared. The final product
 10 was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 70/30 (A/B), 180 min.

¹H NMR

FD-MS, m/e 427 (MH⁺)


15 Analysis for C₂₃H₃₄N₆O₂ · 2HCl:

Calc: C, 55.31; H, 7.26; N, 16.82; Cl, 14.20;

Found: C, 55.05; H, 7.23; N, 16.55; Cl, 14.24.

Example 39

20



D-1-Piq-Pro-p-NHCH₂CH₂C₆H₄NHC(NH)NH₂ · 2HCl

25 By methods substantially equivalent to those described in example 35, 1.03 g of D-1-Piq-Pro-p-NHCH₂CH₂C₆H₄NHC(NH)NH₂ · 2HCl was prepared. The final

product was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 70/30 (A/B), 180 min.

5 ^1H NMR

FD-MS, m/e 441 (M^+)


Analysis for $\text{C}_{24}\text{H}_{36}\text{N}_6\text{O}_2 \cdot 2\text{HCl} \cdot 1.5\text{H}_2\text{O}$:

Calc: C, 53.33; H, 7.65; N, 15.55; Cl, 13.12;

Found: C, 53.41; H, 7.45; N, 15.37; Cl, 13.48.

10

Example 40



15

D-1-Piq-Pro-m-NHCH₂C₆H₄NHC(NH)NH₂ · 2HCl

By methods substantially equivalent to those described in example 36, 1.04 g of D-1-Piq-Pro-m-NHCH₂C₆H₄NHC(NH)NH₂ · 2HCl was prepared. The final product was purified by RPHPLC (method 2, 98/2 (A/B), ramp to 70/30 (A/B), 180 min.


^1H NMR

FD-MS, m/e 427 (M^+)

25 Analysis for $\text{C}_{23}\text{H}_{34}\text{N}_6\text{O}_2 \cdot 2\text{HCl} \cdot \text{H}_2\text{O}$:

Calc: C, 53.38; H, 7.40; N, 16.24; Cl, 13.70;

Found: C, 53.25; H, 7.50; N, 16.23; Cl, 13.88.

Example 415 D-1-Piq-Pro-m-NHCH₂CH₂C₆H₄NHC(NH)NH₂ · 2HCl

By methods substantially equivalent to those described
in example 37, 0.96 g D-1-Piq-Pro-m-
NHCH₂CH₂C₆H₄NHC(NH)NH₂ · 2HCl of was prepared. The final
10 product was purified by RPHPLC (method 2, 98/2 (A/B), ramp
to 70/30 (A/B), 180 min.


¹H NMRFD-MS, m/e 441 (M⁺)15 Analysis for C₂₄H₃₆N₆O₂ · 2.1HCl · 1.5H₂O:

Calc: C, 52.97; H, 7.61; N, 15.44; Cl, 13.68;

Found: C, 52.80; H, 7.57; N, 15.46; Cl, 13.35.

Example 42

20

D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂ · 2HCl

25 A) Preparation of Cbz-DL-1-Piq-cis-Ohi-OEt.

By a method substantially equivalent to that described in Example 1-A, 16.6 g (100%) of Cbz-DL-1-Piq-cis-Ohi-OEt was prepared from Cbz-DL-1-Piq-OH and cis-Ohi-OEt·HCl.

¹H NMR

5 FD-MS, m/e 496 (M⁺)

Analysis for C₂₉H₄₀N₂O₅:

Calc: C, 70.13; H, 8.12; N, 5.64;

Found: C, 69.96; H, 8.23; N, 5.73.

10 B) Preparation of Cbz-D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz.

By methods substantially equivalent to those described in Example 1-D and 18-E, Cbz-D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz and Cbz-L-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz were prepared from Cbz-D,L-1-Piq-Pro-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl. These diastereomers were separated by silica gel chromatography using an ethyl acetate/hexanes gradient. The fractions containing the leading diastereomer (R_f = 0.31, ethyl acetate) were pooled and concentrated to give 1.3 g of Cbz-L-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz. The fractions containing the trailing diastereomer (R_f = 0.19, ethyl acetate) were pooled and concentrated to give 1.5 g of Cbz-D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz as a white foam.

¹H NMR

FD-MS, m/e 735 (MH⁺)

Analysis for C₄₃H₅₁N₅O₆:

30 Calc: C, 70.37; H, 7.00; N, 9.54;

Found: C, 70.20; H, 7.22; N, 9.36.

C) Preparation of D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·2HCl.

35

By a method substantially equivalent to that described in Example 18-F, 0.61 g (63%) of D-1-Piq-cis-Ohi-p-

111


NHCH₂C₆H₄C(NH)NH₂·2HCl were prepared from Cbz-D-1-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NHCbz. The HPLC gradient used in this case was a ramp of 98/2 A/B to 70/30 A/B over 120 minutes.

5 ¹H NMRFAB-MS, m/e 466.4 (MH⁺)Analysis for C₂₇H₃₉N₅O₂·2HCl·1H₂O:

Calc: C, 58.27; H, 7.79; N, 12.58;

Found: C, 58.66; H, 7.56; N, 12.78.

10

Example 43

15

D-3-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·2HClA) Preparation of D-3-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·2HCl


By methods substantially equivalent to those described in Example 1-A, 1-D, 18-E, and 18-F, 1.3 g of 3-Piq-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·2HCl was prepared. The HPLC gradient used in this case was a ramp of 98/2 A/B to 70/30 A/B over 120 minutes.

25 ¹H NMRFAB-MS, m/e 466.4 (MH⁺)Analysis for C₂₇H₃₉N₅O₂·2HCl:

Calc: C, 60.22; H, 7.67; N, 13.00;

Found: C, 59.95; H, 7.73; N, 12.89.

30

Example 44

5 EtSO₂-Phg-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 ((S-cis)-N-[[4-(aminoiminomethyl)phenyl]methyl]-1-
 [N-(ethylsulfonyl)-D-phenylglycyl]-1H-indole-2-carboxamide
 hydrochloride)

10 A) Preparation of Boc-D-Phg-cis-Ohi-OEt

By a method substantially equivalent to that described
 in Example 1-A, 14.9 g (58%) of Boc-D-Phg-cis-Ohi-OEt was
 prepared from Boc-D-Phg-OH and (S)-cis-octahydroindole-2-
 15 carboxylic acid ethyl ester·HCl.

¹H NMR

FD-MS, m/e 430 (M⁺)

Analysis for C₂₄H₃₄N₂O₅:

20 Calc: C, 66.95; H, 7.96; N, 6.51;
 Found: C, 66.69; H, 8.02; N, 6.40.

B) Preparation of D-Phg-cis-Ohi-OEt·HCl.

25 To a cold (0°C), stirring solution of Boc-D-Phg-cis-Ohi-
 OEt in ethyl acetate was bubbled HCl gas for 10 min. After
 stirring for 2 h while warming to room temperature, the
 solvent was removed in vacuo. The resulting solid was
 suspended in diethyl ether and subsequently isolated by
 30 filtration to give 10.7 g (97%) of D-Phg-cis-Ohi-OEt·HCl.

¹H NMR

FD-MS, m/e 331 (M⁺)

Analysis for C₁₉H₂₇N₂O₃Cl:

Calc: C, 62.20; H, 7.41; N, 7.64;
5 Found: C, 62.42; H, 7.36; N, 7.85.

C) Preparation of EtSO₂-D-Phg-cis-Ohi-OEt.

To a solution of D-Phg-cis-Ohi-OEt·HCl (10 g, 27 mmol)
10 and N,N-diisopropylethylamine (10.7 mL, 61 mmol) in THF (200 mL) at -78 °C, was added dropwise via an addition funnel a solution of ethanesulfonyl chloride (3.9 g, 30 mmol) in THF (20 mL). The cold bath was then left unattended and the solution warmed slowly to room temperature. After about 18
15 h, the solution was concentrated in vacuo. The residue was dissolved in ethyl acetate (200 mL), washed twice each with 1 N citric acid (200 mL), saturated aqueous NaHCO₃ (200 mL) and brine (200 mL). The organic phase was then dried with MgSO₄, filtered, and concentrated in vacuo to give 11.2 g (97%) of a
20 yellow foam.

¹H NMR

FD-MS, m/e 422 (M⁺)

Analysis for C₂₁H₃₀N₂O₅S:

25 Calc: C, 59.69; H, 7.16; N, 6.63;
Found: C, 59.94; H, 7.08; N, 6.78.


D) Preparation of EtSO₂-Phg-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl
30

By methods substantially equivalent to those described in Examples 1-D, 18-E and 18-F, 0.62 g of EtSO₂-Phg-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·2HCl were obtained. The HPLC gradient used in this case was a ramp of 90/10 A/B to 60/40 A/B over
35 120 minutes.

¹H NMRFAB-MS, m/e 526.3 (MH⁺)Analysis for C₂₇H₃₅N₅O₄S·HCl:

Calc: C, 57.69; H, 6.45; N, 12.46;

5 Found: C, 57.47; H, 6.48; N, 12.20.

Example 45

10

EtSO₂-Phe-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 ((S-cis)-N-[[4-(aminoiminomethyl)phenyl]methyl]-1-[N-(ethylsulfonyl)-D-phenylalanyl]-1H-indole-2-carboxamide
 hydrochloride)

15


A) Preparation of EtSO₂-Phe-cis-Ohi-p-
 NHCH₂C₆H₄C(NH)NH₂·2HCl

By methods substantially equivalent to those described
 20 in Example 44, 1.5 g EtSO₂-Phe-cis-Ohi-p-
 NHCH₂C₆H₄C(NH)NH₂·HCl were prepared from Boc-D-Phe-OH.

¹H NMRFAB-MS, m/e 540.3 (MH⁺)25 Analysis for C₂₈H₃₇N₅O₄S·HCl·H₂O:

Calc: C, 56.51; H, 6.94; N, 11.77;

Found: C, 56.24; H, 6.55; N, 11.72.

Example 46

5 HOOCCH₂-D-Phe-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 (N-(carboxymethyl)-D-phenylalanyl-N-
 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
 hydrochloride)

10 A) Preparation N-(t-BuOOCCH₂)-N-Boc-D-Phe-Pro-OBn.

To a solution of D-Phe-Pro-OBn·HCl (20 g, 51 mmol) in DMF (100 mL) was added t-butyl bromoacetate (9.9 g, 56 mmol) in one portion and N,N-diisopropylethylamine (17.4 mL, 101
 15 mmol) dropwise over 30 min. This mixture was allowed to stir for 18 hrs at room temperature. Di-t-butyl dicarbonate (16.6 g, 76 mmol) and N,N-diisopropylethylamine (13.2 mL, 76 mmol) were then added in one portion and the reaction was allowed to stir an additional 24 h. The solvent was removed in vacuo
 20 and the residue was partitioned between ethyl acetate (1 L) and 1 M aqueous citric acid (500 mL). The layers were separated and the organic phase was washed once with 1 M aqueous citric acid, twice with saturated aqueous sodium bicarbonate, and once with brine (500 mL each). The organic
 25 phase was dried (Na₂SO₄), filtered, and concentrated in vacuo. The amber oil was purified by silica gel chromatography eluting with a ethyl acetate/hexanes gradient (hexanes to 30% ethyl acetate/hexanes). Fractions containing product were combined and concentrated to give 19.0 g (66%)
 30 as a colorless oil which slowly crystallized upon standing.

¹H NMRFD-MS, m/e 566 (M⁺)Analysis for C₃₂H₄₂N₂O₇:

- 5 Calc: C, 67.82; H, 7.47; N, 4.94;
 Found: C, 68.06; H, 7.33; N, 5.17.

B) Preparation of N-(t-BuOOCCH₂)-N-Boc-D-Phe-Pro-OH.

- 10 To a solution of N-(t-BuOOCCH₂)-N-Boc-D-Phe-Pro-OBn
 (18.5 g, 33 mmol) in ethyl acetate (250 mL) was added 5% Pd/C
 catalyst (5 g). This solution was degassed *in vacuo* several
 times and placed under an atmosphere of hydrogen for 2 h with
 stirring. The balloon was removed, diatomaceous earth was
15 added and the slurry was filtered over a pad of diatomaceous
 earth. The filtrate was concentrated *in vacuo* to give 13.2 g
 (84%) of a white foam.

¹H NMR

- 20 FD-MS, m/e 476 (M⁺)

Analysis for C₂₅H₃₆N₂O₇:

- Calc: C, 63.01; H, 7.61; N, 5.88;
 Found: C, 63.23; H, 7.73; N, 5.59.

- 25 C) Preparation of N-(t-BuOOCCH₂)-N-Boc-D-Phe-Pro-p-
 NHCH₂C₆H₄C(NH)NHCbz.

- By a method substantially equivalent to Example 18-E,
 2.7 g (90%) of N-(t-BuOOCCH₂)-N-Boc-D-Phe-Pro-p-
30 NHCH₂C₆H₄C(NH)NHCbz was prepared from N-(t-BuOOCCH₂)-N-Boc-
 D-Phe-Pro-OH and p-H₂NCH₂C₆H₄C(NH)NHCbz·2HCl.

¹H NMRFD-MS, m/e 743 (MH⁺)

- 35 Analysis for C₄₁H₅₁N₅O₈:

- Calc: C, 66.38; H, 6.93; N, 9.44;
 Found: C, 66.08; H, 6.92; N, 9.16.

D) Preparation of $\text{HOOCCH}_2\text{-D-Phe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$.

To a cooled (0°C) solution of $\text{N-(t-BuOOCCH}_2\text{)-N-Boc-D-Phe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NHCBz}$ (2.2 g, 3 mmol) in dioxane (100 mL) was bubbled HCl gas for 10 minutes. After stirring for 3 h while warming to room temperature, the dioxane was removed in vacuo. The residue was dissolved in a mixture of absolute ethanol (150 mL), water (75 mL) and 1 N HCl (6 mL). To this was added 5% Pd/C (1 g). After degassing using vacuum, this mixture was placed under a hydrogen atmosphere for 16 h while stirring at room temperature. Diatomaceous earth was added and the resulting slurry was filtered over a pad of diatomaceous earth. The filtrate was concentrated in vacuo to a residue and this was immediately purified by HPLC Method 2 (98/2 A/B to 70/30 A/B ramp over 2 hrs). Fractions containing pure product were pooled and lyophilized to give 1.1 g (72%) of a white solid.

$^1\text{H NMR}$


FAB-MS, m/e 452.3 (MH^+)

Analysis for $\text{C}_{24}\text{H}_{29}\text{N}_5\text{O}_4\cdot 2\text{HCl}$:

Calc: C, 54.97; H, 5.96; N, 13.35;

Found: C, 55.21; H, 6.11; N, 13.39.

Example 47



$\text{HOOCCH}_2\text{-D-Phe-cis-Ohi-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

A) Preparation t-BuOOCCH₂-D-Phe-cis-Ohi-OEt.

To a solution of D-Phe-cis-Ohi-OEt·HCl (30 g, 79 mmol)
5 in acetonitrile (400 mL) was added N,N-diisopropylethylamine
(41 mL, 236 mmol) and t-butyl bromoacetate (14 mL, 87 mmol).
This solution was warmed to 50 °C and maintained there for
3 h. After cooling to room temperature, the solution was
concentrated *in vacuo*. The residue was dissolved in ethyl
10 acetate (300 mL) and this solution was washed twice with
saturated aqueous ammonium chloride (200 mL), twice with
saturated aqueous sodium bicarbonate (200 mL), and twice with
brine (200 mL). The organic layer was dried (MgSO₄),
filtered, and concentrated *in vacuo* to give an orange oil
15 which was purified by silica gel chromatography eluting with
a gradient of hexanes to 1:1 hexanes/ethyl acetate.
Fractions containing product (as judged by TLC) were combined
and concentrated to give 33.2 g (92%) of a colorless oil.

20 ¹H NMRFD-MS, m/e 458 (M⁺)Analysis for C₂₆H₃₈N₂O₅:

Calc: C, 68.10; H, 8.35; N, 6.11;

Found: C, 68.37; H, 8.47; N, 5.90.

25

B) Preparation of N-(t-BuOOCCH₂)-N-Boc-D-Phe-cis-Ohi-OH.

To a solution of t-BuOOCCH₂-D-Phe-cis-Ohi-OEt (30 g, 65
mmol) in THF (200 mL) was added N,N-diisopropylethylamine (17
30 mL, 98 mmol) and di-t-butyl dicarbonate (15.7 g, 72 mmol).
This solution was brought to gentle reflux and maintained for
16 hrs. Heating was discontinued, and once cool, the
solution was concentrated *in vacuo*. The residue was
dissolved in ethyl acetate (400 mL) and washed twice with 1.0
35 M citric acid (200 mL), twice with saturated aqueous sodium
bicarbonate (200 mL), and twice with brine (200 mL). The
organic solution was dried (MgSO₄), filtered, and

concentrated in *vacuo* to give a yellow oil. A portion of this oil (24.8 g, 44 mmol) was dissolved in 300 mL of dioxane. To this was added a solution consisting of 2.05 g LiOH·H₂O (49 mmol) in 150 mL water. This mixture was allowed
5 to stir for 5 hrs at room temp at which time 100 mL of saturated aqueous ammonium chloride was added. Solvents were removed in *vacuo* and the residue was partitioned between saturated aqueous sodium bicarbonate and diethyl ether. The layers were separated and the aqueous layer was acidified to
10 pH 3 with citric acid. The acidic aqueous solution was extracted 3 times with diethyl ether (200 mL) and these were combined, dried (MgSO₄), filtered and concentrated to give 24.3 g of N-(t-BuOOCCH₂)-N-Boc-D-Phe-cis-Ohi-OH as a white foam.

15

¹H NMRFD-MS, m/e 530 (M⁺)Analysis for C₂₉H₄₂N₂O₇:

Calc: C, 65.64; H, 7.98; N, 5.28;

20

Found: C, 65.39; H, 8.04; N, 5.39.

C) Preparation of HOOCCH₂-D-Phe-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl

25


By methods substantially equivalent to those described in Example 18-E and 46-D, 1.2 g (67%) of HOOCCH₂-D-Phe-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl were obtained.

¹H NMR30 FAB-MS, m/e 506.3 (MH⁺)Analysis for C₂₈H₃₅N₅O₄·2HCl·H₂O:

Calc: C, 57.24; H, 6.52; N, 11.92;

Found: C, 57.40; H, 6.30; N, 11.69.

120

Example 48

5 HOOCCH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 (N-(carboxymethyl)-D-cyclohexylalanyl-N-[[4-(
 (aminoiminomethyl)phenyl]methyl]-L-prolinamide)

A) Preparation of HOOCCH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl

10

By methods substantially equivalent to those described
 in Example 46, 0.92 g of HOOCCH₂-D-Cha-Pro-p-
 NHCH₂C₆H₄C(NH)NH₂·HCl were prepared.

15 ¹H NMR


FD-MS, m/e 458 (M⁺)

Analysis for C₂₄H₃₅N₅O₄·HCl·0.5H₂O:

Calc: C, 57.30; H, 7.41; N, 13.92;

Found: C, 57.52; H, 7.29; N, 13.83.

20

Example 49

HOOCCH₂-D-Cha-cis-Ohi-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 ((S-cis)-N-[[4-(aminoiminomethyl)phenyl]methyl]-
 1-[N-(carboxymethyl)-D-cyclohexylalanyl]-1H-indole-
 2-carboxamide hydrochloride)

- 5 A) Preparation of HOOCCH₂-D-Phe-cis-Ohi-p-
 NHCH₂C₆H₄C(NH)NH₂·HCl

By methods substantially equivalent to those described
 in Example 1-A and 1-D, 18-E, 44-B, 47-A(using benzyl
 10 bromoacetate) and 18-F, 0.75 g of HOOCCH₂-D-Cha-cis-Ohi-p-
 NHCH₂C₆H₄C(NH)NH₂·HCl were obtained starting from Boc-
 D-Cha and cis-Ohi-OEt·HCl. HPLC Method 2 was used in the
 purification of this material using a gradient of 98/2 A/B to
 70/30 A/B over 3 h.

15


¹H NMR

FAB-MS, m/e 512.3 (MH⁺)

Analysis for C₂₈H₄₁N₅O₄·HCl:

20 Calc: C, 61.36; H, 7.72; N, 12.78;
 Found: C, 61.08; H, 7.47; N, 12.53.

Example 50



25

HOOCCH₂-D-Phe-Pro(4-cis-isoamyl)-p-NHCH₂C₆H₄C(NH)NH₂·HCl

- A) Preparation of Cbz-Pro(4-trans-OH)-OEt.

To a solution of Cbz-Pro(4-trans-OH)-OH (33 g, 124 mmol) in ethanol (500 mL) was added p-toluenesulfonic acid (1 g) and the solution was heated to reflux. After 16 h, the solution was cooled to room temperature, and the solvent was removed in vacuo. The residue was dissolved in ethyl acetate (400 mL) and washed twice with saturated aqueous NaHCO₃, and twice with a saturated aqueous sodium chloride solution. The ethyl acetate solution was dried with MgSO₄, filtered and concentrated in vacuo to give 34.5 g (95%) of a colorless oil.

¹H NMR

FD-MS, m/e 293 (M⁺)

Analysis for C₁₅H₁₉NO₅:

Calc: C, 61.42; H, 6.53; N, 4.77;
Found: C, 61.20; H, 6.65; N, 4.73.

B) Preparation of Cbz-Pro(4-oxo)-OEt.

Cbz-Pro(4-trans-OH)-OEt (32.7 g, 111 mmol) was dissolved in dichloromethane (500 mL) with mechanical stirring in a 1 L round bottom flask. To this solution was added 3 Å molecular sieves (100 g) and pyridinium chlorochromate (60 g, 278 mmol), in portions small enough to maintain efficient stirring. After stirring for 12 h at room temperature, diethyl ether (200 mL) was added and the black slurry was decanted from a tarry residue and flushed through a column of silica gel (200 g). The residue was washed twice with dichloromethane (200 mL) and the combined washings were also passed through the silica plug. The filtrate was flushed through a silica gel column with 1:1 ethyl acetate/hexanes (4 L) and 500 mL fractions were collected. All fractions containing product, as judged by TLC, were combined and concentrated in vacuo to give 23.8 g (74%) of a colorless oil.

¹H NMR

FD-MS, m/e 291 (M⁺)

Analysis for C₁₅H₁₇NO₅:

Calc: C, 61.85; H, 5.88; N, 4.81;
5 Found: C, 61.57; H, 5.82; N, 4.71.

C) Preparation of Cbz-Pro(4-isobutylmethyldiene)-OEt.

Potassium t-butoxide (34 g, 288 mmol) was suspended in
10 tetrahydrofuran (800 mL) in an oven dried 2-neck 2 L round
bottom flask equipped with a nitrogen inlet, magnetic stir
bar, and addition funnel. To this suspension was added, in
several portions, isoamyltriphenylphosphonium bromide (120 g,
288 mmol). After stirring for 30 min, a solution of Cbz-
15 Pro(4-oxo)-OEt (70 g, 240 mmol) in tetrahydrofuran (150 mL)
was added dropwise via an addition funnel over a 1 h period.
After stirring for an additional 2 h, saturated aqueous NH₄Cl
(100 mL) was added. This solution was diluted with ethyl
acetate (750 mL) and the layers were separated. The organic
20 layer was washed two times with 1 N citric acid, twice with
saturated aqueous NaHCO₃, and twice with a saturated aqueous
sodium chloride solution. The organic solution was dried
with MgSO₄, filtered and concentrated to give a yellow oil.
This oil was purified by flash chromatography over silica
25 gel, eluting with 2:1 hexanes/ethyl acetate. Fractions
containing product (as judged by TLC) were combined and
concentrated in vacuo to give 37 g (45%) of a colorless oil.

¹H NMR

30 FD-MS, m/e 345 (M⁺)

Analysis for C₂₀H₂₇NO₄:

Calc: C, 69.54; H, 7.88; N, 4.05;
Found: C, 69.74; H, 7.85; N, 3.99.

35 D) Preparation of Pro(4-cis-isoamyl)-OEt·HCl.

To a solution of Cbz-Pro(4-isobutylmethyldene)-OEt (37 g, 107 mmol) in ethanol (500 mL) was added 5% Pd/C (5 g). Nitrogen gas was bubbled through this solution for 5 min and then hydrogen gas was bubbled through for 3 h. The solution
5 was filtered through a pad of diatomaceous earth. Hydrogen chloride gas was then bubbled through the solution until saturation, and then the solution was concentrated in vacuo to give 26 g (97%) of amber oil.

10 ^1H NMR

FD-MS, m/e 214 (M^+)

Analysis for $\text{C}_{12}\text{H}_{24}\text{ClNO}_2$:

Calc: C, 57.70; H, 9.68; N, 5.61;

Found: C, 57.46; H, 9.50; N, 5.82.

15

E) Preparation of $\text{HOOCCH}_2\text{-D-Phe-Pro(4-cis-isoamyl)-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

By methods substantially equivalent to those described
20 in Example 1-A and 1-D, 18-E, 44-B, 47-A(using benzyl bromoacetate) and 18-F, 0.27 g of $\text{HOOCCH}_2\text{-D-Phe-Pro(4-cis-isoamyl)-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$ were obtained starting from Boc-D-Cha and Pro(4-cis-isoamyl)-OEt·HCl. HPLC Method 2 was used in the purification of this material using a gradient of
25 98/2 A/B to 50/50 A/B over 3 h.

^1H NMR


FAB-MS, m/e 528.4 (MH^+)

Analysis for $\text{C}_{29}\text{H}_{45}\text{N}_5\text{O}_4\cdot 1.9\text{HCl}$:

30 Calc: C, 58.34; H, 7.92; N, 11.73; Cl, 11.28;

Found: C, 58.30; H, 7.85; N, 11.83; Cl, 11.27.

125

Example 51

5 D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂ · 2HCl
 (D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-
 prolinamide dihydrochloride

A) Preparation of Boc-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NHCBz

10

By methods substantially equivalent to those described
 in Examples 1-A, 46-E, and 18-E, 32.5 g (94%) of Boc-D-Cha-
 Pro-p-NHCH₂C₆H₄C(NH)NHCBz was prepared starting from Boc-D-
 Cha-Pro-OH.

15

¹H NMR

FD-MS, m/e 634 (M⁺)

Analysis for C₃₅H₄₇N₅O₆:

Calc: C, 66.33; H, 7.47; N, 11.05;

20

Found: C, 66.30; H, 7.47; N, 11.26.

B) Preparation of D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NHCBz · 2HCl

By a method substantially equivalent to that described
 25 in Example 23-A, 9.6 g (101%) of D-Cha-Pro-p-
 NHCH₂C₆H₄C(NH)NHCBz · 2HCl was prepared from Boc-D-Cha-Pro-p-
 NHCH₂C₆H₄C(NH)NHCBz.

¹H NMR

30 FD-MS, m/e 534 (M⁺)

Analysis for $C_{30}H_{41}N_5O_4Cl_2$:

Calc: C, 59.40; H, 6.81; N, 11.54; Cl, 11.69;

Found: C, 59.54; H, 6.80; N, 11.77; Cl, 11.21.

5 C) Preparation of D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·2HCl

By a method substantially equivalent to that described in Example 18-F, 0.74 g (62%) of D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·2HCl was prepared. HPLC Method 2 was used in the purification of this material using a gradient of 98/2 A/B to 75/25 A/B over 2.5 hrs.

¹H NMR


FAB-MS, m/e 400.3 (M⁺)

15 Analysis for $C_{22}H_{33}N_5O_2 \cdot 2.1HCl$:

Calc: C, 55.50; H, 7.43; N, 14.71; Cl, 15.64;

Found: C, 55.64; H, 7.50; N, 14.65; Cl, 15.81.

Example 52



HOOCCH₂CH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl

(N-(2-carboxyethyl)-D-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
hydrochloride)

25

A) Preparation of HOOCCH₂CH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl

D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NHCbz·2HCl (2.5 g, 4.1 mmol) was suspended in EtOAc (100 mL) with stirring. A solution of 1 M KHCO₃ (100 mL) was added to the suspension and the mixture was stirred until all solid had dissolved. The mixture was transferred to a separatory funnel and the layers separated. The organic layer was dried (MgSO₄), filtered and concentrated in vacuo to give 1.24 g of the free base as a white solid. This solid was dissolved in EtOH (100 mL). Benzyl acrylate (0.41 g, 2.6 mmol) was added and the solution was stirred for 2 days at room temperature. Then, to this solution was added water (50 mL), 1 N HCl (4.6 mL) and 5% Pd/C (0.5 g) and the stirring suspension was degassed and placed under an atmosphere of hydrogen. After 16 h, diatomaceous earth was added and the slurry was filtered over a pad of diatomaceous earth. The filtrate was concentrated to a volume of 25 mL in vacuo and purified by preparative RPHPLC method 2, using a gradient of 98/2 A/B to 75/25 A/B over 2.5 h. Fractions containing pure product (as judged by analytical HPLC) were pooled, concentrated and lyophilized to give 0.27 g (23%) of HOOCCH₂CH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl.

¹H NMR


FAB-MS; m/e 472.4 (MH⁺)

Analysis for C₂₅H₃₇N₅O₄·1.9HCl:

Calc: C, 55.50; H, 7.25; N, 12.94; Cl, 12.45;

Found: C, 55.26; H, 7.26; N, 13.21; Cl, 12.85.

Example 53



HO₂CCH₂-D-Cha-Pro-4-NHCH₂-piperidine·HCl

A) Preparation of Boc-4-(aminomethyl)pyridine.

By a method substantially equivalent to that described
5 in example 34-A, 19 g (87%) of Boc-4-(aminomethyl)pyridine
was prepared from 4-(aminomethyl)pyridine.

¹H NMR

10 B) Preparation of 4-BocNHCH₂-N-Cbz-piperidine.

4-BocNHCH₂-pyridine (10 g, 48 mmol) was dissolved in
ethanol (280 mL) and 5% Rh/C (10 g) was added. The
suspension was shaken under an atmosphere of hydrogen (4.1
15 bar, 60 psi) at 60 °C overnight. The catalyst was then
filtered off and the solution was concentrated in vacuo to
give 9.0 g of a gray solid. A 3.2 g portion of the solid was
dissolved in tetrahydrofuran (75 mL) and an aqueous solution
(75 mL) of potassium carbonate (4.2 g, 30 mmol) was added.
20 To this stirring solution was added benzyl chloroformate (2.3
mL, 16 mmol). After 15 min, the solution was concentrated in
vacuo to about 1/2 the original volume and then diluted with
ethyl acetate. The organic phase was separated and washed
with brine, then dried (MgSO₄), filtered and concentrated in
25 vacuo to give 4.6 g (76%) of a white solid.

¹H NMR

FD-MS, m/e 349 (M⁺)

30 C) Preparation of 4-NH₂CH₂-N-Cbz-piperidine·HCl.

By a method substantially equivalent to that described
in example 23-A, 3 g (84%) of 4-NH₂CH₂-N-Cbz-piperidine·HCl
was prepared from 4-BocNHCH₂-N-Cbz-piperidine.

IR

¹H NMRFD-MS, m/e 249 (MH⁺)5 D) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-OH.

N-(t-BuO₂CCH₂)-N-Boc-D-Phe-Pro-OH (13 g, 27 mmol) was dissolved in ethanol (750 mL) and PtO₂ (13 g) was added. The suspension was shaken under an atmosphere of hydrogen at 4.1
10 bar (60 psi) at 40 °C for 16 h. The catalyst was then filtered off and the filtrate was concentrated in vacuo to give 11.7 g (90%) of a white foam.

IR

15 ¹H NMRFD-MS, m/e 483 (M⁺)Analysis for C₂₅H₄₂N₂O₇:

Calc: C, 62.22; H, 8.77; N, 5.80;

Found: C, 62.99; H, 8.96; N, 5.48.

20

E) Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-piperidine·HCl.


By methods substantially equivalent to those described in example 1-G and example 46-D, 1.1 g of HO₂CCH₂-D-Cha-Pro-
25 4-NHCH₂-piperidine·HCl was prepared from N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-OH and HCl·4-NH₂CH₂-N-Cbz-piperidine. The product was purified by RPHPLC method 2, ramping from 98/2 (A/B) through 70/30 (A/B) over 2 h.

30 IR

¹H NMRFD-MS, m/e 423 (M⁺)Analysis for C₂₂H₃₈N₄O₄·2HCl·1.5H₂O:

Calc: C, 50.57; H, 8.29; N, 10.72;

35 Found: C, 50.31; H, 8.46; N, 10.93.

Example 54HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂-piperidine·HCl

5

Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂-piperidine·HCl.

By methods substantially equivalent to those described in example 52, 0.59 g of HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂-
 10 piperidine·HCl was prepared from 4-aminoethylpyridine. The product was purified by RPHPLC method 2, ramping from 98/2 (A/B) through 70/30 (A/B) over 2 h.


IR

15 ¹H NMRFD-MS, m/e 437 (M⁺)Analysis for C₂₃H₄₀N₄O₄·2.5HCl·1.5H₂O:

Calc: C, 49.80; H, 8.27; N, 10.10;

Found: C, 49.95; H, 8.08; N, 10.34.

20

Example 55HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂CH₂-piperidine·HCl

A) Preparation of 4-hydroxypropyl-N-Cbz-piperidine.

By methods substantially equivalent to those described
5 in example 53-B, 28 g (67%) of 4-hydroxypropyl-N-Cbz-
piperidine was prepared from 4-hydroxypropylpyridine.

¹H NMR

10 B) Preparation of 4-(NH₂CH₂CH₂CH₂)-N-Cbz-piperidine·HCl.

By methods substantially equivalent to those described
in examples 30-B, 30-C and 30-D, 7.3 g of 4-(NH₂CH₂CH₂CH₂)-N-
Cbz-piperidine·HCl were prepared from 4-hydroxypropyl-N-Cbz-
15 piperidine.

¹H NMR

FD-MS, m/e 276 (M⁺)

20 C) Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂CH₂-
piperidine·HCl.

By methods substantially equivalent to those described
in example 53-D and 53-E, 0.39 g of HO₂CCH₂-D-Cha-Pro-4-
25 NHCH₂CH₂CH₂-piperidine·HCl was prepared from N-(t-BuO₂CCH₂)-N-
Boc-D-Cha-Pro-OH and 4-aminopropyl-N-Cbz-piperidine·HCl. The
product was purified by RPHPLC method 2, ramping from 98/2
(A/B) through 70/30 (A/B) over 2 h.

30 IR


¹H NMR

IS-MS, m/e 451.4 (MH⁺)

Analysis for C₂₄H₄₂N₄O₄·2HCl·H₂O:

Calc: C, 53.23; H, 8.56; N, 10.35;

35 Found: C, 53.43; H, 8.63; N, 10.19.

Example 56

- 5 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-1-amidinopiperidine}\cdot\text{HCl}$
 (N-(carboxymethyl)-D-cyclohexylalanyl-N-[[1-(
 (aminoiminomethyl)-hexahydropyridin-4-yl]methyl]-L-
 prolinamide hydrochloride)

- 10 Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-1-}$
 amidinopiperidine·HCl.

 By methods substantially equivalent to those described
 in examples 34-D, 23-A, 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-
 15 Pro-OH), 18-E, and 1-H, 0.35 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-1-}$
 1-amidinopiperidine·HCl was prepared from
 4-BocNHCH₂piperidine. The final product was purified by
 RPHPLC method 2 (98/2 (A/B) ramp to 75/25 (A/B), 150 min).


- 20 IR

¹H NMR

 FAB-MS, m/e 465 (MH⁺)

 Analysis for C₂₃H₄₀N₆O₄·2HCl:

- Calc: C, 51.39; H, 7.88; N, 15.63; Cl, 13.19;
 25 Found: C, 51.66; H, 7.98; N, 15.80; Cl, 13.48.

Example 575 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-1-amidinopiperidine}\cdot\text{HCl}$

Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-1-amidinopiperidine}\cdot\text{HCl}$.

- 10 By methods substantially equivalent to those described in examples 34-D, 23-A, 1-G (using $\text{N-(t-BuO}_2\text{CCH}_2\text{)-N-Boc-D-Cha-Pro-OH}$), 18-E, and 1-H, 0.34 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-1-amidinopiperidine}\cdot\text{HCl}$ were prepared from 4-Boc $\text{NHCH}_2\text{CH}_2\text{-piperidine}$. The final product was purified by
- 15 RPHPLC method 2 (98/2 (A/B) ramp to 75/25 (A/B), 150 min).

IR


^1H NMR

FAB-MS, m/e 479.4 (MH^+)

20 Analysis for $\text{C}_{24}\text{H}_{42}\text{N}_6\text{O}_4\cdot 2\text{HCl}$:

Calc: C, 52.26; H, 8.07; N, 15.24; Cl, 12.86;

Found: C, 52.56; H, 8.15; N, 15.37; Cl, 13.07.

Example 58

5 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3,4-dehydro-piperidine}\cdot\text{HCl}$

A) Preparation of 4-BocNHCH₂-N-methyl-pyridinium iodide.

To a stirring solution of 4-BocNHCH₂-pyridine (20 g, 96
10 mmol) in acetonitrile (200 mL) was added iodomethane (8.9 mL,
144 mmol). After 16 h, the solution was concentrated in
vacuo to give 33.8 g (96%) of a thick light yellow oil.

FD-MS, m/e 223.1 (M⁺)

15

B) Preparation of 4-BocNHCH₂-N-Fmoc-3,4-dehydro-piperidine.

To a stirring solution of 4-BocNHCH₂-N-methyl-pyridinium
iodide (7.7 g, 34 mmol) in 1,2-dichloroethane (100 mL) was
20 added 1,8-bis(dimethylamino)naphthalene (1.5 g, 6.8 mmol)
followed by 2-chloroethyl chloroformate (5.3 g, 37 mL). The
solution was heated to reflux and after 2 h, the solution was
cooled to room temperature and the solvent was removed in
vacuo and the residue was quickly flushed through a column of
25 silica gel with 20% ethyl acetate/hexanes. The organic
solvents were removed in vacuo and the residue was dissolved
in methanol (300 mL) and heated to reflux for 20 min.
Saturated aqueous NaHCO₃ (100 mL) was then added and the
solvents were removed in vacuo. The residue was dissolved in
30 water (200 mL) and washed twice with hexanes, then saturated

with solid NaCl and extracted several times with ethyl acetate. The combined ethyl acetate extracts were dried (MgSO₄), filtered, and concentrated in vacuo to give a light yellow oil which was dissolved in dichloromethane (75 mL).

- 5 To this stirring solution was then added N,N-diisopropylethylamine (2.1 mL, 12.2 mmol) followed by 9-fluorenylmethyl chloroformate (3.2 g, 12.2 mmol). After 2 h, the solvent was removed in vacuo and the residue was dissolved in ethyl acetate (250 mL) and washed twice with 1 N
10 citric acid, once with brine, twice with saturated aqueous NaHCO₃ and finally once with brine. The organic phase was then dried (MgSO₄), filtered and concentrated in vacuo and the residue was chromatographed over a silica gel column, eluting with a step gradient of 5% ethyl acetate/hexanes
15 through 50% ethyl acetate/hexanes. The product containing fractions (judged by TLC) were combined and concentrated to give 4 g (27%) of a white solid.

IR

20 ¹H NMR

FD-MS, m/e 435 (M⁺)

C) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-N-Fmoc-3,4-dehydro-piperidine.

25

By methods substantially equivalent to those described in examples 23-A and 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-OH), 2.5 g of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-N-Fmoc-3,4-dehydro-piperidine was prepared from 4-BocNHCH₂-N-Fmoc-

30 3,4-dehydro-piperidine.

IR

¹H NMR

FD-MS, m/e 799 (M⁺)

35

D) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-3,4-dehydro-piperidine.

N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-N-Fmoc-3,4-dehydro-piperidine (1.5 g, 1.9 mmol) was dissolved in morpholine (25 mL) and after stirring for 5 h, the solvent was evaporated in vacuo. The residue was dissolved in ethyl acetate and washed twice with saturated aqueous NaHCO₃, dried (MgSO₄), filtered and concentrated in vacuo. The residue was then dissolved in a small volume of chloroform and chromatographed over silica gel, eluting with a gradient of 5% to 10% A/B (A = 9:1 methanol/conc. NH₄OH; B = chloroform). The product containing fractions as judged by TLC were combined and concentrated in vacuo to give 890 mg (82%) of white solid.

¹H NMR

FD-MS, m/e 576 (MH⁺)

E) Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-3,4-dehydro-piperidine·HCl.

HCl gas was bubbled through a solution of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-3,4-dehydro-piperidine (820 mg, 1.4 mmol) and anisole (1 mL) in dioxane (25 mL) at 0 °C for 10 min. After stirring for 12 h, the solvent was removed in vacuo and the residue was dissolved in water (50 mL) and washed twice with diethyl ether. The aqueous phase was then concentrated to a volume of about 20 mL in vacuo and purified by RPHPLC (Method 2, 98/2 (A/B) through 70/30 (A/B), 2 h). The product containing fractions as judged by analytical RPHPLC were combined, partially concentrated in vacuo and lyophilized to give 442 mg (68%) of white solid.

IR

¹H NMR

FD-MS, m/e 423 (MH⁺)


137

Analysis for $C_{22}H_{36}N_4O_4 \cdot 2HCl \cdot 1.5H_2O$:

Calc: C, 50.57; H, 8.29; N, 10.72;

Found: C, 50.31; H, 8.46; N, 10.93.

5

Example 59 HO_2CCH_2 -D-Cha-Pro-4-NHCH₂CH₂-3,4-dehydro-piperidine·HCl

10

Preparation of HO_2CCH_2 -D-Cha-Pro-4-NHCH₂CH₂-3,4-dehydro-piperidine·HCl.


By methods substantially equivalent to those described in example 58, 73 mg of HO_2CCH_2 -D-Cha-Pro-4-NHCH₂CH₂-3,4-dehydro-piperidine·HCl was prepared from 4-BocNHCH₂CH₂-pyridine. The final product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

20 IR

¹H NMRIS-MS, m/e 435.2 (MH⁺)Analysis for $C_{23}H_{38}N_4O_4 \cdot 2.3HCl \cdot 3H_2O$:

Calc: C, 48.26; H, 8.15; N, 9.79; Cl, 14.24;

25 Found: C, 48.31; H, 7.93; N, 9.66; Cl, 14.56.

Example 60

5 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{CH}_2\text{-3,4-dehydro-piperidine}\cdot\text{HCl}$

Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{CH}_2\text{-3,4-dehydro-piperidine}\cdot\text{HCl}$.

10 By methods substantially equivalent to those described in example 58, 205 mg of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{CH}_2\text{-3,4-dehydro-piperidine}\cdot\text{HCl}$ was prepared from 4-BocNHCH₂CH₂CH₂-pyridine.

15 IR


¹H NMR

IS-MS, m/e 449.2 (MH⁺)

Analysis for C₂₄H₄₀N₄O₄·2.3 HCl·H₂O:

Calc: C, 52.37; H, 3.11; N, 10.18;

20 Found: C, 51.64; H, 7.72; N, 10.31; Cl, 14.69.

Example 61

25 (1R,4aR,8aR)-1-Piq-Pro-Agm·3HCl

(N-[4-[(aminoiminomethyl)amino]butyl]-1-[[4aR,8aR)-decahydro-1(R)-isoquinoliny]carbonyl]-1-prolinamide trihydrochloride)

5 The R_f values in this example were determined by silica gel thin layer chromatography (Kieselgel 60 F-254) in the following systems (v/v):

(A) Chloroform:Methanol:Acetic Acid 135:15:1

(B) Ethyl acetate:Acetic Acid:Absolute Ethanol
10 90:10:10

(C) Ethyl acetate:Hexane (70:30)

(D) Chloroform

A) N-Methoxycarbonylphenethylamine
15

To a stirred solution of phenethylamine (75.2 mL, 0.6 mol) and triethylamine (83 mL, 0.6 mol) in THF (500 mL) was added slowly methyl chloroformate 46.2 mL, 0.6 mol) dissolved in THF (50 mL). After the reaction was stirred for
20 an additional 1 h at room temperature, diethyl ether (2 L) and 1 N HCl (800 mL) were added. The organic layer was washed with water, dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to give a clear oil of pure title compound (102 g, 95%).

25

B) 2-Methoxycarbonyl-DL-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid

To a solution of N-methoxycarbonyl phenethylamine (102
30 g, 0.57 mol) in trifluoroacetic acid (300 mL) was added glyoxylic acid (63 g, 0.68 mol), and the mixture was heated to reflux temperature. After 4 h at reflux the reaction was cooled to room temperature, solvent removed in vacuo, and diethyl ether (800mL) / water (100 mL) was added to the
35 residue. The reaction mixture pH was raised to 12 with 5 N NaOH and the aqueous layer separated. To the aqueous layer was added diethyl ether (500 mL), and the solution was

acidified to pH 2.5 with 5 N HCl. The organic layer was separated, dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to afford an oil of pure title compound (107 g, 80%); FAB-MS 236 (MH⁺).

5

C) 2-Methoxycarbonyl-DL-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid t-butyl ester

To a stirred, cooled (0 °C), solution of
10 2-methoxycarbonyl-DL-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid (2) (105 g, 0.45 mol) in CH₂Cl₂ (200 mL) was added t-butanol (52 mL, 0.54 mol) and DCC (92 g, 0.45 mol). After 2 h at 0 °C and 24 h at room temperature the solvent was removed in vacuo, and ethyl acetate (800mL) / 1 N NaHCO₃
15 (300 mL) was added to the residue. The organic layer was separated, washed with water, 1.5 N citric acid, and water. The organic layer was dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to afford an oil of pure title compound (106 g, 81%); FAB-MS 292 (MH⁺); TLC R_f (A)
20 0.61; elemental analysis (calcd) C₁₆H₂₁NO₄: C, 65.96; H, 7.27; N, 4.81; Found: C, 66.24, H, 7.28, N, 4.73.

D) 2-Methoxycarbonyl-(1RS,4aSR,8aSR)-perhydroisoquinoline-1-carboxylic acid t-butyl ester

25

A solution of 2-methylcarbonyl-DL-1,2,3,4-tetrahydroisoquinoline-1-carboxylic acid t-butyl ester (105 g, 0.36 mol) in t-butanol (800 mL) was reduced over 5% Rh/Al₂O₃ (52.5 g) at 55 bar (800 psi) of hydrogen in a high pressure
30 apparatus at 50 °C for 24 hours. The reaction mixture was filtered through a pad of diatomaceous earth, and the filtrate was concentrated in vacuo. The resulting oil was dried to give pure title compound (96.5 g, 90%) FD-MS 298 (MH⁺); TLC R_f (C) 0.63.

35

E) 2-Methoxycarbonyl-(1RS,4aRS,8aRS)-perhydroisoquinoline-1-carboxylic acid ethyl ester

To a solution of 2-methoxycarbonyl-(1RS,4aSR,8aSR)-perhydoisoquinoline-1-carboxylic acid t-butyl ester (81.2 g, 273 mmol) in EtOH (500 mL) was added sodium ethoxide (21% in ethanol) (88.4 mL, 273 mmol) and the reaction mixture was refluxed (24 h). The organic solvent was evaporated in vacuo, ethyl acetate (400mL) and water (100 mL) was added to the residue. The organic layer was separated, washed twice with water, dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to afford an oil of pure title compound (70 g, .95%); FAB-MS 270 (MH⁺); TLC R_f (A) 0.61.

F) 2-Methoxycarbonyl-(1RS,4aRS,8aRS)-perhydoisoquinoline-1-carboxylic acid

To a solution of the product of step E (70 g, 260 mmol) in THF (250 mL) was added 2 N NaOH (156 mL, 312 mmol) and the reaction mixture stirred at room temperature (30 h). The organic solvent was evaporated in vacuo, diethyl ether (400 mL) and water (100 mL) was added to the residue. The aqueous layer separated and ethyl acetate (400 mL) was added. The pH of the solution was adjusted to 2.0 with 5 N HCl. The organic layer was dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to give a clear oil. The oil was crystallized from hexane (200 mL) to afford pure title compound (46.4 g, .74%); FAB-MS 242 (MH⁺); TLC R_f (A) 0.36; elemental analysis (calcd) C₁₂H₁₉NO₄: C, 59.74; H, 7.94; N, 5.81; Found: C, 59.95, H, 7.88, N, 5.54. NMR assignments were made by homonuclear decoupling, COSY, HMQC, and DEPT experiments.

G) 2-Cbz-(1RS,4aRS,8aRS)-perhydoisoquinoline-1-carboxylic acid

To a stirred solution of the product of step F (46 g, 191 mmol), at room temperature, in anhydrous CH₃CN (200 mL) under an inert atmosphere was added a solution of

iodotrimethylsilane (62.4 mL, 440 mmol) in CH₃CN (60 mL). The reaction mixture was stirred at 55 °C for 30 min and cooled to room temperature. The reaction was quenched with water (100 mL) followed by sodium metabisulfite (1 g). The
5 pH of the reaction was raised to 10.0 with 5 N NaOH, and benzyl chloroformate (27.3 mL, 191 mmol) was added dropwise while the pH maintained at 10 with 2 N NaOH. After the reaction was stirred for an additional 30 min at room temperature, the organic solvent was evaporated in vacuo, and
10 diethyl ether (200 mL) was added. The reaction was allowed to stand at room temperature (2 h) and ethyl acetate (200 mL) was added. The aqueous solution was acidified to pH 2.5 with 5 N HCl; the organic layer was separated, dried (MgSO₄), filtered, and the filtrate was concentrated in vacuo to give
15 pure title compound as an oil (39.5 g, 65%); FAB-MS 318 (MH⁺); elemental analysis (calcd) C₁₈H₂₃NO₄: C, 68.12; H, 7.30; N, 4.41; Found: C, 66.37, H, 7.52, N, 4.37.

H) 2-Cbz-(1RS,4aRS,8aRS)-perhydoisoquinoline-1-carbonyl-
20 Pro-O-t-Bu

To a stirred, cooled (0 °C) solution of the product of step G (39 g, 123 mmol) in DMF (200 mL) was added proline t-butyl ester (21.1 g, 123 mmol), 1-hydroxybenzotriazole
25 (16.6 g, 123 mmol), and DCC (25.3 g, 123 mmol). The reaction mixture was stirred for 2 h at 0 °C and 24 h at room temperature. The reaction precipitate was filtered and the filtrate concentrated in vacuo to an oil. The oil was dissolved in EtOAc (200 mL) and water (100 mL). The organic
30 layer was washed sequentially with 1 N NaHCO₃, water, 1.5 N citric acid, and water. The organic layer was dried (MgSO₄), filtered, and the filtrate evaporated to an amorphous solid of the title compound as a mixture of diastereomers (52.7 g, 91%) FAB-MS 471 (MH⁺).

35

I) 2-Cbz-(4aR,8aR)-perhydoisoquinoline-1(R)-carbonyl-Pro-OH

To a stirred solution of the product of step H (52.4 g, 111 mmol) in CH_2Cl_2 (20 mL) was added trifluoroacetic acid (70 ml) and anisole (5 ml). The reaction mixture was stirred at room temperature for 1 h and concentrated in vacuo without heating. The residue was diluted with diethyl ether (400 ml), water (100 mL), and the pH of the solution was adjusted to 10.0 with 5 N NaOH. The aqueous layer separated and ethyl acetate (300 mL) was added. The pH of the solution was adjusted to 2.5 with 5 N HCl; the organic layer was separated, dried (MgSO_4), filtered, and the filtrate was concentrated in vacuo to give a clear oil. The oil was dissolved in diethyl ether (500 mL) and (L)-(-)-alpha-methylbenzylamine was added to the solution. The solution was allowed to stand at room temperature (24 h). The resulting solid was filtered, washed with diethyl ether and dried. The solid was suspended in ethyl acetate, washed with 1.5 N citric acid, and water. The organic layer was dried (MgSO_4), filtered, and the filtrate evaporated to give the title compound as an oil (20.2 g, 44%) FAB-MS 415 (MH^+); $[\alpha]_D = 3.2^\circ$ ($C = 0.5$, MeOH); elemental analysis (calcd) $\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_5$: C, 66.65; H, 7.30; N, 6.76. Found: C, 66.38, H, 7.36, N, 6.63.

J) 2-Cbz-(4aR,8aR)-perhydroisoquinoline-1(R)-carbonyl-Pro-NH-(CH_2)₄-NH-Boc

In flask 1 the product of step I (1.06 g, 2.55 mmole) was dissolved in DMF (10 ml), cooled to -15°C , and N-methylmorpholine (0.28 ml, 2.55 mmole) was added, followed by isobutyl chloroformate (0.33 ml, 2.55 mmole). The reaction mixture was stirred at -15°C for 2 min.

In flask 2 N-Boc-1,4-diamino-butane (0.48 g, 2.55 mmole) was dissolved in DMF (10 ml), cooled to 0°C , and N-methylmorpholine (0.28 ml, 2.55 mmole) was added to the solution. The reaction mixture was stirred at 0°C for 2 min.

The contents of flask 2 was added to flask 1, and the reaction mixture was stirred for 4 h (-15 °C) and 24 h at room temperature. To the reaction was added 1 N NaHCO₃ (1 ml) and the reaction solvent was removed *in vacuo* to an oil.

5 The residue was dissolved in EtOAc (200 ml) and washed sequentially with 1.5 N citric acid, water, 1 N NaHCO₃ (100 ml), and water. The organic solution was dried (MgSO₄), filtered, and concentrated to dryness *in vacuo* to give crude title compound as a solid (1.47 g, 99%): FAB-MS 585 (MH⁺);

10 TLC R_f (A) 0.70.

K) (4aR,8aR)-Perhydroisoquinoline-1(R)-carbonyl-Pro-Agm·3HCl

To a stirred solution of the product of step J (1.4 g, 2.4 mmol) in CH₂Cl₂ (2 mL) was added trifluoroacetic acid (25 ml) and anisole (2.5 ml). The reaction mixture was stirred at room temperature for 30 min and concentrated *in vacuo* without heating. The reaction was diluted with diethyl ether (100 ml) and the supernatant decanted. The resulting oil was

20 triturated twice with diethyl ether and dried. The dried oil was dissolved in THF (20 mL), triethylamine (0.66 mL, 4.8 mmol), and bis-Cbz-S-methylisothiourea (0.859 g, 2.4 mmol) was added to the mixture. The reaction mixture was stirred at room temperature for 48 h. The organic solvent was



25 evaporated *in vacuo* and the residue was dissolved in EtOAc (200 ml) and washed sequentially with 1 N NaHCO₃ (100 ml), and water. The organic solution was dried (MgSO₄), filtered, and concentrated to dryness *in vacuo* to give a crude solid (1.5 g, 79%): TLC R_f (D) 0.33. The crude solid (1.5 g, 1.93

30 mmol) dissolved in ethanol (50 mL), water (10 mL), and 1 N HCl (5.8 mL, 5.8 mmol) was hydrogenated in the presence of 5% Pd/C catalyst (2.5 g) at ambient temperature and pressure. The catalyst was removed by filtration and the filtrate concentrated to an oil *in vacuo*. The oil was dissolved in

35 trifluoroacetic acid (10 mL), thioanisole (1.0 mL) and trifluoromethanesulfonic acid (1.0 mL) were added to the mixture. The reaction mixture was stirred at room

temperature for 0.5 h and diethyl ether (100 ml) was added. The supernatant was decanted, and the resulting oil was triturated twice with diethyl ether and dried in vacuo to give a crude solid (1.3 g). The solid (1.3 g) was dissolved in 0.05% HCl and applied to a 5 X 25 cm column of Vydac C₁₈ resin. A gradient of increasing concentrations of CH₃CN (2% to 25%) was used to elute the peptide from the column. Fractions were collected and pooled on the basis of analytical RPHPLC profile and lyophilized to afford the pure title compound (0.139 g, 15%): FAB-MS 393 (MH⁺); elemental analysis (calcd) C₂₀H₃₆N₆O₂·5HCl·3H₂O: C, 38.28. Found: C, 38.34.

Example 62



Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂C₆H₄NH₂·HCl.

By methods substantially equivalent to those described in examples 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH), 23-D, and 58-E, 0.17 g of HO₂CCH₂-D-Cha-Pro-4-NHCH₂C₆H₄NH₂·HCl was prepared from 4-nitrobenzylamine hydrochloride. The final product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

IR

¹H NMR


146

FAB-MS, m/e 431.3 (MH⁺)Analysis for C₂₃H₃₄N₄O₄·2.2 HCl·1.5 H₂O:

Calc: C, 51.37; H, 7.35; N, 10.42; Cl, 14.50;

Found: C, 50.87; H, 6.72; N, 10.41; Cl, 14.18.

5

Example 63

10

HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂C₆H₄NH₂·HClPreparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂C₆H₄NH₂·HCl.

By methods substantially equivalent to those described in example 62, 0.19 g of HO₂CCH₂-D-Cha-Pro-4-NHCH₂CH₂C₆H₄NH₂·HCl was prepared from 4-nitrophenethylamine hydrochloride. The final product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).


20 IR

¹H NMRFAB-MS, m/e 445.3 (MH⁺)Analysis for C₂₄H₃₆N₄O₄·2.2HCl·0.5H₂O:

Calc: C, 54.00; H, 7.40; N, 10.50; Cl, 14.61;

25

Found: C, 53.65; H, 7.59; N, 10.24; Cl, 14.33.

Example 64

5 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$

A) Preparation of 4-Boc₂NCH₂-3-F-C₆H₃NO₂.

To a stirring solution of 2-fluoro-4-nitrotoluene (5 g,
 10 32 mmol) in carbon tetrachloride (160 mL) was added
 N-bromosuccinimide (5.7 g, 32 mmol) followed by benzoyl
 peroxide (0.78 g, 3.2 mmol) and the solution was heated to
 reflux. After 12 h, the heat was removed and the mixture was
 diluted with carbon tetrachloride (100 mL) and washed with
 15 water. The organic phase was then diluted with ethyl acetate
 (300 mL), dried (MgSO₄), filtered and concentrated *in vacuo*.
 The residue was dissolved in tetrahydrofuran (50 mL) and
 added to a stirring solution of NaH (60% dispersion in oil;
 1.3 g, 32 mmol) and di-*t*-butyl iminodicarboxylate (6.9 g, 32
 20 mmol) in tetrahydrofuran (100 mL). After stirring overnight
 the solvent was removed *in vacuo* and the residue was
 chromatographed over silica gel, eluting with a step gradient
 of hexanes through 20% ethyl acetate/hexanes. The product
 containing fractions (judged by TLC) were combined and
 25 concentrated *in vacuo* to give 3.9 g (33%) of a white solid.

IR

¹H NMR

FD-MS, *m/e* 370 (M⁺)

30 Analysis for C₁₇H₂₃N₂O₆:

Calc: C, 55.13; H, 6.26; N, 7.56;

Found: C, 55.27; H, 6.23; N, 7.44.

B) Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$.


By methods substantially equivalent to those described
 5 in example 23-A, 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH), 23-D and 58-E, 0.44 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$ was prepared. The final product was purified using RPHPLC method 2, 98/2 (A/B) through 70/30 (A/B), 2 h).

10 IR

¹H NMRFAB-MS, m/e 449.3 (MH⁺)Analysis for C₂₃H₃₃N₄O₄F·1.3HCl:

Calc: C, 55.70; H, 6.97; N, 11.30; Cl, 9.29;

15 Found: C, 55.38; H, 6.97; N, 11.05; Cl, 9.31.

Example 65

20

 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-amidinothiophene}\cdot\text{HCl}$

(N-(carboxymethyl)-D-cyclohexylalanyl-N-

[[5-(aminoiminomethyl)thiophen-2-yl]methyl]-L-prolinamide
hydrochloride)

25 A) Preparation of 2-cyano-5-formylthiophene

To a flame-dried 3 neck 1L round bottom flask was
 added diisopropylamine (9 mL, 66 mmol) and THF (150 mL)
 under a nitrogen atmosphere. The flask was cooled to an
 30 internal temperature of -78 °C (dry ice/acetone). To this

stirring solution was added n-butyllithium (1.6 M in hexanes, 41.3 mL, 66.1 mmol) via syringe and the mixture was allowed to stir for 5 min. To this solution was added a solution of 2-thiophenecarbonitrile (6.55 g, 60 mmol) in THF (30 mL) over 10 min. The resulting bright red solution was allowed to stir at -78 °C for 45 min, at which time dimethylformamide (23.3 mL, 300 mmol) was added via syringe. This mixture was allowed to stir for 2 h at -78 °C and then solid citric acid (about 10 g) was added followed by water (60 mL). Volatile solvents were removed in vacuo and the residue was partitioned between diethyl ether and brine (200 mL each). Layers were separated and the aqueous phase was washed once with diethyl ether. The combined organic phase was washed once with brine, dried (MgSO₄), filtered and concentrated in vacuo to give a yellow solid which was purified by silica gel chromatography using an ethyl acetate/hexanes gradient (hexanes to 50% ethyl acetate/hexanes). Fractions containing pure product were pooled and concentrated in vacuo to give 6.9 g (84%) of 2-cyano-5-formyl-thiophene.

¹H NMR

B) Preparation of 2-cyano-5-(hydroxymethyl)thiophene

25

To a solution of 2-cyano-5-formyl-thiophene (6.9 g, 50 mmol) in EtOH (100 mL) was added sodium borohydride (1.9 g, 50 mmol) in portions. After 5 min of stirring, the solvent was removed in vacuo and the residue was partitioned between ethyl acetate and brine. The layers were separated and the organic phase was washed once with 1 M citric acid and once with brine, then dried (MgSO₄), filtered and concentrated in vacuo to give 6.1 g (88%) of 2-cyano-5-(hydroxymethyl)thiophene.

35

¹H NMR

FD-MS, m/e 140 (M⁺)

Analysis for C_6H_5NOS :

Calc: C, 51.78; H, 3.62; N, 10.06;

Found: C, 51.54; H, 3.62; N, 9.86.

5 C) Preparation of 2-cyano-5-(bromomethyl)thiophene

To a solution of 2-cyano-5-(hydroxymethyl)thiophene (6.0 g, 43 mmol) in THF (50 mL) was added triphenylphosphine (15.7 g, 47 mmol) and carbon
10 tetrabromide (12.3 g, 47 mmol). After stirring overnight under nitrogen atmosphere at room temperature, the solvent was removed in vacuo and the residue was dissolved in chloroform, then adsorbed onto silica gel and loaded onto a silica gel column. The product was eluted using an ethyl
15 acetate/hexanes gradient. Fractions containing pure product (as judged by TLC) were pooled and concentrated in vacuo to give 6.5 g (75%) of 2-cyano-5-(bromomethyl)thiophene.

20 1H NMRFD-MS, m/e 203 (M^+)Analysis for C_6H_4NSBr :

Calc: C, 35.66; H, 1.99; N, 6.93;

Found: C, 35.71; H, 2.03; N, 6.95.

25

D) Preparation of 2-cyano-5-(aminomethyl)thiophene·HCl

To a cold (0°C) solution of 2-cyano-5-(bromomethyl)thiophene (6.0 g, 30 mmol) in THF (50 mL)
30 under nitrogen was added NaH (60% dispersion in oil, 1.3 g, 33 mmol) in portions. To this stirring suspension was added a solution of di-t-butyl iminodicarboxylate (7.1 g, 33 mmol) in THF (50 mL) over 30 min. After stirring for 3 h, saturated aqueous ammonium chloride (100 mL) was added.
35 Volatile solvents were then removed in vacuo and the residue was partitioned between ethyl acetate and water. The layers were separated and the organic phase was washed

twice with brine, dried (MgSO_4), filtered and concentrated in vacuo to give 10.5 g (100%) of 2-cyano-5-Boc₂NCH₂-thiophene which crystallized upon standing. This solid was dissolved in EtOAc (200 mL) and cooled to 0 °C using an ice/water bath. Anhydrous HCl gas was bubbled through the solution for 10 min and the mixture was stirred for 2 h while warming to room temperature. Solvents were removed in vacuo and the resulting solid was suspended in diethyl ether and isolated by filtration. The white solid was dried overnight under vacuum to give 5.2 g (100%) of 2-cyano-5-(aminomethyl)thiophene·HCl.

¹H NMR

FD-MS, m/e 139 (M⁺)

Analysis for C₆H₇N₂SCl:

Calc: C, 41.26; H, 4.04; N, 16.04;

Found: C, 41.19; H, 4.12; N, 15.82.

E) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-cyanothiophene.

By a method substantially equivalent to that described in example 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH), 4.6 g (93%) of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-cyanothiophene was prepared from 2-cyano-5-(aminomethyl)thiophene·HCl.

IR

¹H NMR

FD-MS, m/e 602 (M⁺)

F) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-C(NH)NHBoc-thiophene.

Hydrogen sulfide gas was bubbled through a solution of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-cyanothiophene (1.5 g, 2.5 mmol) and triethylamine (4.5 mL) in pyridine (45 mL)

for 5 min and then the vessel was sealed and allowed to stand overnight. The next morning, nitrogen was bubbled through the solution for 5 min and the solvent was removed *in vacuo*. The residue was dissolved in ethyl acetate and washed once
5 with water and once with brine, then dried (MgSO_4), filtered and concentrated *in vacuo*. The residue was then dissolved in toluene and concentrated *in vacuo* two times.

The residue was then dissolved in acetone (100 mL) and iodomethane (5 mL) was added. After stirring overnight at
10 room temperature, the solvents were removed *in vacuo*. The resulting gold foam was then dissolved in methanol (20 mL), NH_4OAc (0.39 g, 5 mmol) was added, and the solution was heated to reflux. After 1 h, the solvent was removed *in vacuo* and the residue was dissolved in tetrahydrofuran (10
15 mL). To this stirring solution was added a solution of K_2CO_3 (1.73 g, 12.5 mmol) in water (10 mL) followed by di-*t*-butyl dicarbonate (2.2 g, 10 mmol). After stirring for 1 h, the suspension was diluted with ethyl acetate (400 mL) and washed with water followed by brine. The organic phase was then
20 concentrated *in vacuo* and chromatographed over silica gel, eluting with a step gradient of 10% ethyl acetate/hexanes through 75% ethyl acetate/hexanes. The product containing fractions, as judged by TLC, were combined and concentrated *in vacuo* to give 1.1 g (61%) of a white foam.

25

 ^1H NMRFD-MS, m/e 720 (M^+)Analysis for $\text{C}_{36}\text{H}_{57}\text{N}_5\text{O}_8\text{S}$:

	Calc:	C, 60.06; H, 7.98; N, 9.73;
30	Found:	C, 59.76; H, 8.07; N, 9.52.

G) Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-amidinothiophene}\cdot\text{HCl}$.

35

By a method substantially equivalent to that described in example 58-E, 500 mg of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-}$

amidinothiophene·HCl were prepared. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

IR

5 ¹H NMR

FAB-MS, m/e 464.2 (MH⁺)


Analysis for C₂₂H₃₃N₅O₄S·2HCl·H₂O:

Calc: C, 47.65; H, 6.73; N, 12.63; Cl, 12.79;

Found: C, 47.53; H, 6.57; N, 12.59; Cl, 12.67.

10

Example 66



15

HO₂CCH₂-D-Cha-Pro-2-NHCH₂-5-amidinopyridine·HCl

(N-(carboxymethyl)-D-cyclohexylalanyl-N-

[[5-(aminoiminomethyl)pyridin-2-yl]methyl]-L-prolinamide
hydrochloride)

20

A) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-2-NHCH₂-5-cyanopyridine.

By methods substantially equivalent to those described in example 64-A, 23-A and 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-

25

Cha-ProOH), 4.4 g of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-2-NHCH₂-5-cyanopyridine was prepared from 2-methyl-5-cyanopyridine.

154

IR

 ^1H NMRFD-MS, m/e 597 (M^+)

- 5 B) Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-2-NHCH}_2\text{-5-amidinopyridine}\cdot\text{HCl}$.


By methods substantially equivalent to those described in example 65-F and 65-G, 130 mg of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-2-NHCH}_2\text{-5-amidinopyridine}\cdot\text{HCl}$ was prepared from N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-2-NHCH₂-5-cyanopyridine. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

IR

15 ^1H NMRFAB-MS, m/e 459.3 (MH^+)HRMS (FAB), m/e calcd. for $\text{C}_{23}\text{H}_{35}\text{N}_6\text{O}_4$: 459.2720

Found: 459.2707

20

Example 67

$\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-2-NHCH}_2\text{-5-amidinopiperidine}\cdot\text{HCl}$

25

Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-2-NHCH}_2\text{-5-amidinopiperidine}\cdot\text{HCl}$.

N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-2-NHCH₂-5-cyanopyridine
30 (1.2 g, 2 mmol) was elaborated by methods substantially

equivalent to those described in example 23-A, 23-B, and 1-B. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h). The fractions containing the minor product as judged by analytical RPHPLC were combined, partially concentrated in vacuo and lyophilized to give 93 mg (9%) of a pale green solid.

IR

¹H NMR


10 IS-MS, m/e 465.5 (MH⁺)

HRMS (FAB), m/e calcd. for C₂₃H₄₁N₆O₄: 465.3189

Found: 465.3191

Example 68

15



HO₂CCH₂-D-Cha-Pro-2-NHCH₂-5-amidino-5,6-dehydro-piperidine·HCl

20

(N-(carboxymethyl)-D-cyclohexylalanyl-N-[[5-(aminoiminomethyl)-1,2,3,4-tetrahydropyridin-2-yl]methyl]-L-prolinamide hydrochloride)

Preparation of HO₂CCH₂-D-Cha-Pro-2-NHCH₂-5-amidino-5,6-dehydro-piperidine·HCl.

25

N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-2-NHCH₂-5-cyanopyridine (1.2 g, 2 mmol) was elaborated by methods substantially equivalent to those described in example 23-A, 23-B, and 1-B. The product was purified by RPHPLC method 2 (98/2 (A/B)

30

156


through 70/30 (A/B), 2 h). The fractions containing the major product as judged by analytical RPHPLC were combined, partially concentrated and lyophilized to give 422 mg (39%) of a white solid.

5

IR

¹H NMRIS-MS, m/e 463.3 (MH⁺)Analysis for C₂₃H₃₈N₆O₄•2.9HCl•2H₂O:

10 Calc: C, 45.71; H, 7.49; N, 13.91; Cl, 17.01;
 Found: C, 45.51; H, 6.83; N, 13.66; Cl, 16.83.

Example 69

15

HO₂CCH₂-D-Cha-Pro-3-NHCH₂-6-amidino-pyridazine•3HCl

(N-(carboxymethyl)-D-cyclohexylalanyl-N-

[[6-(aminoiminomethyl)pyridazin-3-yl]methyl]-L-prolinamide

20

trihydrochloride

A) Preparation of 3-methyl-6-cyanopyridazine.

To a stirring solution of 3-methylpyridazine (11 g, 118
 25 mmol) in dichloromethane (200 mL) was added AlCl₃ (0.05 g)
 followed by trimethylsilylcyanide (21 g, 211 mmol). After 20
 min, a solution of p-toluenesulfonyl chloride (38 g, 201
 mmol) in dichloromethane (50 mL) was added via addition
 funnel and the solution continued to stir overnight. The
 30 next morning, the solvent was removed in vacuo and the

residue was suspended in ethanol with stirring for 15 min and then filtered to give a white solid. The solid was dissolved in tetrahydrofuran (200 mL) and to this stirring solution was added 1,8-diazabicyclo[5.4.0]undec-7-ene (16 mL, 105 mmol).

- 5 After 1 h, the solvent was removed in vacuo and the residue was partitioned between hexanes and saturated aqueous NH_4Cl . The phases were separated and the aqueous phase was basified with solid Na_2CO_3 , then extracted three times with ethyl acetate. The combined ethyl acetate phases were dried
10 (MgSO_4), filtered and concentrated in vacuo to give 9 g (64%) of white solid.

IR

^1H NMR

- 15 FD-MS, m/e 119.1 (M^+)

B) Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-3-NHCH}_2\text{-6-amidino-pyridazine}\cdot\text{HCl}$.

- 20 By methods substantially equivalent to those described in example 66, 90 mg of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-3-NHCH}_2\text{-6-amidinopyridazine}\cdot\text{HCl}$ was prepared from 3-methyl-6-cyanopyridazine. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

25

IR


^1H NMR

FAB-MS, m/e 460.3 (MH^+)

Analysis for $\text{C}_{22}\text{H}_{33}\text{N}_7\text{O}_4\cdot 3\text{HCl}\cdot 2\text{H}_2\text{O}$:

- 30 Calc: C, 43.68; H, 6.66; N, 16.21;
 Found: C, 44.04; H, 6.45; N, 15.57.

Example 70



5 HO₂CCH₂-D-Cha-Pro-4-NHCH₂-1-amidino-3,4-dehydro-
 piperidine·HCl
 (N-(carboxymethyl)-D-cyclohexylalanyl-N-[[1-
(aminoiminomethyl)-1,2,3,6-tetrahydropyridin-4-yl]methyl]-
 L-prolinamide hydrochloride)

10

A) Preparation of N,N'-Boc₂-thiourea.

To a stirring suspension of NaH (60% oil dispersion, 9.4 g, 234 mmol) in tetrahydrofuran (500 mL) at 0 °C was added thiourea (4.0 g, 52 mmol). After 30 min, the cold bath was removed and the reaction was allowed to stir for 30 min at room temperature. Once again, the vessel was cooled to 0 °C and a solution of di-*t*-butyl dicarbonate (25 g, 115 mmol) in tetrahydrofuran (100 mL) was added via addition funnel. After stirring for 30 min at 0 °C and an additional 2 h at room temperature, saturated aqueous NaHCO₃ was added. The solution was then concentrated to about half the original volume *in vacuo* and ethyl acetate was added. The organic phase was then washed with saturated aqueous NaHCO₃, followed by brine and was then dried with MgSO₄, filtered and concentrated to give 11.9 g (83%) of a white solid.

IR

¹H NMR

30 FD-MS, m/e 276 (M⁺)

Analysis for $C_{11}H_{20}N_2O_4S$:

Calc: C, 47.81; H, 7.30; N, 10.14;

Found: C, 47.69; H, 7.28; N, 10.34.

- 5 B) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-1-(N,N'-Boc₂-amidino)-3,4-dehydro-piperidine.

To a stirring solution of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-3,4-dehydro-piperidine (0.6 g, 1 mmol) and
10 triethylamine (0.35 g, 3.4 mmol) in dimethylformamide (10 mL) was added N,N'-Boc₂-thiourea (0.28 g, 1 mmol) followed by
HgCl₂ (0.28 g, 1 mmol). After 4 h, the solvent was removed
in vacuo, and the residue was dissolved in ethyl acetate and
washed twice with brine. The organic phase was then dried
15 with MgSO₄, filtered and concentrated in vacuo. The product
was purified by chromatography over silica gel, eluting with
20% ethyl acetate/hexanes through 75% ethyl acetate hexanes.
The product containing fractions as judged by TLC were
combined and concentrated in vacuo to give 800 mg (94%) of a
20 white foam.

IR

¹H NMR

FD-MS, m/e 820 (MH⁺)

- 25 Analysis for $C_{42}H_{70}N_6O_{10}$:

Calc: C, 61.59; H, 8.61; N, 10.26;

Found: C, 61.81; H, 8.79; N, 10.45.

- 30 C) Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-1-amidino-3,4-dehydro-piperidine·HCl.


By methods substantially equivalent to those described
in example 58-E, 0.22 g (55%) of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-1-
amidino-3,4-dehydro-piperidine·HCl was prepared from N-(t-
35 BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-1-(N,N'-Boc₂-amidino)-3,4-
dehydro-piperidine. The product was purified by RPHPLC
method 2 (98/2 (A/B) through 70/30 (A/B), 2 h).

160

IR

 ^1H NMRFAB-MS, m/e 463.3 (MH^+)Analysis for $\text{C}_{23}\text{H}_{38}\text{N}_6\text{O}_4 \cdot 2.2\text{HCl} \cdot 2\text{H}_2\text{O}$:

5 Calc: C, 47.73; H, 7.70; N, 14.52; Cl, 13.47;
 Found: C, 47.49; H, 7.64; N, 14.55; Cl, 13.48.

Example 71

10

 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-benzamidinium}\cdot\text{HCl}$

(N-(carboxymethyl)-D-cyclohexylalanyl-N-

15 [[4-(aminoiminomethyl)-2-fluorophenyl]methyl]-L-prolinamide
 hydrochloride)


Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-benzamidinium}\cdot\text{HCl}$.

20 By methods substantially equivalent to those described
 in example 66, 0.27 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-}$
 benzamidinium $\cdot\text{HCl}$ was prepared from 3-F-4-Me-benzonitrile. The
 product was purified by RPHPLC method 2 (98/2 (A/B) through
 70/30 (A/B), 2 h).

25 IR

 ^1H NMRFAB-MS, m/e 476.3 (MH^+)Analysis for $\text{C}_{24}\text{H}_{34}\text{N}_5\text{O}_4\text{F} \cdot 2\text{HCl} \cdot 1.5\text{H}_2\text{O}$:

30 Calc: C, 50.09; H, 6.83; N, 12.17; Cl, 12.32;
 Found: C, 49.89; H, 6.65; N, 12.17; Cl, 12.42.

Example 72

5

HO₂CCH₂-D-Cha-Pro-4-NHCH₂-3,5-F₂-benzamidine·HCl
 (N-(carboxymethyl)-D-cyclohexylalanyl-N-
 [[4-(aminoiminomethyl)-2,6-difluorophenyl]methyl]-L-
 prolinamide hydrochloride)

10

Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-3,5-F₂-
 benzamidine·HCl.

By methods substantially equivalent to those described
 in example 65, 0.28 g of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-3,5-F₂-
 benzamidine·HCl was prepared from 3,5-F₂-benzonitrile. The
 product was purified by RPHPLC method 2 (98/2 (A/B) through
 70/30 (A/B), 150 min).


20 IR

¹H NMRFAB-MS, m/e 494.2 (MH⁺)Analysis for C₂₄H₃₃N₅O₄F₂·2HCl·1.5H₂O:

Calc: C, 48.57; H, 6.45; N, 11.80; Cl, 11.95;

25

Found: C, 48.26; H, 6.17; N, 11.89; Cl, 11.90.

Example 73

5 HO₂CCH₂-D-Cha-Pro-4-NHCH₂-2-aminopyridine·HCl

A) Preparation of 4-methyl-2-phthalimidopyridine.

To a stirring solution of 4-methyl-2-aminopyridine (50
 10 g, 460 mmol) in acetic acid (1 L) was added phthalic
 anhydride (68 g, 460 mmol) and the reaction was heated to
 reflux. After 12 h, acetic anhydride (43 mL, 460 mmol) was
 added and the solution continued to stir at reflux for an
 additional 48 h. The solvent was then removed *in vacuo* and
 15 the solid residue was suspended in toluene and concentrated
in vacuo twice. The solid was then suspended in ethyl
 acetate with vigorous stirring and filtered. After repeating
 this ethyl acetate washing procedure the solid was dried
 overnight *in vacuo* to give 46.6 g (42%) of a white solid.

20

IR

¹H NMR

FD-MS, m/e 238 (M⁺)

Analysis for C₁₄H₁₀N₂O₂:

25 Calc: C, 70.58; H, 4.23; N, 11.76;

 Found: C, 70.42; H, 4.29; N, 11.70.

B) Preparation of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-phthalimidopyridine.

30

By methods substantially equivalent to those described in example 64-A, 23-A and 1-G (using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH), 2.4 g of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-phthalimidoopyridine was prepared from 4-methyl-2-phthalimidoopyridine.

IR

¹H NMR

FD-MS, m/e 717.7 (M⁺)

10

C) Preparation of HO₂CCH₂-D-Cha-Pro-4-NHCH₂-2-aminopyridine·HCl.

To a stirring solution of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-Pro-4-NHCH₂-2-phthalimidoopyridine (1.6 g, 2.2 mmol) in ethanol (25 mL) was added hydrazine hydrate (0.52 mL, 10.4 mmol). After 1 h, the solvents were removed in vacuo and the residue was dissolved in ethyl acetate and concentrated in vacuo twice. The residue was elaborated by a procedure substantially equivalent to that described in example 58-E to give 380 mg (37%) of a white solid. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 150 min).

IR

25 ¹H NMR


FAB-MS, m/e 432.3 (MH⁺)

Analysis for C₂₂H₃₃N₅O₄·2.1 HCl·H₂O:

Calc: C, 50.23; H, 7.11; N, 13.31;

Found: C, 50.05; H, 7.08; N, 13.54.

30

Example 74

HO₂CCH₂-D-Cha-Pro-5-NHCH₂-2-aminopyridine·HCl

5

Preparation of HO₂CCH₂-D-Cha-Pro-5-NHCH₂-2-aminopyridine·HCl.

By methods substantially equivalent to those described
in example 73, 0.88 g of HO₂CCH₂-D-Cha-Pro-5-NHCH₂-2-
10 aminopyridine·HCl was prepared from 5-methyl-2-aminopyridine.
The product was purified by RPHPLC method 2 (98/2 (A/B)
through 70/30 (A/B), 150 min).

IR

15 ¹H NMR


FAB-MS; m/e 432.3 (MH⁺)

Analysis for C₂₂H₃₃N₅O₄·2HCl·H₂O:

Calc: C, 50.58; H, 7.14; N, 13.40;

Found: C, 50.79; H, 7.20; N, 13.58.

20

Example 75

HO₂CCH₂-D-Cha-Pro-5-NHCH₂CH₂-2-aminopyridine·HCl

A) Preparation of 5-Me-2-Boc₂N-pyridine.

To a stirring solution of 5-methyl-2-aminopyridine (10.5 g, 100 mmol) in dichloromethane (200 mL) at 0 °C was added
5 N,N-diisopropylethylamine (25.8 g, 200 mmol), followed by
di-t-butyl dicarbonate (55 g, 250 mmol) and finally 4-(N,N-
dimethylamino)pyridine (12.2 g, 100 mmol). The cold bath was
removed and the solution was allowed to continue stirring
overnight. The mixture was then diluted with ethyl acetate
10 (600 mL) and washed three times with saturated aqueous NH₄Cl,
once with brine, twice with saturated aqueous NaHCO₃, and
once again with brine. The organic phase was then dried
(MgSO₄), filtered, concentrated *in vacuo* and chromatographed
over silica gel eluting with 10% ethyl acetate/hexanes
15 through 75% ethyl acetate/hexanes. The fractions containing
product, as judged by TLC, were combined and concentrated *in vacuo*
to give 12.8 g (42%) of a white solid.

IR

20 ¹H NMRFD-MS, m/e 308 (M⁺)Analysis for C₁₆H₂₄N₂O₄:

Calc: C, 62.32; H, 7.84; N, 9.08;

Found: C, 62.51; H, 8.11; N, 9.37.

25

B) Preparation of 5-BrCH₂-2-Boc₂N-pyridine.

By methods substantially equivalent to those described
in example 64-A, approximately 11.6 g of 5-BrCH₂-2-Boc₂N-
30 pyridine (which was contaminated with starting material) was
prepared from 5-Me-2-Boc₂N-pyridine.

¹H NMRFD-MS, m/e 386.3 (M⁺)35 Analysis for C₁₆H₂₃N₂O₄Br:

Calc: C, 49.62; H, 5.99; N, 7.23;

Found: C, 49.86; H, 6.00; N, 7.07.

C) Preparation of 5-NCCH₂-2-Boc₂N-pyridine.

To a stirring solution of slightly impure 5-BrCH₂-2-Boc₂N-pyridine (9.7 g, 25 mmol) in dimethylformamide (150 mL) was added 18-crown-6 (1.32 g, 5 mmol) followed by KCN (1.95 g, 30 mmol). After stirring for 6 h, the solvent was removed *in vacuo* and the residue was chromatographed over silica gel, eluting with a step gradient of hexanes through 40% ethyl acetate/hexanes. The product containing fractions as judged by TLC, were combined and concentrated *in vacuo* to give a 2.6 g (31% over 2 steps) of a white solid.

IR

15 ¹H NMRFD-MS, m/e 333.4 (M⁺)Analysis for C₁₇H₂₃N₃O₄:

Calc: C, 61.25; H, 6.95; N, 12.60;

Found: C, 61.09; H, 6.92; N, 12.53.

20

D) Preparation of HO₂CCH₂-D-Cha-Pro-5-NHCH₂CH₂-2-aminopyridine·HCl.

To a stirring solution of 5-NCCH₂-2-Boc₂N-pyridine (2.5 g, 7.5 mmol) in methanol (150 mL) was added CoCl₂ (0.97 g, 7.5 mmol) and water (0.81 g, 45 mmol). After 5 min, NaBH₄ (2.84 g, 75 mmol) was added in small portions over 15 min. After an additional 15 min, the solvent was removed *in vacuo* and the residue was dissolved in concentrated aqueous NH₄OH and extracted several times with ethyl acetate. The combined ethyl acetate extracts were dried with MgSO₄, filtered and concentrated *in vacuo*.


Then, by methods substantially equivalent to those described in example 1-A and 73-C, the residue was elaborated to give 1.2 g (33%) of HO₂CCH₂-D-Cha-Pro-5-NHCH₂CH₂-2-aminopyridine·HCl. The product was purified by RPHPLC method 2 (98/2 (A/B) through 60/40 (A/B), 150 min).

167

IR

 ^1H NMRFAB-MS, m/e 446.3 (MH^+)5 HRMS (FAB) calc. for $\text{C}_{23}\text{H}_{36}\text{N}_5\text{O}_4$: 446.2767

Found: 446.2769

Example 76

10

 $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-2-aminopyridine}\cdot\text{HCl}$ A) Preparation of 4-BocNHCH₂CH₂-2-CN-pyridine.

15

To a solution of 4-BocNHCH₂CH₂-pyridine (2.22 g, 10 mmol) in acetone (50 mL) was added a solution of m-chloro-perbenzoic acid (5.2 g, 30 mmol) in acetone (50 mL) via an addition funnel over 10 min. After stirring overnight, the solvent was removed in *vacuo* and the residue was partitioned between water (100 mL) and diethyl ether (100 mL). The organic phase was separated and extracted three times with water. The combined aqueous phase was then saturated with solid NaCl and extracted three times with dichloromethane (100 mL). The combined dichloromethane extracts were washed once with brine, dried with Na_2SO_4 , filtered and concentrated to a small volume in *vacuo* and then diethyl ether was added. The white precipitate (2.0 g) was filtered and dried in *vacuo*.

20

25

One half of the isolated solid (4.2 mmol) was dissolved in dichloromethane (10 mL) and to this stirring solution was added trimethylsilyl cyanide (0.84 mL, 6.3 mmol) followed by N,N-dimethylcarbamoyl chloride (0.58 mL, 6.3 mmol). After
5 stirring overnight, 1 M aq. KHCO_3 (1 mL) was added slowly and the mixture was partitioned between ethyl acetate and water. The organic phase was then washed twice with brine, dried with MgSO_4 , filtered and concentrated in vacuo to give 0.6 g (58%) of an amber oil which crystalized upon standing.

10

IR

 ^1H NMRFD-MS, m/e 247 (M^+)Analysis for $\text{C}_{13}\text{H}_{17}\text{N}_3\text{O}_2$:

15

Calc: C, 63.31; H, 7.02; N, 16.99;

Found: C, 63.31; H, 7.02; N, 16.71.

B) Preparation of 4-BocNHCH₂CH₂-2-CbzNH-pyridine.

20

To a stirring solution of 4-BocNHCH₂CH₂-2-CN-pyridine (0.5 g, 2 mmol) in methanol (2.4 mL), was added 5 N NaOH (1.6 mL, 8 mmol) and the solution was heated to reflux. After 24 h, the solution was cooled to room temperature and allowed to stir for an additional 48 h. The pH was then adjusted to 7
25 with 1 N HCl and the solvents were removed in vacuo.

The residue was suspended in toluene (50 mL) and heated to reflux. To this stirring solution was sequentially added, triethylamine (0.36 mL, 2.6 mmol), benzyl alcohol (0.27 mL, 2.6 mmol) and diphenylphosphoryl azide (0.72 g, 2.6 mmol).

30

After stirring at reflux overnight, the solution was allowed to cool and then diluted with ethyl acetate (200 mL) and washed twice with saturated aqueous NH_4Cl and twice with brine. The organic phase was then dried with MgSO_4 , filtered and concentrated in vacuo. The residue was then

35

chromatographed over silica gel eluting with a step gradient of hexanes through 50% ethyl acetate/hexanes. The product

containing fractions as determined by TLC were combined and concentrated in vacuo to give 0.37 g (50%) of a white solid.

^1H NMR

5 FD-MS, m/e 371.2 (M^+)

Analysis for $\text{C}_{20}\text{H}_{25}\text{N}_3\text{O}_4$:

Calc: C, 64.67; H, 6.78; N, 11.31;

Found: C, 64.90; H, 7.07; N, 11.06.

10 C) Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-2-aminopyridine}\cdot\text{HCl}$.

By methods substantially equivalent to those described in example 23-A, 1-G (using $\text{N-(t-BuO}_2\text{CCH}_2)\text{-N-Boc-D-Cha-ProOH}$),
 15 23-D and 58-E, 48 mg of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{CH}_2\text{-2-aminopyridine}\cdot\text{HCl}$ was prepared from 4-BocNHCH₂CH₂-2-CbzNH-pyridine. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 150 min).

20 ^1H NMR

FAB-MS, m/e 446.4 (MH^+)


Analysis for $\text{C}_{23}\text{H}_{35}\text{N}_5\text{O}_4\cdot 1.5\text{HCl}\cdot\text{H}_2\text{O}$:

Calc: C, 53.30; H, 7.49; N, 13.51;

Found: C, 53.64; H, 7.27; N, 13.80.

25

Example 77



30

$\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-F-aniline}\cdot\text{HCl}$

Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-F-aniline}\cdot\text{HCl}$.

By methods substantially equivalent to those described
 5 in example 64, 0.55 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-2-F-}$
 $\text{aniline}\cdot\text{HCl}$ was prepared from 3-F-4- NO_2 -toluene. The product
 was purified by RPHPLC method 2 (98/2 (A/B) through 60/40
 (A/B), 180 min).

10 IR

^1H NMR


FAB-MS, m/e 449.3 (MH^+)

Analysis for $\text{C}_{23}\text{H}_{33}\text{N}_4\text{O}_4\text{F}\cdot 0.9\text{HCl}\cdot\text{H}_2\text{O}$:

Calc: C, 55.32; H, 7.25; N, 11.22; Cl, 6.39;

15 Found: C, 55.49; H, 6.93; N, 11.15; Cl, 6.23.

Example 78



20

$\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2\cdot\text{HCl}$

Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2\cdot\text{HCl}$.

25 By methods substantially equivalent to those described
 in example 10, using $\text{N-(t-BuO}_2\text{CCH}_2\text{)-N-Boc-D-Cha-ProOH}$ in place
 of Boc-D-Phe-ProOH , 0.53 g of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-}$
 $\text{NHCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2\cdot\text{HCl}$ was prepared. The product was purified by
 RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 150 min).

30

IR

¹H NMR

FD-MS, m/e 445.4 (MH⁺)


5 Analysis for $C_{24}H_{36}N_4O_4 \cdot 2.2HCl \cdot 0.5H_2O$:

Calc: C, 54.00; H, 7.40; N, 10.50; Cl, 14.61;

Found: C, 54.18; H, 7.54; N, 10.31; Cl, 14.86.

Example 79

10


$$\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_{10}\text{CH}_2\text{NH}_2 \cdot \text{HCl}$$

15 Preparation of $\text{HO}_2\text{CCH}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_{10}\text{CH}_2\text{NH}_2\cdot\text{HCl}$.

By methods substantially equivalent to those described in example 12, using N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH in place of Boc-D-Phe-ProOH, 0.04 g of HO₂CCH₂-D-Cha-Pro-4-

20 $\text{NHCH}_2\text{C}_6\text{H}_{10}\text{CH}_2\text{NH}_2 \cdot \text{HCl}$ was prepared. The product was purified by RPHPLC method 2 (98/2 (A/B) through 70/30 (A/B), 150 min.).

¹H NMR


FD-MS, m/e 451 (MH⁺)

25 Analysis for $C_{24}H_{42}N_4O_4 \cdot 2.7HCl \cdot 0.5H_2O$:

Calc: C, 51.65; H, 8.25; N, 10.04;

Found: C, 51.47; H, 7.87; N, 9.97.

172

Example 80

5 EtSO₂-D-Cha-Pro-4-NHCH₂C₆H₄C(NH)NH₂ · HCl
 (N-(ethylsulfonyl)-D-cyclohexylalanyl-N-
 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
 hydrochloride)

10 Preparation of EtSO₂-D-Cha-Pro-4-NHCH₂C₆H₄C(NH)NH₂ · HCl.

By methods substantially equivalent to those described
 in example 18, using EtSO₂-D-Cha-ProOH in place of Cbz-D-1-
 Piq-ProOH, 3.6 g of EtSO₂-D-Cha-Pro-4-NHCH₂C₆H₄C(NH)NH₂ · HCl
 15 was prepared. The product was purified by RPHPLC method 2,
 (90/10 (A/B) through 50/50 (A/B), 180 min).

IR

¹H NMR


20 FAB-MS, m/e 492.3 (MH⁺)

Analysis for C₂₄H₃₇N₅O₄S · HCl:

Calc: C, 54.58; H, 7.25; N, 13.26; Cl, 6.71;

Found: C, 54.31; H, 7.31; N, 13.37; Cl, 6.71.

25

Example 81

EtSO₂-D-Cha-Pro-4-NHCH₂C₆H₄CH₂NH₂ · HCl

Preparation of $\text{EtSO}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2\cdot\text{HCl}$.

By methods substantially equivalent to those described in example 10, using $\text{EtSO}_2\text{-D-Cha-ProOH}$ in place of Boc-D-Phe-ProOH , $\text{EtSO}_2\text{-D-Cha-Pro-4-NHCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2\cdot\text{HCl}$ was prepared. The product was purified by RPHPLC method 2, (90/10 (A/B) through 50/50 (A/B), 180 min).

IR

$^1\text{H NMR}$


FAB-MS, m/e 479.4 (MH^+)

Analysis for $\text{C}_{24}\text{H}_{38}\text{N}_4\text{O}_4\text{S}\cdot\text{HCl}\cdot\text{H}_2\text{O}$:

Calc: C, 54.07; H, 7.75; N, 10.51; Cl, 6.65;

Found: C, 54.13; H, 7.44; N, 10.51; Cl, 6.78.

Example 82



20

$\text{EtSO}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$

Preparation of $\text{EtSO}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$.

By methods substantially equivalent to those described in example 64, using $\text{EtSO}_2\text{-D-Cha-ProOH}$ in place of $\text{N-(t-BuO}_2\text{CCH}_2\text{)-N-Boc-D-Cha-ProOH}$, 0.53 g of $\text{EtSO}_2\text{-D-Cha-Pro-4-NHCH}_2\text{-3-F-C}_6\text{H}_3\text{NH}_2\cdot\text{HCl}$ was prepared. The product was purified by RPHPLC method 2, (90/10 (A/B) through 50/50 (A/B), 180 min).

30

IR

$^1\text{H NMR}$

FAB-MS, m/e 483.3 (MH⁺)


Analysis for C₂₃H₃₅N₄O₄SF•1.1HCl•0.5H₂O:

Calc: C, 51.95; H, 7.03; N, 10.54; Cl, 7.33;

Found: C, 52.09; H, 6.94; N, 10.39; Cl, 7.24.

5

Example 83



10

EtSO₂-D-Cha-Pro-4-NHCH₂-2-aminopyridine·HCl

Preparation of EtSO₂-D-Cha-Pro-4-NHCH₂-2-aminopyridine·HCl.

By methods substantially equivalent to those described in example 73, using EtSO₂-D-Cha-ProOH in place of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH, 0.22 g of EtSO₂-D-Cha-Pro-4-NHCH₂-2-aminopyridine·HCl was prepared from 4-methyl-2-aminopyridine. The product was purified by RPHPLC method 2, (90/10 (A/B) through 50/50 (A/B), 180 min.).

20

¹H NMR

FAB-MS, m/e 466.4 (MH⁺)


Analysis for C₂₂H₃₅N₅O₄S•1.1HCl:

Calc: C, 52.25; H, 7.19; N, 13.85; Cl, 7.71;

25

Found: C, 52.49; H, 6.96; N, 13.96; Cl, 7.76.

175

Example 84

5 EtSO₂-D-Cha-Pro-5-NHCH₂-2-aminopyridine·HCl

Preparation of EtSO₂-D-Cha-Pro-5-NHCH₂-2-aminopyridine·HCl.

By methods substantially equivalent to those described
 10 in example 73, using EtSO₂-D-Cha-ProOH in place of N-(t-BuO₂CCH₂)-N-Boc-D-Cha-ProOH, 0.24 g of EtSO₂-D-Cha-Pro-5-NHCH₂-2-aminopyridine·HCl was prepared from 5-methyl-2-aminopyridine. The product was purified by RPHPLC method 2, (90/10 (A/B) through 50/50 (A/B), 180 min).


15

¹H NMR

FAB-MS, m/e 466.4 (MH⁺)

Analysis for C₂₂H₃₅N₅O₄S·1.15HCl:

Calc:	C, 52.06; H, 7.18; N, 13.80; Cl, 8.03;
20 Found:	C, 52.38; H, 6.97; N, 14.20; Cl, 8.46.

Example 85

HOOCCH₂CH₂CH₂-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
(N-(3-carboxypropyl)-D-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
hydrochloride)

A) Preparation of Cbz-MeOOCCH=CHCH₂-D-Cha-Pro-p-
NHCH₂C₆H₄C(NH)NHCbz

To a solution of D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NHCbz·2HCl
(2.5 g, 4.1 mmol) in DMF (50 mL) was added N,N-
diisopropylethylamine (2.2 mL, 12.2 mmol) and methyl
3-bromocrotonate (0.95 g, 4.5 mmol). After stirring for 48
h, Cbz-Cl (0.7 mL, 5 mmol) and additional N,N-
diisopropylethylamine (0.85 mL, 5 mmol) were added. After an
additional 16 h of stirring, the volatiles were removed in
vacuo. The residue was partitioned between ethyl acetate
(100 mL) and saturated aqueous ammonium chloride (100 mL).
The layers were separated and the organic phase was washed
once with saturated aqueous ammonium chloride (50 mL), twice
with saturated aqueous sodium bicarbonate (50 mL) and twice
with brine (50 mL). The organic phase was dried (MgSO₄),
filtered, and concentrated in vacuo. The crude residue was
purified by flash chromatography (ethyl acetate/hexanes
gradient) to yield 660 mg (22%) of a white foam.

¹H NMR

FD-MS, m/e 766 (M⁺)

Analysis for C₄₃H₅₁N₅O₈:

Calc: C, 67.43; H, 6.71; N, 9.14;
Found: C, 67.22; H, 6.57; N, 8.98.

B) Preparation of HOOCCH₂CH₂CH₂-D-Cha-Pro-p-
NHCH₂C₆H₄C(NH)NH₂·HCl

35


To a solution of Cbz-MeOOCCH=CHCH₂-D-Cha-Pro-p-
NHCH₂C₆H₄C(NH)NHCbz (0.5 g, 0.65 mmol) in ethanol (5 mL) was

177

added 1 N sodium hydroxide (0.65 mL). After stirring for 5 h at room temperature, 1 N HCl (3 mL), 10% Pd/C (0.5 g), H₂O (15 mL) and ethanol (25 mL) were added. The stirring suspension was degassed under vacuum, then placed under an atmosphere of hydrogen for 18 h. Diatomaceous earth was added and the slurry was filtered. The filtrate was concentrated *in vacuo* and purified by RPHPLC (method 2, 98/2 (A/B), ramp to 75/25 (A/B) over 150 minutes). Fractions containing pure product were pooled and lyophilized to give 46 mg (13%).

¹H NMRFAB-MS, m/e 486.3 (MH⁺)Analysis for C₂₆H₃₉N₅O₄·2.1HCl:

Calc: C, 55.15; H, 7.35; N, 12.19; Cl, 13.24;
Found: C, 55.55; H, 7.37; N, 12.19; Cl, 13.58.

Example 86

HOOCCH₂-D-Chg-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
(N-(carboxymethyl)-D-cyclohexylglycyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
hydrochloride)

A). Preparation of D-cyclohexylglycine

By a method substantially equivalent to that described in Example 53-B, 16.1 g (16%) of D-cyclohexylglycine was prepared starting from D-phenylglycine.

5 ¹H NMRFD-MS, m/e 117 (M⁺)Analysis for C₈H₁₅NO₂:

Calc: C, 61.12; H, 9.62; N, 8.91;

Found: C, 61.23; H, 9.56; N, 8.73.

10

B) Preparation of Boc-D-cyclohexylglycine

By a method substantially equivalent to that described in Example 17-A (using di-tert-butyl dicarbonate), 22 g (90%)
15 of Boc-D-cyclohexylglycine was prepared.

¹H NMRFD-MS, m/e 258 (M⁺)Analysis for C₁₃H₂₃NO₄:

20 Calc: C, 60.68; H, 9.01; N, 5.44;

Found: C, 60.91; H, 9.18; N, 5.38.

C) Preparation of Boc-Pro-p-NHCH₂C₆H₄C(NH)NHCbz

25 By a method substantially equivalent to that described in Example 1-G, 20.5 g (76%) of Boc-Pro-p-NHCH₂C₆H₄C(NH)NHCbz was prepared from Boc-Pro-OH and NH₂CH₂C₆H₄C(NH)NHCbz·2HCl.

¹H NMR30 FD-MS, m/e 481 (M⁺)Analysis for C₂₆H₃₂N₄O₅:

Calc: C, 64.98; H, 6.71; N, 11.66;

Found: C, 64.76; H, 6.78; N, 11.62.

35 D) Preparation of Pro-p-NHCH₂C₆H₄C(NH)NHCbz·2HCl

By a method substantially equivalent to that described in Example 23-A, 18.4 g (100%) of Pro-p-NHCH₂C₆H₄C(NH)NHCbz·2HCl was prepared.

5 ¹H NMR

FD-MS, m/e 381 (M⁺)

Analysis for C₂₁H₂₆N₄O₃Cl₂:

Calc: C, 55.63; H, 5.78; N, 12.36;

Found: C, 54.19; H, 6.27; N, 12.15.

10

E) Preparation of Boc-D-Chg-Pro-p-NHCH₂C₆H₄C(NH)NHCbz

By a method substantially equivalent to that described in Example 1-G, 3.6 g (97%) of Boc-D-Chg-Pro-p-NHCH₂C₆H₄C(NH)NHCbz were prepared.

15

¹H NMR

FD-MS, m/e 619 (M⁺)

Analysis for C₃₄H₄₅N₅O₆:

20 Calc: C, 65.89; H, 7.32; N, 11.30;

Found: C, 67.59; H, 8.07; N, 10.99.

F) Preparation of Cbz-t-BuOOCCH₂-D-Chg-Pro-p-NHCH₂C₆H₄C(NH)NHCbz

25

By methods substantially equivalent to those described in Examples 23-A and 85-A (using t-butyl bromoacetate and benzyl chloroformate), 1.6 g (45%) of N-Cbz-N-(t-BuOOCCH₂)-D-Chg-Pro-p-NHCH₂C₆H₄C(NH)NHCbz were prepared.

30

¹H NMR

FD-MS, m/e 769 (M⁺)

Analysis for C₄₃H₅₃N₅O₈:

Calc: C, 67.26; H, 6.96; N, 9.12;

35 Found: C, 67.50; H, 6.97; N, 9.11.

180


G) Preparation of $\text{HOOCCH}_2\text{-D-Chg-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

By methods substantially equivalent to those described in Examples 23-A (using dioxane as solvent) and 18-F, 411 mg (61%) of $\text{HOOCCH}_2\text{-D-Chg-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$ were prepared.

 ^1H NMRFAB-MS, m/e 444.3 (MH^+)10 Analysis for $\text{C}_{23}\text{H}_{33}\text{N}_5\text{O}_4\cdot 2.5\text{HCl}$:

Calc: C, 51.67; H, 6.69; N, 13.10;

Found: C, 51.84; H, 6.50; N, 13.15.

Example 87 $\text{HOOCCH}_2\text{-D-hPhe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

(N-(carboxymethyl)-D-homophenylalanyl-N-

20 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
hydrochloride)A) Preparation of $\text{HOOCCH}_2\text{-D-hPhe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$


By methods substantially equivalent to those described in Example 86, 335 mg of $\text{HOOCCH}_2\text{-D-hPhe-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$ were prepared starting from Boc-D-hPhe-OH.

181

¹H NMRFAB-MS, m/e 466.3 (MH⁺)Analysis for C₂₅H₃₁N₅O₄·2.1HCl·H₂O:

Calc: C, 53.61; H, 6.32; N, 12.50; Cl, 13.29;

5 Found: C, 53.58; H, 6.08; N, 12.59; Cl, 13.67.

Example 88

10

HOOCCH₂-D-hCha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
 (N-(carboxymethyl)-D-homocyclohexylalanyl-N-
 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
 hydrochloride)

15

A) Preparation of Boc-D-hCha-OH

By a method substantially equivalent to that described
 in Example 53-D, 5.1 g (100%) of Boc-D-hCha-OH was prepared
 20 from Boc-D-hPhe-OH.

¹H NMRFD-MS, m/e 240 (M⁺)Analysis for C₁₅H₂₇NO₄:

25 Calc: C, 63.13; H, 9.54; N, 4.91;

Found: C, 63.38; H, 9.39; N, 5.12.

B) Preparation of HOOCCH₂-D-hCha-Pro-p-
 NHCH₂C₆H₄C(NH)NH₂·HCl

30

By methods substantially equivalent to those described in Example 86, 135 mg of $\text{HOOCCH}_2\text{-D-hCha-Pro-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$ were prepared.

5 ^1H NMR

FAB-MS, m/e 472.3 (MH^+)


Analysis for $\text{C}_{25}\text{H}_{37}\text{N}_5\text{O}_4\cdot 2.2\text{HCl}\cdot 0.5\text{H}_2\text{O}$:

Calc: C, 53.54; H, 7.22; N, 12.49; Cl, 13.91;

Found: C, 53.29; H, 7.01; N, 12.46; Cl, 14.30.

10

Example 89



15 $\text{HOOCCH}_2\text{-Gly-N-C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{Gly-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

A) Preparation of $\text{HOOCCH}_2\text{-Gly-N-C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{Gly-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$

20 By methods substantially equivalent to those described in Examples 1-G, 1-D, 1-G, 23-A, 85-A, and 18-F, 365 mg of $\text{N-HOOCCH}_2\text{-Gly-N-C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{Gly-p-NHCH}_2\text{C}_6\text{H}_4\text{C(NH)NH}_2\cdot\text{HCl}$ were prepared.

25 ^1H NMR


FAB-MS, m/e 426.2 (MH^+)

Analysis for $\text{C}_{22}\text{H}_{27}\text{N}_5\text{O}_4\cdot 2.2\text{HCl}\cdot 1.5\text{H}_2\text{O}$:

Calc: C, 49.60; H, 6.09; N, 13.15; Cl, 14.64;

Found: C, 49.79; H, 5.71; N, 13.31; Cl, 14.49.

30

Example 90

5 (C₂H₅)₂CHCO-Glu-Pro-p-NHCH₂C₆H₄C(NH)NH₂ · HCl

A) Preparation of Boc-(γ-OBn)-Glu-Pro-p-NHCH₂C₆H₄C(NH)NHCBz

10 By a method substantially equivalent to that described in Example 1-G, 2.7 g (64%) of Boc-(γ-OBn)-Glu-Pro-p-NHCH₂C₆H₄C(NH)NHCBz were prepared starting from Boc-(γ-OBn)-Glu-OH and Pro-p-NHCH₂C₆H₄C(NH)NHCBz · 2HCl.

15 ¹H NMR

FD-MS, m/e 700 (M⁺)

Analysis for C₃₈H₄₅N₅O₈:

Calc: C, 65.22; H, 6.48; N, 10.01;

Found: C, 65.00; H, 6.56; N, 10.06.

20

B) Preparation of (γ-OBn)-Glu-Pro-p-NHCH₂C₆H₄C(NH)NHCBz · 2HCl

25 By a method substantially equivalent to that described in Example 23-A, 2.38 g (98%) of (γ-OBn)-Glu-Pro-p-NHCH₂C₆H₄C(NH)NHCBz · 2HCl were prepared.

¹H NMR

FD-MS, m/e 600 (M⁺)

30 Analysis for C₃₃H₃₉N₅O₆Cl₂:

184

Calc: C, 58.93; H, 5.84; N, 10.41;
Found: C, 58.64; H, 6.00; N, 10.63.

C) Preparation of (C₂H₅)₂CHCO-(γ-OBn)-Glu-Pro-p-
5 NHCH₂C₆H₄C(NH)NHCbz

To a stirring mixture of (γ-OBn)-Glu-Pro-p-
NHCH₂C₆H₄C(NH)NHCbz·2HCl (1.3 g, 2 mmol) in THF/H₂O (50 mL
each) was added K₂CO₃ (1.38 g, 10 mmol) and 2-ethylbutyryl
10 chloride (0.3 g, 2.2 mmol). After stirring 10 min,
volatiles were removed in vacuo. The resulting residue was
partitioned between water and ethyl acetate (100 mL each).
Layers were separated and the organic phase was washed
twice with each saturated aqueous ammonium chloride and
15 brine, dried (MgSO₄), filtered and concentrated in vacuo to
give 1.45 g (100%).

¹H NMR

FD-MS, m/e 698 (M⁺)

20 Analysis for C₃₉H₄₇N₅O₇:

Calc: C, 67.13; H, 6.79; N, 10.04;
Found: C, 67.11; H, 6.70; N, 9.74.

D) Preparation of (C₂H₅)₂CHCO-Glu-Pro-p-
25 NHCH₂C₆H₄C(NH)NH₂·HCl


By a method substantially equivalent to that described
in example 18-F, 425 mg (47%) of (C₂H₅)₂CHCO-Glu-Pro-p-
NHCH₂C₆H₄C(NH)NH₂·HCl were prepared. HPLC Method 2 (Ramp
30 98/2 A/B to 75/25 A/B over 150 min) was used to purify the
product.

¹H NMR

FAB-MS, m/e 474.3 (MH⁺)

35 Analysis for C₂₄H₃₅N₅O₅·1.5HCl·1.1H₂O:

Calc: C, 51.10; H, 6.91; N, 12.41; Cl, 9.43;
Found: C, 51.10; H, 6.81; N, 12.41; Cl, 9.62.

Example 91

5

$(C_2H_5)_2CHCO-Met(O_2)-Pro-p-NHCH_2C_6H_4C(NH)NH_2 \cdot HCl$

A) Preparation of $(C_2H_5)_2CHCO-Met(O_2)-Pro-p-NHCH_2C_6H_4C(NH)NH_2 \cdot HCl$

10

By methods substantially equivalent to those described in Example 90, 530 mg of $(C_2H_5)_2CHCO-Met(O_2)-Pro-p-NHCH_2C_6H_4C(NH)NH_2 \cdot HCl$ were prepared.

15 1H NMR


FAB-MS, m/e 508.2 (MH^+)

Analysis for $C_{24}H_{37}N_5O_5S \cdot 1.1HCl$:

Calc: C, 52.63; H, 7.01; N, 12.79; Cl, 7.12;

Found: C, 52.42; H, 7.03; N, 12.80; Cl, 6.99.

20

Example 92

MeSO₂CH₂CO-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl
(N-(methylsulfonylacetyl)-L-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide
hydrochloride)

5

Preparation of MeSO₂CH₂CO-D-Cha-Pro-p-NHCH₂C₆H₄C(NH)NH₂·HCl

By methods substantially equivalent those described in
Examples 1-G and 18-F, 550 mg of MeSO₂CH₂CO-D-Cha-Pro-p-
10 NHCH₂C₆H₄C(NH)NH₂·HCl were prepared.

¹H NMR

FAB-MS, m/e 520.5 (MH⁺)

Analysis for C₂₅H₃₇N₅O₅S·1.2HCl·H₂O:

15 Calc: C, 51.64; H, 6.97; N, 12.04; Cl, 7.32;
 Found: C, 51.58; H, 6.84; N, 12.18; Cl, 7.61.

The compounds of the invention are believed to
selectively inhibit thrombin over other proteinases and
20 nonenzyme proteins involved in blood coagulation without
appreciable interference with the body's natural clot lysing
ability (the compounds have a low inhibitory effect on
fibrinolysis). Further, such selectivity is believed to
permit use with thrombolytic agents without substantial
25 interference with thrombolysis and fibrinolysis. Further,
the compounds of the present invention are believed to be
orally active.

The invention in one of its aspects provides a
method of inhibiting thrombin in mammals comprising
30 administering to a mammal in need of treatment an effective
(thrombin inhibiting) dose of a compound of Formula I.

The thrombin inhibition contemplated by the present
method includes both medical therapeutic and/or prophylactic
treatment as appropriate.

35 In a further embodiment the invention relates to
treatment, in a human or animal, of conditions where
inhibition of thrombin is required. The compounds of the

invention are expected to be useful in animals, including man, in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disease states in which the compounds have a potential utility are in treatment or prophylaxis of thrombosis and hypercoagulability in blood and tissues. Disease states in which the compounds have a potential utility, in treatment and/or prophylaxis, include venous thrombosis and pulmonary embolism, arterial thrombosis, such as in myocardial ischemia, myocardial infarction, unstable angina, thrombosis-based stroke and peripheral arterial thrombosis. Further, the compounds have expected utility in prophylaxis of atherosclerotic diseases such as coronary arterial disease, cerebral arterial disease and peripheral arterial disease. Further, the compounds are expected to be useful together with thrombolytics in myocardial infarction. Further, the compounds have expected utility in the treatment or prophylaxis for reocclusion after thrombolysis, percutaneous transluminal angioplasty (PTCA) and coronary bypass operations. Further, the compounds have expected utility in prevention of rethrombosis after microsurgery. Further, the compounds are expected to be useful in anticoagulant treatment in connection with artificial organs and cardiac valves. Further, the compounds have expected utility in anticoagulant treatment in hemodialysis and disseminated intravascular coagulation. A further expected utility is in rinsing of catheters and mechanical devices used in patients *in vivo*, and as an anticoagulant for preservation of blood, plasma and other blood products *in vitro*. Still further, the compounds have expected utility in other diseases where blood coagulation could be a fundamental contributing process or a source of secondary pathology, such as cancer, including metastasis, and inflammatory diseases, including arthritis and diabetes. The anti-coagulant compound is administered orally, or parenterally, e.g., by intravenous infusion (iv), intramuscular injection (im) or subcutaneous injection (sc).

The specific dose of a compound administered according to this invention to obtain therapeutic and/or prophylactic effects will, of course, be determined by the particular circumstances surrounding the case, including, for example, the compound administered, the rate of administration, and the condition being treated.

A typical daily dose for each of the above utilities is between about 0.01 mg/kg and about 1000 mg/kg. The dose regime may vary, e.g., for prophylactic use a single daily dose may be administered or multiple doses such as 3 or 5 times daily may be appropriate. In critical care situations a compound of the invention is administered by iv infusion at a rate between about 0.01 mg/kg/h and about 20 mg/kg/h and preferably between about 0.1 mg/kg/h and about 5 mg/kg/h.

The method of this invention also is practiced in conjunction with a clot lysing agent, e.g., tissue plasminogen activator (t-PA), modified t-PA, streptokinase or urokinase. In cases when clot formation has occurred and an artery or vein is blocked, either partially or totally, a clot lysing agent is usually employed. A compound of the invention can be administered prior to or along with the lysing agent or subsequent to its use alone and preferably further is administered along with aspirin to prevent the reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with a platelet glycoprotein receptor (IIb/IIIa) antagonist, that inhibits platelet aggregation. A compound of the invention can be administered prior to or along with the IIb/IIIa antagonist or subsequent to its use to prevent the reoccurrence of clot formation.

The method of this invention is also practiced in conjunction with aspirin. A compound of the invention can be administered prior to or along with aspirin or subsequent to its use to prevent the reoccurrence of clot formation. As stated above, preferably a compound of the present invention

is administered in conjunction with a clot lysing agent and aspirin.

This invention also provides pharmaceutical formulations for use in the above described therapeutic method. Pharmaceutical formulations of the invention comprise an effective thrombin inhibiting amount of a compound of Formula I in association with a pharmaceutically acceptable carrier, excipient or diluent. For oral administration the antithrombotic compound is formulated in gelatin capsules or tablets which may contain excipients such as binders, lubricants, disintegration agents and the like. For parenteral administration the antithrombotic is formulated in a pharmaceutically acceptable diluent, e.g., physiological saline (0.9%), 5% dextrose, Ringer's solution, and the like.

The compound of the present invention can be formulated in unit dosage formulations comprising a dose between about 0.1 mg and about 1000 mg. Preferably the compound is in the form of a pharmaceutically acceptable salt such as for example the sulfate salt, acetate salt or a phosphate salt. An example of a unit dosage formulation comprises 5 mg of a compound of the present invention as a pharmaceutically acceptable salt in a 10 mL sterile glass ampoule. Another example of a unit dosage formulation comprises about 10 mg of a compound of the present invention as a pharmaceutically acceptable salt in 20 mL of isotonic saline contained in a sterile ampoule.

The compounds can be administered by a variety of routes including oral, rectal, transdermal, subcutaneous, intravenous, intramuscular, and intranasal. The compounds of the present invention are preferably formulated prior to administration. Therefore, another embodiment of the present invention is a pharmaceutical formulation comprising an effective amount of a compound of Formula I or a pharmaceutically acceptable salt or solvate thereof in association with a pharmaceutically acceptable carrier, diluent or excipient therefor.

The active ingredient in such formulations comprises from 0.1% to 99.9% by weight of the formulation. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other
5 ingredients of the formulation and not deleterious to the recipient thereof.

The present pharmaceutical formulations are prepared by known procedures using well known and readily available ingredients. In making the compositions of the
10 present invention, the active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material
15 which acts as a vehicle, excipient or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols, (as a solid or in a liquid medium), soft and hard gelatin capsules,
20 suppositories, sterile injectable solutions, sterile packaged powders, and the like. The compositions of this invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

25 The following formulation examples are illustrative only and are not intended to limit the scope of the invention in any way. "Active ingredient," of course, means a compound according to Formula I or a pharmaceutically acceptable salt or solvate thereof.

Formulation 1

Hard gelatin capsules are prepared using the following ingredients:

5

	Quantity <u>(mg/capsule)</u>
Active ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

Formulation 2

A tablet is prepared using the ingredients below:

10

	Quantity <u>(mg/capsule)</u>
Active ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

The components are blended and compressed to form tablets each weighing 665 mg.

15

Formulation 3

An aerosol solution is prepared containing the following components:

	<u>Weight</u>
Active ingredient	0.25
Ethanol	25.75
Propellant 22 (Chlorodifluoromethane)	<u>70.00</u>
Total	100.00

The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30 °C and transferred to a filling device. The required amount is then fed to a stainless steel container and diluted with the remainder of the propellant. The valve units are then fitted to the container.

Formulation 4

10

Tablets, each containing 60 mg of active ingredient, are made as follows:

Active ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

15

The active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50 °C and passed through a No. 18 mesh U.S. Sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each weighing 150 mg.

25

Formulation 5

Capsules, each containing 80 mg of active ingredient, are made as follows:

5

Active ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg

The active ingredient, cellulose, starch, and magnesium stearate are blended, passed through a No. 45 mesh U.S. sieve, and filled into hard gelatin capsules in 200 mg quantities.

10

Formulation 6

Suppositories, each containing 225 mg of active ingredient, are made as follows:

15

Active ingredient	225 mg
Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

The active ingredient is passed through a No. 60 mesh U.S. sieve and suspended in the saturated fatty acid glycerides previously melted using the minimum heat necessary. The mixture is then poured into a suppository mold of nominal 2 g capacity and allowed to cool.

20

Formulation 7

Suspensions, each containing 50 mg of active ingredient per 5 mL dose, are made as follows:

5

Active ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 mL
Benzoic acid solution	0.10 mL
Flavor	q.v.
Color	q.v.
Purified water to total	5 mL

The active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

10

Formulation 8

15

An intravenous formulation may be prepared as follows:

Active ingredient	100 mg
Isotonic saline	1,000 mL

20

The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 mL per minute.

25

The compounds provided by the invention (Formula I) are orally active and selectively inhibit the action of thrombin in mammals.


The ability of the compounds of the present invention to be an effective and orally active thrombin

inhibitor is evaluated in one or more of the following assays.

The inhibition of thrombin is demonstrated by in vitro inhibition of the amidase activity of thrombin as measured in an assay in which thrombin hydrolyzes the chromogenic substrate, N-benzoyl-L-phenylalanyl-L-valyl-L-arginyl-p-nitroanilide.

The assay is carried out by mixing 50 μ L buffer (0.03M Tris, 0.15M NaCl, pH 7.4) with 25 μ L of bovine thrombin or human thrombin solution (0.21 mg/mL of thrombostat bovine thrombin, Parke-Davis, or purified human thrombin, Enzyme Research Laboratories, South Bend, Indiana, at about 8 NIH units/mL, in the same buffer) and 25 μ L of test compound in a solvent (in 50% aqueous methanol, v:v). The 150 μ L of an aqueous solution of the chromogenic substrate (at 0.25 mg/mL) are added and the rates of hydrolysis of the substrate are measured by monitoring the reactions at 405 nm for the release of p-nitroaniline. Standard curves are constructed by plotting free thrombin concentration against hydrolysis rate. The hydrolysis rates observed with test compounds are then converted to "free thrombin" values in the respective assays by use of the standard curves. The bound thrombin (bound to test compound) is calculated by subtracting the amount of free thrombin observed in each assay from the known initial amount of thrombin used in the assay. The amount of free inhibitor in each assay is calculated by subtracting the number of moles of bound thrombin from the number of moles of added inhibitor (test compound).

The Kass value is the hypothetical equilibrium constant for the reaction between thrombin and the test compound (I).



$$\text{Kass} = \frac{[\text{Thrombin-I}]}{[(\text{Thrombin}) \times (\text{I})]}$$

Kass is calculated for a range of concentrations of test compounds and the mean value is reported in units of liter per mole.

5 By substantially following the procedures described above for human thrombin, and using other human blood coagulation system serine proteases and proteases of the fibrinolytic system with the appropriate chromogenic substrates, identified below, selectivity of the compounds of
10 the present invention with respect to the coagulation factor serine proteases and with respect to the fibrinolytic system serine proteases are evaluated as well as their substantial lack of interference with serine proteases of the fibrinolytic system. Thrombin inhibitors preferably should
15 spare fibrinolysis induced by urokinase, tissue plasminogen activator (t-PA) and streptokinase. This would be important to the therapeutic use of such agents as an adjunct to streptokinase, t-PA or urokinase thrombolytic therapy and to the use of such agents as an endogenous fibrinolysis-sparing
20 (with respect to t-PA and urokinase) antithrombotic agent. In addition to the lack of interference with the amidase activity of the fibrinolytic proteases, such fibrinolytic system sparing can be studied by the use of human plasma clots and their lysis by the respective fibrinolytic
25 plasminogen activators.

Human factors X, Xa, IXa, XIa, and XIIa are purchased from Enzyme Research Laboratories, South Bend, Indiana; human urokinase from Leo Pharmaceuticals, Denmark; and recombinant activated Protein C (aPC) is prepared at Eli
30 Lilly and Co. substantially according to U.S. Patent 4,981,952. Chromogenic substrates: N-Benzoyl-Ile-Glu-Gly-Arg-p-nitroanilide (for factor Xa); N-Cbz-D-Arg-Gly-Arg-p-nitroanilide (for factor IXa assay as the factor Xa substrate); Pyroglutamyl-Pro-Arg-p-nitroanilide (for Factor
35 XIa and for aPC); H-D-Pro-Phe-Arg-p-nitroanilide (for factor XIIa); and Pyroglutamyl-Gly-Arg-p-nitroanilide (for urokinase); are purchased from KabiVitrum, Stockholm, Sweden,

or from Midwest Biotech, Fishers, Indiana. Bovine trypsin is purchased from Worthington Biochemicals, Freehold, New Jersey, and human plasma kallikrein from Kabi Vitrum, Stockholm, Sweden. Chromogenic substrate H-D-Pro-Phe-Arg-p-nitroanilide for plasma kallikrein is purchased from Kabi Vitrum, Stockholm, Sweden. N-Benzoyl-Phe-Val-Arg-p-nitroanilide, the substrate for human thrombin and for trypsin, is synthesized according to procedures described above for the compounds of the present invention, using known methods of peptide coupling from commercially available reactants or purchased from Midwest Biotech, Fishers, Indiana.

Human plasmin is purchased from Boehringer Mannheim, Indianapolis, Indiana; nt-PA is purchased as single chain activity reference from American Diagnostica, Greenwich, Connecticut; modified-t-PA6 (mt-PA6) is prepared at Eli Lilly and Company by procedure known in the art (See, Burck, et al., J. Biol. Chem., 265, 5120-5177 (1990)). Plasmin chromogenic substrate H-D-Val-Leu-Lys-p-nitroanilide and tissue plasminogen activator (t-PA) substrate H-D-Ile-Pro-Arg-p-nitroanilide are purchased from Kabi Vitrum, Stockholm, Sweden.

In the chromogenic substrates described above the three-letter symbols Ile, Glu, Gly, Pro, Arg, Phe, Val, Leu and Lys are used to indicate the corresponding amino acid group isoleucine, glutamic acid, glycine, proline, arginine, phenylalanine, valine, leucine and lysine, respectively.

Table 1 which follows lists the Kass values obtained with the indicated compound represented by the Formula I.

Table 1
Inhibition Properties

-----Kass X 10 ⁶ (L/mole)-----						
5	<u>Example</u>	Human <u>Thrombin</u>	<u>Xa</u>	<u>Trypsin</u>	<u>Plasmin</u>	<u>t-PA</u>
	1	3.3	0.0017	0.38	0.010	0.082
	2	4.4	0.0092	1.0	0.045	0.023
10	3	2.1	0.0021	0.24	0.0046	0.033
	4	13.	0.0057	0.076	0.0033	0.074
	5	0.035	0.00039	0.00015	0.00022	<0.001
	6	2.3	0.00024	0.0016	0.00023	<0.001
	7	1.9	0.0041	0.0081	0.00022	0.0037
15	8	5.3	0.00038	0.022	0.00080	0.0012
	9	0.033	0.00058	0.00017	<0.001	<0.001
	10	1.4	0.000024	0.077	0.0045	0.00055
	11	0.064	0.000068	0.0036	0.0060	<0.001
	12	0.44	0.000042	0.0022	0.00087	0.000042
20	13	0.49	0.00098	0.00066	0.00011	<0.001
	14	0.021	0.00014	0.000056	0.000046	<0.001
	15	220	0.0070	2.6	0.017	0.004
	16	0.97	0.0051	0.02	0.0017	0.0012
	17	33	0.049	3.8	0.024	0.0034
25	18	180	0.21	14	0.085	0.0094
	19	11	0.14	41	0.16	0.0056
	20	10	0.10	19	0.10	0.0077
	21	3.6	0.62	0.96	0.031	0.002
	22	3.0	0.043	8.5	0.042	0.004
30	23	770	0.14	21	0.44	0.31
	24	0.78	<0.001	0.03	<0.001	<0.001
	25	0.43	<0.001	0.01	<0.001	0.01
	26	13.	0.01	0.21	0.01	0.01
	27	0.29	0.02	0.02	<0.001	0.02
35	28	0.95	0.01	0.11	<0.001	<0.001
	29	0.76	0.02	0.02	<0.001	<0.001
	30	0.55	0.02	0.03	<0.001	<0.001

199

	31	0.07	0.14	0.05	<0.001	<0.001
	32	0.13	0.03	0.04	<0.001	0.01
	33	0.04	0.04	0.02	<0.001	<0.001
	34	0.65	<0.001	0.49	0.01	0.01
5	35	0.09	<0.001	0.04	<0.001	<0.001
	36	0.06	<0.001	0.01	<0.001	<0.001
	37	0.02		<0.009		
	38	1.1	<0.001	0.07	<0.001	
	39	0.12	<0.001	<0.02	<0.001	
10	40	0.03	<0.001	0.01	<0.001	
	41	0.01	<0.001	<0.001	<0.001	
	42	38.	0.07	43.	0.12	0.01
	43	1.4	0.05	100.	0.30	0.01
	44	630	0.66	2,600.	1.8	0.04
15	45	610	1.5	110.	1.1	0.77
	46	240	0.07	48.	0.13	0.01
	47	45	0.04	82.	0.18	0.01
	48	4,100	5.5	250.	0.43	0.05
	49	400	3.4	940.	0.84	0.04
20	50	40	4.3	2,200.	1.8	0.04
	51	1,300	0.84	27.	0.13	0.01
	52	700	0.46	47.	0.11	0.01
	53	0.36	<0.001	0.1	<0.001	<0.001
	54	2.8	<0.001	0.52	0.01	<0.001
25	55	0.01	<0.001	0.01	<0.001	<0.001
	56	490	0.20	3.6	0.01	<0.001
	57	39	0.01	0.24	0.01	<0.001
	58	0.60	<0.001	0.40	<0.001	<0.001
	59	1.10	<0.001	0.21	<0.001	<0.001
30	60	0.24	<0.001	0.03	<0.001	<0.001
	61	63	0.03	1.0	0.004	0.008
	62	7.6				
	63	0.31				
	64	15	0.01			
35	65	4,500		570.		
	66	120		4.6		
	67	11				

			200	
	68	600		
	69	340	3.7	
	70	430	9.8	
	71	1,400	240.	
5	72	2,300	280	
	73	28		
	74	29		
	75	0.45		
	76	12	0.36	
10	77	11		
	78	11	8.6	
	79	2.0	0.24	
	80	1,700.	90.	
	81	5.3	2.1	
15	82	5.0		
	83	2.4	0.03	
	84	2.9	0.07	
	85	265	30.8	
	86	200	0.63	
20	87	30	2.5	
	88	410	3.9	
	89	0.14	0.071	
	90	0.61	0.002	
	91	3.6	0.007	
25	92	130	7.7	

It is noted that, unexpectedly, compounds in which X contains a D-cyclohexylalanyl moiety have improved potency with respect to inhibition of thrombin and display especially surprisingly enhanced potency with respect to the inhibition of factor Xa when compared with, for example, the corresponding compounds in which X contains a D-phenylalanyl moiety.

Materials

Dog plasma is obtained from conscious mixed-breed hounds (either sex Hazelton-LRE, Kalamazoo, Michigan, U.S.A.) by venipuncture into 3.8 percent citrate. Fibrinogen is prepared from fresh dog plasma and human fibrinogen is prepared from in-date ACD human blood at the fraction I-2 according to previous procedures and specifications. Smith, Biochem. J., 185, 1-11 (1980); and Smith, et al., Biochemistry, 11, 2958-2967, (1972). Human fibrinogen (98 percent pure/plasmin free) is from American Diagnostica, Greenwich, Connecticut. Radiolabeling of fibrinogen I-2 preparations is performed as previously reported. Smith, et al., Biochemistry, 11, 2958-2967, (1972). Urokinase is purchased from Leo Pharmaceuticals, Denmark, as 2200 Ploug units/vial. Streptokinase is purchased from Hoechst-Roussel Pharmaceuticals, Somerville, New Jersey.

Methods - Effects on Lysis of Human Plasma Clots by t-PA

Human plasma clots are formed in micro test tubes by adding 50 μ L thrombin (73 NIH unit/mL) to 100 μ L human plasma which contains 0.0229 μ Ci 125-iodine labeled fibrinogen. Clot lysis is studied by overlaying the clots with 50 μ L of urokinase or streptokinase (50, 100, or 1000 unit/mL) and incubating for 20 hours at room temperature. After incubation the tubes are centrifuged in a Beckman Microfuge. 25 μ L of supernate is added into 1.0 mL volume of 0.03 M tris/0.15 M NaCl buffer for gamma counting. Counting controls 100 percent lysis are obtained by omitting thrombin (and substituting buffer). The thrombin inhibitors are evaluated for possible interference with fibrinolysis by including the compounds in the overlay solutions at 1, 5, and 10 ug/mL concentrations. Rough approximations of IC₅₀ values are estimated by linear extrapolations from data points to a value which would represent 50 percent of lysis for that particular concentration of fibrinolytic agent.

Anticoagulant ActivityMaterials

Dog plasma and rat plasma is obtained from conscious mixed-breed hounds (either sex, Hazelton-LRE, Kalamazoo, Michigan, U.S.A.) or from anesthetized male Sprague-Dawley rats (Harlan Sprague-Dawley, Inc., Indianapolis, Indiana, U.S.A.) by venipuncture into 3.8 percent citrate. Fibrinogen is prepared from in-date ACD human blood as the fraction I-2 according to previous procedures and specifications. Smith, Biochem. J., **185**, 1-11 (1980); and Smith, et al., Biochemistry, **11**, 2958-2967 (1972). Human fibrinogen is also purchased as 98 percent pure/plasmin free from American Diagnostica, Greenwich, Connecticut. Coagulation reagents ACTIN, Thromboplastin, and Human plasma are from Baxter Healthcare Corp., Dade Division, Miami, Florida. Bovine thrombin from Parke-Davis (Ann Detroit, Michigan) is used for coagulation assays in plasma.

Methods20 Anticoagulation Determinations

Coagulation assay procedures are as previously described. Smith, et al., Thrombosis Research, **50**, 163-174 (1988). A CoAScreener coagulation instrument (American LABor, Inc.) is used for all coagulation assay measurements. The prothrombin time (PT) is measured by adding 0.05 mL saline and 0.05 mL Thromboplastin-C reagent to 0.05 mL test plasma. The activated partial thromboplastin time (APTT) is measured by incubation of 0.05 mL test plasma with 0.05 mL Actin reagent for 120 seconds followed by 0.05 mL CaCl₂ (0.02 M). The thrombin time (TT) is measured by adding 0.05 mL saline and 0.05 mL thrombin (10 NIH units/mL) to 0.05 mL test plasma. The compounds of Formula I are added to human or animal plasma over a wide range of concentrations to determine prolongation effects on the APTT, PT, and TT assays. Linear extrapolations are performed to estimate the concentrations required to double the clotting time for each assay.

Animals

Male Sprague Dawley rats (350-425 gm, Harlan Sprague Dawley Inc., Indianapolis, IN) are anesthetized with xylazine (20 mg/kg, s.c.) and ketamine (120 mg/kg, s.c.) and maintained on a heated water blanket (37 °C). The jugular vein(s) is cannulated to allow for infusions.

Arterio-Venous shunt model

The left jugular vein and right carotid artery are cannulated with 20 cm lengths of polyethylene PE 60 tubing. A 6 cm center section of larger tubing (PE 190) with a cotton thread (5 cm) in the lumen, is friction fitted between the longer sections to complete the arterio-venous shunt circuit. Blood circulated through the shunt for 15 minutes before the thread is carefully removed and weighed. The weight of a wet thread is subtracted from the total weight of the thread and thrombus (see J.R. Smith, Br. J. Pharmacol., 77,29 (1982)).

When the compound of Example 48 was compared with D-MePhe-Pro-Arg-H (discussed above at page 2) in the Arterio-Venous shunt model, the antithrombotic potency was found to be 9-fold greater during continuous intravenous infusion. To reduce thrombus weight to the same extent (to approximately 20% of control), the compound of Example 48 prolonged plasma thrombin time about 3-fold, whereas plasma thrombin time was prolonged greater than 20-fold during infusion of the standard compound. Prothrombin time and APTT were extended only to about 120% of control (a few seconds) during infusion of the compound of Example 48.

FeCl₃ model of arterial injury

The carotid arteries are isolated via a midline ventral cervical incision. A thermocouple is placed under each artery and vessel temperature is recorded continuously on a strip chart recorder. A cuff of tubing (0.058 ID x 0.077 OD x 4 mm, Baxter Med. Grade Silicone), cut longitudinally, is placed around each carotid directly above the thermocouple. FeCl₃ hexahydrate is dissolved in water and the concentration

(20%) is expressed in terms of the actual weight of FeCl_3 only. To injure the artery and induce thrombosis, 2.85 μl is pipetted into the cuff to bathe the artery above the thermocouple probe. Arterial occlusion is indicated by a rapid drop in temperature. The time to occlusion is reported in minutes and represented the elapsed time between application of FeCl_3 and the rapid drop in vessel temperature (see K.D. Kurz, Thromb. Res., 60, 269 (1990)).

10 Spontaneous thrombolysis model

In vitro data suggested that the peptide thrombin inhibitors inhibit thrombin and other serine proteases, such as plasmin and tissue plasminogen activator. To assess if the compounds inhibited fibrinolysis in vivo, the rate of spontaneous thrombolysis is determined by implanting a labeled whole blood clot into the pulmonary circulation. Rat blood (1 mL) is mixed rapidly with bovine thrombin (4 IU, Parke Davis) and ^{125}I human fibrogen (5 μCi , ICN), immediately drawn into silastic tubing and incubated at 37 °C for 1 hour. The aged thrombus is expelled from the tubing, cut into 1 cm segments, washed 3X in normal saline and each segment is counted in a gamma counter. A segment with known counts is aspirated into a catheter that is subsequently implanted into the jugular vein. The catheter tip is advanced to the vicinity of the right atrium and the clot is expelled to float into the pulmonary circulation. One hour after implant, the heart and lungs are harvested and counted separately. Thrombolysis is expressed as a percentage where:

30
$$\% \text{Thrombolysis} = \frac{(\text{injected cpm} - \text{lung cpm})}{\text{injected cpm}} \times 100$$

The fibrinolytic dissolution of the implanted clot occurs time-dependently (see J.P. Clozel, Cardiovas. Pharmacol., 12, 520 (1988)).

Coagulation parameters

Plasma thrombin time (TT) and activated partial thromboplastin time (APTT) are measured with a fibrometer. Blood is sampled from a jugular catheter and collected in syringe containing sodium citrate (3.8%, 1 part to 9 parts blood). To measure TT, rat plasma (0.1 mL) is mixed with saline (0.1 mL) and bovine thrombin (0.1 mL, 30 U/mL in TRIS buffer; Parke Davis) at 37 °C. For APTT, plasma (0.1 mL) and APTT solution (0.1 mL, Organon Teknika) are incubated for 5 minutes (37 °C) and CaCl₂ (0.01 mL, 0.025 M) is added to start coagulation. Assays are done in duplicate and averaged.

Index of Bioavailability

A measure of bioactivity, plasma thrombin time (TT), served as a substitute for the assay of parent compound on the assumption that increments in TT resulted from thrombin inhibition by parent only. The time course of the effect of the thrombin inhibitor upon TT is determined after i.v. bolus administration to anesthetized rats and after oral treatment of fasted conscious rats. Due to limitations of blood volume and the number of points required to determine the time course from time of treatment to the time when the response returned to pretreatment values, two populations of rats are used. Each sample population represented alternating sequential time points. The average TT over the time course is used to calculate area under the curve (AUC). The index of bioavailability is calculated by the formula shown below and is expressed as percent relative activity.

The area under the curve (AUC) of the plasma TT time course is determined and adjusted for the dose. This index of bioavailability is termed "% Relative Activity" and is calculated as

$$\% \text{Relative Activity} = \frac{\text{AUC po}}{\text{AUC iv}} \times \frac{\text{Dose iv}}{\text{Dose po}} \times 100$$

Compounds

Compound solutions are prepared fresh daily in normal saline and are injected as a bolus or are infused starting 15 minutes before and continuing throughout the experimental perturbation which is 15 minutes in the arteriovenous shunt model and 60 minutes in the FeCl₃ model of arterial injury and in the spontaneous thrombosis model. Bolus injection volume is 1 mL/kg for i.v., and 5 mL/kg for p.o. and infusion volume is 3 mL/hr.

10

Statistics

Results are expressed as means +/- SEM. One-way analysis of variance is used to detect statistically significant differences and then Dunnett's test is applied to determine which means are different. Significance level for rejection of the null hypothesis of equal means is $P < 0.05$.

Animals

Male dogs (Beagles; 18 months - 2 years; 12-13 kg, Marshall Farms, North Rose, New York 14516) are fasted overnight and fed Purina certified Prescription Diet (Purina Mills, St. Louis, Missouri) 240 minutes after dosing. Water is available ad libitum. The room temperature is maintained between 66-74°F; 45-50% relative humidity; and lighted from 0600-1800 hours.

Pharmacokinetic model.

Test compound is formulated immediately prior to dosing by dissolving in sterile 0.9% saline to a 5 mg/mL preparation. Dogs are given a single 2 mg/kg dose of test compound by oral gavage. Blood samples (4.5 mL) are taken from the cephalic vein at 0.25, 0.5, 0.75, 1, 2, 3, 4 and 6 hours after dosing. Samples are collected in citrated Vacutainer tubes and kept on ice prior to reduction to plasma by centrifugation. Plasma samples are derivatized with dinitrophenylhydrazine and analyzed by HPLC (Zorbax SB-C8 column) eluting with methanol/500 mM sodium acetate adjusted to pH 7 with

phosphoric acid (60:40, v/v). Plasma concentration of test compound is recorded and used to calculate the pharmacokinetic parameters: elimination rate constant, K_e ; total clearance, Cl_t ; volume of distribution, V_D ; time of maximum plasma test compound concentration, T_{max} ; maximum concentration of test compound at T_{max} , C_{max} ; plasma half-life, $t_{0.5}$; area under the curve, A.U.C.; and fraction of test compound absorbed, F .

10 Canine Model of Coronary Artery Thrombosis

Surgical preparation and instrumentation of the dogs are as described in Jackson, et al., Circulation, 82, 930-940 (1990). Mixed-breed hounds (aged 6-7 months, either sex, Hazelton-LRE, Kalamazoo, MI, U.S.A.) are anesthetized with sodium pentobarbital (30 mg/kg intravenously, i.v.), intubated, and ventilated with room air. Tidal volume and respiratory rates are adjusted to maintain blood PO_2 , PCO_2 , and pH within normal limits. Subdermal needle electrodes are inserted for the recording of a lead II ECG.

20 The left jugular vein and common carotid artery are isolated through a left mediolateral neck incision. Arterial blood pressure (ABP) is measured continuously with a precalibrated Millar transducer (model MPC-500, Millar Instruments, Houston, TX, U.S.A.) inserted into the carotid artery. The jugular vein is cannulated for blood sampling during the experiment. In addition, the femoral veins of both hindlegs are cannulated for administration of test compound.

30 A left thoracotomy is performed at the fifth intercostal space, and the heart is suspended in a pericardial cradle. A 1- to 2-cm segment of the left circumflex coronary artery (LCX) is isolated proximal to the first major diagonal ventricular branch. A 26-gauge needle-tipped wire anodal electrode (Teflon®-coated, 30-gauge silverplated copper wire) 35 3-4 mm long is inserted into the LCX and placed in contact with the intimal surface of the artery (confirmed at the end

of the experiment). The stimulating circuit is completed by placing the cathode in a subcutaneous (s.c.) site. An adjustable plastic occluder is placed around the LCX, over the region of the electrode. A precalibrated electromagnetic flow probe (Carolina Medical Electronics, King, NC, U.S.A.) is placed around the LCX proximal to the anode for measurement of coronary blood flow (CBF). The occluder is adjusted to produce a 40-50% inhibition of the hyperemic blood flow response observed after 10-s mechanical occlusion of the LCX. All hemodynamic and ECG measurements are recorded and analyzed with a data acquisition system (model M3000, Modular Instruments, Malvern, PA, U.S.A.).

Thrombus Formation and Compound Administration Regimens

Electrolytic injury of the intima of the LCX is produced by applying 100- μ A direct current (DC) to the anode. The current is maintained for 60 minutes and then discontinued whether the vessel has occluded or not. Thrombus formation proceeds spontaneously until the LCX is totally occluded (determined as zero CBF and an increase in the S-T segment). Compound administration is started after the occluding thrombus is allowed to age for 1 hour. A 2-hour infusion of the compounds of the present invention at doses of 0.5 and 1 mg/kg/h is begun simultaneously with an infusion of thrombolytic agent (e.g. tissue plasminogen activator, streptokinase, APSAC). Reperfusion is followed for 3 hours after administration of test compound. Reocclusion of coronary arteries after successful thrombolysis is defined as zero CBF which persisted for ≥ 30 minutes.

Hematology and template bleeding time determinations

Whole blood cell counts, hemoglobin, and hematocrit values are determined on a 40- μ l sample of citrated (3.8%) blood (1 part citrate:9 parts blood) with a hematology analyzer (Cell-Dyn 900, Sequoia-Turner, Mount View, CA, U.S.A.). Gingival template bleeding times are determined with a Simplate II bleeding time device (Organon Teknika Durham, N.C., U.S.A.).

The device is used to make 2 horizontal incisions in the gingiva of either the upper or lower left jaw of the dog. Each incision is 3 mm wide x 2 mm deep. The incisions are made, and a stopwatch is used to determine how long bleeding occurs. A cotton swab is used to soak up the blood as it oozes from the incision. Template bleeding time is the time from incision to stoppage of bleeding. Bleeding times are taken just before administration of test compound (0 minutes), 60 minutes into infusion, at conclusion of administration of the test compound (120 minutes), and at the end of the experiment.

All data are analyzed by one-way analysis of variance (ANOVA) followed by Student-Neuman-Kuels post hoc *t* test to determine the level of significance. Repeated-measures ANOVA are used to determine significant differences between time points during the experiments. Values are determined to be statistically different at least at the level of $p < 0.05$. All values are mean \pm SEM. All studies are conducted in accordance with the guiding principles of the American Physiological Society. Further details regarding the procedures are described in Jackson, et al., J. Cardiovasc. Pharmacol., 21, 587-599 (1993).

25

Table 2

Human Plasma Anticoagulation					
	<u>2x Clotting time (ng/mL)</u>			% oral/i.v.	
	<u>Example</u>	<u>TT</u>	<u>APTT</u>	<u>PT</u>	<u>activity (Rat)</u>
30	1	250	NT	NT	8
	2	170	NT	NT	NT
	3	590	NT	NT	NT
	4	230	NT	NT	NT
35	5	30,000	NT	NT	NT
	6	390	NT	NT	NT
	7	490	NT	NT	NT

210

	8	130	NT	NT	NT
	9	>91,000	NT	NT	NT
	10	420	NT	NT	NT
	11	8,800	NT	NT	NT
5	12	1,700	NT	NT	NT
	13	660	NT	NT	NT
	14	21,000	NT	NT	NT
	15	9	89	200	4
	16	650	8,600	8,500	NT
10	17	26	370	400	15
	18	7	79	170	12
	19	58	300	540	NT
	20	62	550	600	23
	21	160	2,000	1,000	NT
15	22	170	2,300	1,000	NT
	23	7	86	120	12
	24	1,000	25,900	25,000	NT
	25	1,430	34,600	40,400	NT
	26	37	870	750	NC ¹
20	27	1,940	27,800	27,000	NT
	28	520	8,100	8,300	NT
	29	750	9,800	15,800	NT
	30	1,000	14,000	14,400	NT
	31	10,800	34,500	30,000	NT
25	32	3,100	28,500	47,800	NT
	33	18,900	59,300	70,900	NT
	34	530	4,900	5,500	NT
	35	4,500	50,400	78,400	NT
	36	7,700	>91,000	>91,000	NT
30	37	>9,000	>9,000	>9,000	NT
	38	540	6,100	11,100	NT
	39	5,300	69,200	78,600	NT
	40	35,100	>91,000	>91,000	NT
	41	82,200	>91,000	>91,000	NT
35	42	20	270	320	NC ²
	43	280	1,100	930	NT
	44	4	100	170	NT

211

	45	8	200	330	NC ²
	46	2.	67	77	NC ³
	47	12.	140	270	NC ²
	48	2.	33	59	22
5	49	5	130	130	NC ²
	50	35	460	420	NT
	51	2	48	110	NT
	52	6	80	170	NT
	53	1,400	33,800	34,500	NT
10	54	140	3,300	2,200	NT
	55	55,800	>91,000	>91,000	NT
	56	5	160	200	NT
	57	14	360	340	NT
	58	710	14,300	11,400	NT
15	59	420	5,500	7,000	NT
	60	3,200	22,200	77,700	NT
	61	45	679	756	NT
	62	54			NT
	63	680			NT
20	64	23	900	650	NC ³
	65	1	48	85	NT
	66	7.9	180	270	NT
	67	34.	1,800	1,300	NT
	68	4.	49	190	NT
25	69	5.	110	220	NT
	70	2	160	180	NT
	71	1	89	150	NC ¹
	72	1	160	160	NC ¹
	73	21	340	330	NT
30	74	20	420	350	NT
	75				NT
	76	29	490	560	NT
	77	53	1,600	890	NT
	78	46	430	760	NT
35	79	140	1,700	2,400	NT
	80	4	40	130	NT

212

	81	110	1,500	2,700	NT
	82	130	3,800	3,000	NT
	83	91	1,000	1,300	NT
	84	110	1,500	1,900	NT
5	85	4.9	100	197	NT
	86	4	81	180	NT
	87	9	260	310	NT
	88	4	100	200	NT
	89	570	51,000	37,000	NT
10	90	330	14,500	14,600	NT
	91	63	3,000	3,500	NT
	92	8.9	210	340	NT

15 Notes to Table 2

NC¹ indicates screening not completed; low or very low activity observed at 20 mg/kg p.o.

20 NC² indicates screening not completed; low or very low activity observed at 60 mg/kg p.o.

NC³ indicates screening not completed; low or very low activity observed at 50 mg/kg p.o.

25

NT indicates not tested, as does a blank entry.


We claim:

1. A compound having the Formula I

5 $X-Y-NH-(CH_2)_r-G$ I

wherein

X is prolinyl, homoprolinyl,
 $R^m-(CH_2)_g-NH-CH_2-C(O)-$,



10


in which

R^d is carboxy or methylsulfonyl;

R^e is NHR^C , $NHCOR^C$ or $NHCOOR^C$; in which

15 R^C is C_1-C_{10} alkyl, C_3-C_8 cycloalkyl or a (C_3-C_8) cycloalkyl (C_1-C_6) alkyl radical of 4-10 carbons;

T is C_3-C_8 cycloalkyl, C_1-C_8 alkyl,



20



a is 0, 1 or 2; and

Q is $-OH$, C_1-C_4 alkoxy, or $-NH-A$;

A is hydrogen, C_1-C_4 alkyl, $R''SO_2-$, $R''OC(O)-$,




$R''C(O)-$, $R''NC(O)-$ or $-(CH_2)_g-R^m$;

- g is 1, 2, or 3;
 B is hydrogen or C₁-C₄ alkyl;
 R' is hydrogen or C₁-C₄ alkyl;
 R" is C₁-C₄ alkyl, C₁-C₄ perfluoroalkyl, -(CH₂)_d-R^m,
 5 or unsubstituted or substituted aryl, where aryl is phenyl,
 naphthyl, a 5- or 6-membered unsubstituted or substituted
 aromatic heterocyclic ring, having one or two heteroatoms
 which are the same or different and which are selected from
 sulfur, oxygen and nitrogen, or a 9- or 10-membered
 10 unsubstituted or substituted fused bicyclic aromatic
 heterocyclic group having one or two heteroatoms which are
 the same or different and which are selected from sulfur,
 oxygen and nitrogen;
 R^m is -COOR^b, -SO₂(C₁-C₄ alkyl), -SO₃H, -P(O)(OR^b)₂
 15 or tetrazol-5-yl;
 Rⁿ is -COOR^b or tetrazol-5-yl;
 each R^b is independently hydrogen or C₁-C₄ alkyl;
 d is 1, 2, or 3;
 m is 0, 1, or 2;
 20 n is 0, 1, or 2; and
 Z is hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy,
 halo or R_aSO₂NH-, where R_a is C₁-C₄ alkyl;




- 25 in which
 R^g is C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or -(CH₂)_p-L-
 (CH₂)_q-T';
 R^p is hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or
 -(CH₂)_p-L-(CH₂)_q-T';

- where p is 0, 1, 2, 3, or 4; L is a bond, -O-, -S-, or -NH-; q is 0, 1, 2 or 3; and T' is hydrogen, C₁-C₄ alkyl, C₃-C₈ cycloalkyl, -COOH, -CONH₂, or Ar, where Ar is unsubstituted or substituted aryl, where aryl is phenyl, naphthyl, a 5- or 6-membered unsubstituted or substituted aromatic heterocyclic ring, having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen, or a 9- or 10-membered unsubstituted or substituted fused bicyclic aromatic heterocyclic group having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen;
- RY is -CH₂-, -O-, -S-, or -NH-; and
- R^Z is a bond or, when taken with RY and the three adjoining carbon atoms, forms a saturated carbocyclic ring of 5-8 atoms, one atom of which may be -O-, -S-, or -NH-;
- r is 1, 2 or 3; and
- G is -(CH₂)_s-R, where s is 0-5, -CH=CH-(CH₂)_t-R, where t is 0-3, or G is



216



where D and E are each independently N or CH;


k is 0 or 1;

b is 0 or 1;


5

M is S, O, or NH;

each W is independently N or CH; and




10



and wherein one to all of the otherwise unsubstituted carbon atoms of the aromatic or heteroaromatic rings of

15



may bear a fluoro substituent;

or a pharmaceutically acceptable salt thereof;


or a pharmaceutically acceptable solvate of said compound

20

or salt thereof;

provided that A is not hydrogen or

t-butyloxycarbonyl when G is $-(CH_2)_s-NH-C(NH)NH_2$, Y is



unsubstituted prolinyl (R^P is hydrogen), and T is


and provided further that R is not amino or

25

guanidino when $r=1$ and $s=0$;

and provided further that A is not hydrogen, C₁-C₄ alkyl, methylsulfonyl or -(CH₂)_q-R^m when G is


$$-(CH_2)_5-R \text{ in which R is } \begin{array}{c} \text{NH} \\ \parallel \\ -C-NH_2 \end{array} \text{ or } \begin{array}{c} \text{NH} \\ \parallel \\ -NH-C-NH_2 \end{array},$$

5 Y is  , unsubstituted prolinyl
(R^p is hydrogen) or 4-hydroxyprolinyl (R^p is OH), R' is
hydrogen, T is cyclohexyl and Q is -NH-A;


and provided further that R"SO₂ is not arylsulfonyl

when G is $-(CH_2)_5-R$ in which R is $-NH-\overset{NH}{\underset{||}{C}}-NH_2$, Y is unsubstituted prolinyl (R^P is hydrogen) or 4-methylthioprolinyl (R^P is $-SCH_3$), and Q is $-NH-A$;

and provided further that A is not $R^{\infty}SO_2^-$ when G is



' T is C₁-C₈ alkyl,




, and Q is -NH-A.


2. A compound of Formula I or salt or solvate thereof as claimed in Claim 1 wherein

X is prolinyl, homoprolinyl,

218



T is C₃-C₈ cycloalkyl, C₁-C₈ alkyl,



5

a is 0 or 1;

Q is -OH, C₁-C₄ alkoxy, or -NH-A;

A is hydrogen, C₁-C₄ alkyl, R^{''}SO₂-, R^{''}OC(O)-, R^{''}C(O)-, or -(CH₂)_g-COOH;

g is 1, 2, or 3;

10

B is hydrogen or C₁-C₄ alkyl;

R' is hydrogen or C₁-C₄ alkyl;

R^{''} is C₁-C₄ alkyl, C₁-C₄ perfluoroalkyl,

-(CH₂)_d-COOH, or unsubstituted or substituted aryl, where aryl

is phenyl, naphthyl, a 5- or 6-membered unsubstituted or

15

substituted aromatic heterocyclic ring, having one or two

heteroatoms which are the same or different and which are

selected from sulfur, oxygen and nitrogen, or a 9- or 10-

membered unsubstituted or substituted fused bicyclic aromatic

heterocyclic group having one or two heteroatoms which are

20

the same or different and which are selected from sulfur,



oxygen and nitrogen;

d is 1, 2, or 3;

m is 0, 1, or 2;

n is 0, 1, or 2; and

Z is hydrogen, C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxy, halo, or R_aSO₂NH-, where R_a is C₁-C₄ alkyl;



5 in which

R^g is C₁-C₆ alkyl; C₃-C₈ cycloalkyl, or $\text{-(CH}_2\text{)}_p\text{-L-(CH}_2\text{)}_q\text{-T'}$;

R^p is hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, or $\text{-(CH}_2\text{)}_p\text{-L-(CH}_2\text{)}_q\text{-T'}$;

10 where p is 0, 1, 2, 3, or 4; L is a bond, -O-, -S-, or -NH-; q is 0, 1, 2 or 3, and T' is hydrogen, C₁-C₄ alkyl, C₃-C₈ cycloalkyl, -COOH, -CONH₂, or Ar, where Ar is
 15 unsubstituted or substituted aryl, where aryl is phenyl, naphthyl, a 5- or 6-membered unsubstituted or substituted aromatic heterocyclic ring, having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen, or a 9- or 10-membered
 20 unsubstituted or substituted fused bicyclic aromatic heterocyclic group having one or two heteroatoms which are the same or different and which are selected from sulfur, oxygen and nitrogen;


R^y is -CH₂-, -O-, -S-, or -NH-; and

R^z is a bond or, when taken with R^y and the three adjoining carbon atoms, forms a saturated carbocyclic ring of
 25 5-8 atoms, one atom of which may be -O-, -S-, or -NH-;

r is 1 or 2; and

220

G is $-(CH_2)_s-R$, where s is 0-5, $-CH=CH-(CH_2)_t-R$,



- 5
 10
 or a pharmaceutically acceptable salt thereof;
 or a pharmaceutically acceptable solvate of said compound
 or salt thereof;

provided that A is not hydrogen or
 t-butyloxycarbonyl when G is $-(CH_2)_s-NH-C(NH)NH_2$, Y is

- 15
 unsubstituted prolinyl (R^P is hydrogen), and T is ;
 and provided further that R is not amino or
 guanidino when $r=1$ and $s=0$;

and provided further that A is not hydrogen, C_1 - C_4
 alkyl, methylsulfonyl or $-(CH_2)_g-COOH$ when G is $-(CH_2)_s-R$ in

- 20
 which R is $-C(=NH)-NH_2$ or $-NH-C(=NH)-NH_2$,

Y is , unsubstituted prolinyl
 (R^P is hydrogen) or 4-hydroxyprolinyl (R^P is OH), R' is
 hydrogen, T is cyclohexyl and Q is $-NH-A$;


and provided further that $R''SO_2$ is not arylsulfonyl when

- 25
 G is $-(CH_2)_s-R$ in which R is $-NH-C(=NH)-NH_2$, Y is unsubstituted


221

prolinyl (RP is hydrogen) or 4-methyl-thioprolinyl (RP is -SCH₃), and Q is -NH-A;

and provided further that A is not R"SO₂- when



5



, and Q is -NH-A.

3. A compound or salt or solvate thereof as claimed in Claim 1 or 2 wherein

10 alkyl by itself or as part of another substituent is methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl or sec-butyl;

perfluoroalkyl by itself or as part of another substituent is trifluoromethyl, perfluoroethyl, perfluoro-n-propyl, perfluoroisopropyl, perfluoro-n-butyl, perfluoro-t-butyl, perfluoroisobutyl or perfluoro-sec-butyl;

15 C₃-C₈ cycloalkyl is cyclopropyl, methylcyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 4-methylcyclohexyl or cyclooctyl;

20 halo is chloro, fluoro, bromo or iodo;


a 5- or 6-membered heterocyclic ring is furyl, thienyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyranyl, pyridinyl, pyrimidinyl, pyrazinyl, oxazinyl or thiazinyl;

25 a 9- or 10-membered heterocyclic ring is indolyl, benzothienyl, benzofuryl, benzoxazolyl, benzoisoxazolyl, benzopyrazolyl, quinolinyl, isoquinolinyl, benzimidazolyl or benzothiazolyl;

and further where any of the aromatic or
30 heteroaromatic groups listed for the definition of Ar or R" is independently unsubstituted or substituted with one or two substituents that will afford a stable structure

independently selected from halo, hydroxyl, C₁-C₄ alkyl, C₁-C₄ alkoxy, amino (-NH₂), mono(C₁-C₄ alkyl)amino, -(CH₂)_jCOOH, mercapto, -S(O)_h(C₁-C₄ alkyl), -NHS(O)_h(C₁-C₄ alkyl), -NHC(O)(C₁-C₄ alkyl), -S(O)_hNH₂, -S(O)_hNH(C₁-C₄ alkyl), or -S(O)_hN(C₁-C₄ alkyl)₂, h is 0, 1 or 2, and j is 0, 1, 2, 3, or 4.

4. A compound or salt or solvate thereof as claimed in Claim 1, 2 or 3 wherein




10 X is Z , homoprolinyl, 1- or 3-Tiq, or 1- or 3-Piq; Y is prolinyl; and Q is NHA in which A is hydrogen or RⁿSO₂-, R' is hydrogen, Z is hydrogen, and B is hydrogen; and R is a guanidino or an amidino group.

15 5. A compound or salt or solvate thereof as claimed in any one of Claims 1-4 wherein

G is a 4-amidinophenyl group.

20 6. A compound or salt or solvate thereof as claimed in Claim 1, 2 or 3 wherein



X is Q in which T is cyclohexyl, a is 1, R' is hydrogen and Q is -NH-A in which A is hydrogen, RⁿSO₂- or -(CH₂)_g-COOH.

25 7. A compound or salt or solvate thereof as claimed in Claim 6 in which A is RⁿSO₂- and Rⁿ is ethyl.


8. A compound or salt or solvate thereof as claimed in Claim 6 in which A is -(CH₂)_g-COOH and g is 1.

cyclohexylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide.

16. A compound or salt or solvate thereof as claimed in Claim 2 which compound is N-(carboxymethyl)-D-cyclohexylalanyl-N-[[5-(aminoiminomethyl)thiophen-2-yl]methyl]-L-prolinamide.


17. A pharmaceutical formulation comprising in association with a pharmaceutically acceptable carrier, diluent, or excipient, a compound of Formula I, Ia, Ib or Ic, or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of Claims 1-16.

18. A process for preparing a compound having the Formula I



as claimed in any one of Claims 1-9 which comprises

a) removing simultaneously or sequentially the protecting group(s) P of a corresponding compound of Formula II



wherein (P)X represents a radical X which may bear one or more protecting groups P independently selected from an amino acid protecting group P for a compound of Formula I in which X includes a basic NH moiety and a carboxy protecting group P for a compound of Formula I in which X includes a carboxy residue and G(P) represents a radical G which may bear one or more independently selected amino protecting groups P; or

b) for a compound of Formula I in which R is



, by hydrogenolysis of a corresponding compound of



Formula I in which R is $-\text{C}-\text{NH}_2$;

and whereafter, when a salt of the compound of Formula I is required, forming the salt with a pharmaceutically acceptable acid.


- 5 19. A method of inhibiting thrombin in a mammal comprising administering an effective dose of a compound, or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of Claims 1-16 to a mammal requiring thrombin inhibition.

9. A compound or salt or solvate thereof as claimed in any one of Claims 1-3 or 6-8 in which Y is (L)-prolinyl, (S)-cis-octahydro-1H-indole-2-carbonyl, or N-(2-phenylethyl)glycyl.

5

10. A compound, or a pharmaceutically acceptable salt or solvate thereof, which compound is selected from


a) a compound of Formula Ia



wherein the benzamidine ring is unsubstituted or may bear one or two fluoro substituents,


15

b) a compound of Formula Ib



and

c) a compound of Formula Ic




wherein D is N or CH;

and X has any of the values claimed in Claims 1-4 and 6-8.

11. A compound, or salt or solvate thereof, as claimed in Claim 10 which is a compound of Formula Ia in which the benzamidine ring is unsubstituted.

12. A compound, or salt or solvate thereof, as claimed in Claim 10 or 11 wherein



X is in which R' is hydrogen, a is 1, T is cyclohexyl or phenyl and Q is -NH-A in which A is hydrogen, ethylsulfonyl or carboxymethyl.

13. A compound or salt or solvate thereof as claimed in Claim 1 which compound is selected from

- a) D-phenylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
- b) N-[[4-(aminoiminomethyl)phenyl]methyl]-1-[[4aS,8aS]-decahydro-1(R)-isoquinolinyl]carbonyl]-L-prolinamide,
- c) N-(ethylsulfonyl)-D-phenylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,

- d) (S-cis)-N-[[4-(aminoiminomethyl)-phenyl]methyl]-1-[N-(ethylsulfonyl)-D-phenylglycyl]-1H-indole-2-carboxamide,
- e) (S-cis)-N-[[4-(aminoiminomethyl)-phenyl]methyl]-1-[N-(ethylsulfonyl)-D-phenylalanyl]-1H-indole-2-carboxamide,
- 5 f) N-(carboxymethyl)-D-phenylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
- g) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
- 10 h) (S-cis)-N-[[4-(aminoiminomethyl)-phenyl]methyl]-1-[N-(carboxymethyl)-D-cyclohexylalanyl]-1H-indole-2-carboxamide,
- i) D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
- 15 j) N-(2-carboxyethyl)-D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
- k) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[1-(aminoiminomethyl)-hexahydropyridin-4-yl]methyl]-L-prolinamide,
- 20 l) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[5-(aminoiminomethyl)thiophen-2-yl]methyl]-L-prolinamide,
- m) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[5-(aminoiminomethyl)pyridin-2-yl]methyl]-L-prolinamide,
- 25 n) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[5-(aminoiminomethyl)-1,2,3,4-tetrahydropyridin-2-yl]methyl]-L-prolinamide,
- o) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[6-(aminoiminomethyl)pyridazin-3-yl]methyl]-L-prolinamide,
- 30 p) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[1-(aminoiminomethyl)-1,2,3,6-tetrahydropyridin-4-yl]methyl]-L-prolinamide,
- q) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)-2-fluorophenyl]methyl]-L-prolinamide,
- 35 r) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[4-(aminoiminomethyl)-2,6-difluorophenyl]methyl]-L-prolinamide,

- s) N-(ethylsulfonyl)-D-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
t) N-(carboxymethyl)-D-cyclohexylglycyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
5 u) N-(carboxymethyl)-D-homophenylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
v) N-(carboxymethyl)-D-homocyclohexylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide, and
w) N-(methylsulfonylacetyl)-L-cyclohexylalanyl-N-
10 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide.

14. A compound or salt or solvate thereof as
claimed in Claim 1 which compound is selected from

- i) (S-cis)-N-[[4-(aminoiminomethyl)-
15 phenyl]methyl]-1-[N-(ethylsulfonyl)-D-phenylalanyl]-1H-
indole-2-carboxamide,
ii) N-(carboxymethyl)-D-phenylalanyl-N-
[[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
iii) N-(carboxymethyl)-D-cyclohexylalanyl-N-
20 [[4-(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
iv) D-cyclohexylalanyl-N-[[4-(
(aminoiminomethyl)phenyl]methyl]-L-prolinamide,
v) N-(carboxymethyl)-D-cyclohexylalanyl-N-
[[5-(aminoiminomethyl)thiophen-2-yl]methyl]-L-prolinamide,
25 vi) N-(carboxymethyl)-D-cyclohexylalanyl-N-[[1-(
(aminoiminomethyl)-1,2,3,6-tetrahydropyridin-4-yl]methyl]-L-
prolinamide,
vii) N-(carboxymethyl)-D-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)-2-fluorophenyl]methyl]-L-prolinamide,
30 and
vii) N-(carboxymethyl)-D-cyclohexylalanyl-N-
[[4-(aminoiminomethyl)-2,6-difluorophenyl]methyl]-L-
prolinamide.

- 35 15. A compound or salt or solvate thereof as
claimed in Claim 2 which compound is N-(carboxymethyl)-D-

INTERNATIONAL SEARCH REPORT

Int. application No.
PCT/US95/02558

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : Please See Extra Sheet.

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

USPTO APS

search terms: smith, wiley, agmatine, antithrombo?

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BE, 887,224 (BAJUSZ ET AL) 27 July 1981, see pages 3 and 6.	1-4, 17-19
X	US, A, 4,346,078 (BAJUSZ ET AL) 24 August 1982, see col. 2, lines 60+ and col. 3, lines 20+.	1-4, 17-19

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* "A"	document defining the general state of the art which is not considered to be of particular relevance	* "T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
* "E"	earlier document published on or after the international filing date	* "X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
* "L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	* "Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
* "O"	document referring to an oral disclosure, use, exhibition or other means	* "Z"	document member of the same patent family
* "P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

07 APRIL 1995

Date of mailing of the international search report

08 MAY 1995

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SHEELA J. HUFF

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/02558

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 38/00; C07D 205/08, 223/16, 239/02, 211/26, 285/08, 285/14, 417/00, 513/00, 277/04, 277/18, 277/38, 263/04, 263/16, 295/00, 207/00, 207/08, 207/10; C07C 229/00, 241/00, 243/00, 249/00, 233/00, 235/00, 237/00, 239/00

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

540/362, 593, 594, 595; 544/323; 546/199, 246; 548/128, 131, 190, 214, 233, 566, 518, 558, 326.5, 453; 560/34, 35, 168, 169; 562/126, 439, 560; 564/123; 514/19

B. FIELDS SEARCHED

Minimum documentation searched

Classification System: U.S.

540/362, 593, 594, 595; 544/323; 546/199, 246; 548/128, 131, 190, 214, 233, 566, 518, 558, 326.5, 453; 560/34, 35, 168, 169; 562/126, 439, 560; 564/123; 514/19